




UNIVERSITY
OF TURKU

A large, high-magnification microscopic image of a plant structure, possibly a flower or leaf, rendered in a cyan color against a black background. The structure is complex and multi-lobed, with a central point from which several arms or petals radiate outwards.

ON SINGLET OXYGEN,
PHOTOINHIBITION,
PLASTOQUINONE
AND THEIR
INTERCONNECTIONS

Heta Mattila

TURUN YLIOPISTON JULKAISUJA – ANNALES UNIVERSITATIS TURKUENSIS

SARJA – SER. AI OSA – TOM. 634 | ASTRONOMICA – CHEMICA – PHYSICA – MATHEMATICA | TURKU 2020



UNIVERSITY
OF TURKU

**ON SINGLET OXYGEN,
PHOTOINHIBITION,
PLASTOQUINONE
AND THEIR
INTERCONNECTIONS**

Heta Mattila

University of Turku

Faculty of Science and Engineering
Department of Biochemistry
Molecular Plant Biology unit
Doctoral Programme of Molecular Life Sciences

Supervised by

Adjunct Professor, Esa Tyystjärvi
University of Turku
Faculty of Science and Engineering
Department of Biochemistry
Molecular Plant Biology unit

Reviewed by

Assistant Professor, Alizée Malnoë
University of Umeå
Sweden

Professor, Shunichi Takahashi
National Institute of Basic Biology
Okazaki, Japan

Opponent

Associate Professor, Pavel Pospíšil
Palacký University Olomouc
Czech Republic

The originality of this publication has been checked in accordance with the University of Turku quality assurance system using the Turnitin OriginalityCheck service.

Cover Image: Heta Mattila

ISBN 978-951-29-8269-1 (PRINT)
ISBN 978-951-29-8270-7 (PDF)
ISSN 0082-7002 (Print)
ISSN 2343-3175 (Online)
Painosalama, Turku, Finland 2020

--Tämä heleä, kostea multa,
tämä kirkkaus loppumaton--

Saima Harmaja

UNIVERSITY OF TURKU

Faculty of Science and Engineering

Department of Biochemistry

Molecular Plant Biology

HETA MATTILA: On singlet oxygen, photoinhibition, plastoquinone, and their interconnections

Doctoral Dissertation, 151 pp.

Doctoral Programme of Molecular Life Sciences

November 2020

ABSTRACT

In oxygenic photosynthesis light is captured in series by two protein complexes: photosystem II and photosystem I (PSI and PSII). Electron transfer from PSII to PSI is mediated by the plastoquinone (PQ) pool. Despite being the energy source, light also damages the photosynthetic machinery. Singlet oxygen ($^1\text{O}_2$), an excited state of O_2 , may be generated when a charge recombination reaction in PSII re-excites the reaction center chlorophylls (P680), producing a triplet state. In this thesis, mass-spectroscopy-based detection methods for $^1\text{O}_2$ were developed further, to understand the role of this reactive oxygen species in photosynthesis. It was shown that even though both O_2 and $^1\text{O}_2$ are produced in pumpkin thylakoid membranes, most (if not all) of the $^1\text{O}_2$ derives from the ambient dissolved O_2 , not from the nascent O_2 produced due to the water splitting activity of PSII. The result shows that the O_2 evolving ability of PSII as such does not render PSII vulnerable to oxidative damage.

Nevertheless, light inactivates PSII, and no consensus about the mechanism(s) of the photoinhibitory damage exists. Therefore, the temperature dependence of the rate constant of PSII photoinhibition was measured under various conditions and compared with temperature dependencies of $^1\text{O}_2$ production and recombinations. The results show that in plants and cyanobacteria the rate constant of photoinhibition and production of $^1\text{O}_2$ increase similarly with temperature as the miss probability of the oxygen evolving complex. Photoinhibition proceeded under anaerobicity, where no $^1\text{O}_2$ is produced, and was unaffected by quenchers of $^1\text{O}_2$. We suggest that when a miss occurs, but a recombination does not re-reduce the PSII reaction center, P680⁺ lives long enough to oxidize a vital component of PSII, causing the photodamage.

Plants have also ways to adjust to different light conditions. An initial fluorescence screen and subsequent high-performance liquid chromatography measurements revealed that 470 nm, 560 nm and 660 nm light favors PSII over PSI and reduces 80–90 % of the PQ pool, whereas 440 nm, 520 nm and 690 nm favors PSI and oxidizes 90–100 % of the pool in *Arabidopsis thaliana*, when moderate light was used. Light state curvilinearly followed the redox state of the PQ pool; state 2 was reached with 50 % reduction. All tested white lights, including light from the Sun, reduced less than 50 % of the PQ pool. This PSI light character of white light enables plants to respond to the intensity of light via the redox state of the PQ pool.

KEYWORDS: MIMS, photoinactivation, ROS, signaling, state transitions, TEMP

TURUN YLIOPISTO

Luonnontieteiden ja tekniikan tiedekunta

Biokemian laitos

Molekulaarinen kasvibiologia

Heta Mattila: On singlet oxygen, photoinhibition, plastoquinone, and their interconnections

Väitöskirja, 151 s.

Molekulaaristen biotieteiden tohtoriohjelma

marraskuu 2020

TIIVISTELMÄ

Vaikka valo onkin yhteytyksen eli fotosynteesin energianlähde, se myös vahingoittaa yhteyttäviä eliöitä. Hapetta vapauttavassa yhteytyksessä kaksi proteiini-kompleksia, valoreaktiot II ja I (PSII ja PSI), vastaanottavat valoa. Plastokinonivaranto välittää elektroninsiirtoa näiden välillä. Singlettihappi on reaktiivinen happiyhdiste, jota syntyy, kun PSII:n reaktiokeskus-klorofylli (P680) siirtyy triplettilaan elektronin palatessa rekombinaatioreaktioissa takaisin reaktiokeskukselle. Tässä työssä kehitettiin edelleen massaspektrometria-menettelmiä ja havaittiin, että vaikka eristetyt kasvin yhteytyskalvostot tuottavat sekä happea että singlettihapetta, suurin osa singlettihapetta on peräisin ympäristöön liuenneesta hapesta eikä PSII:n veden hajotuksessa vastasyntyneestä hapesta. PSII:n kyky vapauttaa happea ei siis itsessään tee PSII:sta herkkää hapettavalle vahingolle.

Valo kuitenkin vaurioittaa erityisesti PSII:ta, eikä valovahingon mekanismista ole yksimielisyyttä. Tästä syystä PSII:n fotoinhibition, singlettihapen tuoton ja rekombinaatioreaktioiden lämpötilavasteet mitattiin kasveista ja syanobakteereista. Sekä fotoinhibitio että singlettihappi lisääntyivät lämpötilan noustessa. Molemmat lämpötilavasteet selvästi muistuttavat PSII:n happea vapauttavan kompleksin "hutin" lämpötilariippuvuutta. Fotoinhibitio ei kuitenkaan vähentynyt, kun yhteytyskalvostoja valotettiin singlettihapen vaimentajien kanssa tai hapettomissa olosuhteissa, jolloin singlettihapetta ei synny. Tulosten perusteella muotoiltiin hypoteesi: mikäli huti tapahtuu eli happea vapauttava kompleksi on tilapäisesti kykenemätön luovuttamaan elektronin, eikä rekombinaatiokaan uudelleen pelkistä reaktiokeskusta, P680⁺:lla on tarpeeksi aikaa hapettaa jokin PSII:n tärkeä osa.

Kasvit kykenevät myös sopeutumaan valo-olosuhteiden muutoksiin. Plastokinonivarannon hapetus-pelkistystila mitattiin nestekromatografialla; PSII vastaanotti PSI:tä tehokkaammin 470, 560 ja 660 nm:n aallonpituuksien valoa ja nämä valot pelkistivät 80–90 % plastokinonivarannosta. 440, 520 ja 690 nm:n aallonpituudet sen sijaan suosivat PSI:tä ja hapettivat 90–100 % plastokinonivarannosta, kun kohtalaista valovoimakkuutta käytettiin. Kaikki testatut valkoiset valot, mukaan lukien auringonvalo, suosivat PSI:tä, joten kasvit voivat aistia valon voimakkuutta plastokinonivarannon pelkistyneisyyden avulla.

ASIASANAT: HPLC, MIMS, plastokinoni, reaktiiviset happilajit, signalointi, ¹O₂, TEMP, tilasiirtymät, yhteyttäminen

Table of contents

Table of contents	6
Abbreviations	8
List of original publications	11
1 Introduction	12
1.1 Photosynthetic electron transport chain.....	12
1.1.1 Energy transfer: (not simply) hop-hop-hop-hop-hop	13
1.1.2 Structure and function of photosystem II	14
1.1.3 The mediators: plastoquinone pool, cytochrome <i>b₆/f</i> complex and plastocyanin	16
1.1.4 Structure and function of photosystem I	17
1.2 Singlet oxygen	18
1.2.1 Very shortly about the chemistry of singlet oxygen.....	18
1.2.2 Singlet oxygen production by chlorophylls.....	19
1.2.3 Antenna or reaction center?	20
1.2.4 Additional sources of singlet oxygen in plant cells	21
1.2.5 Non-photochemical quenching and other protective measures against singlet oxygen	22
1.2.6 Where, when and why to bother?	23
1.3 Photoinhibition	25
1.3.1 Photodamage to photosystem II in ultraviolet radiation	26
1.3.2 How does visible light damage photosystem II?	26
1.3.3 The repair cycle.....	28
1.3.4 Photoinhibition of photosystem I.....	29
1.4 Responses to changing quantity and quality of light.....	30
1.4.1 Balancing the photosystems: roles for plastoquinone redox state	30
1.4.2 Photosynthetic control	32
1.4.3 Cyclic pathways and other alternative routes	32
2 Aims of the study	34
3 Materials and methods	35
3.1 Organisms and growth conditions	35
3.1.1 Light sources.....	35
3.2 Chlorophyll <i>a</i> fluorescence and P700 signal.....	36
3.3 Quantification of photoinhibition	36
3.4 Measurements of singlet oxygen.....	37

3.5	Measurements of plastoquinone.....	37
4	Overview of the results.....	38
4.1	Mass spectroscopy could be used to quantify singlet oxygen..	38
4.2	Nascent oxygen produced by photosystem II was not converted to singlet oxygen	38
4.3	Photoinhibition of photosystem II.....	40
4.3.1	S ₂ Q _A ⁻ recombination reactions cannot be the reason for photoinhibition caused by visible light or laser pulses..	40
4.3.2	Singlet oxygen did not affect photoinhibition.....	41
4.3.3	Photoinhibition proceeded even at -78.5 °C.....	42
4.4	Redox state of the plastoquinone pool.....	42
4.4.1	F ₀ rise revealed visible light wavelengths favoring photosystem II or photosystem I.....	42
4.4.2	The photochemically active plastoquinone pool was reduced by three and oxidized by four visible light wavelengths	44
4.4.3	The lights favoring photosystem II or photosystem I induced related physiological changes	44
4.4.4	Redox state of the plastoquinone pool in white light	45
4.4.5	Validation of the liquid chromatography method	46
5	Discussion.....	47
5.1	Improved detection methods for singlet oxygen are needed...	47
5.2	Singlet oxygen and photoinhibition	47
5.2.1	Photosystem II is protected against nascent oxygen ...	47
5.2.2	Photoinhibition and misses of oxygen evolving complex: an unexpected resemblance.....	49
5.2.3	Long-lived P680 ⁺ damages photosystem II.....	49
5.2.4	Direct light absorption of the manganese ions contributes to photoinhibition also in visible light.....	50
5.2.5	Why do carotenoids protect photosystem II?	52
5.2.6	Sustained quenching of fluorescence may affect photoinhibition measurements.....	53
5.3	Redox state of the plastoquinone pool senses both the imbalances between the photosystems and the intensity of sunlight.....	54
5.3.1	Delicate balance between the photosystems.....	54
5.3.2	Implications for state transitions	55
5.3.3	Redox state of the plastoquinone pool can and should be measured	57
6	Conclusions and future perspectives	60
	Acknowledgements	61
	List of references.....	62
	Original Publications.....	87

Abbreviations

^1Chl	Singlet (ground state) chlorophyll
$^1\text{O}_2$	Singlet oxygen; $^1\Delta_g\text{O}_2$
$^1\Delta_g\text{O}_2$	Singlet oxygen
$^1\Sigma^+_g\text{O}_2$	Singlet oxygen
^3Chl	Triplet chlorophyll
$^3\text{O}_2$	Triplet (ground state) oxygen; $^3\Sigma^-_g\text{O}_2$
$^3\text{P680}$	PSII reaction center triplet ($^3\text{Chl}_{\text{D1}}$ or $^3\text{P}_{\text{D1}}$)
$^3\text{P700}$	PSI reaction center triplet
$^3\Sigma^-_g\text{O}_2$	Triplet (ground state) oxygen; $^3\text{O}_2$
^{16}O	A stable isotope of oxygen (common in the nature)
^{18}O	A stable isotope of oxygen (rare in the nature)
A_0	The primary electron acceptor of PSI (a chlorophyll)
ADP	Adenosine diphosphate
ATP	Adenosine triphosphate
ATPase	ATP synthase
Car	Carotenoid
Car_{D2}	A β -carotene close to PSII reaction center
Chl	Chlorophyll
Chl^*	Excited singlet chlorophyll
Chl_{D1}	A PSII reaction center chlorophyll
Chl_{D2}	A PSII reaction center chlorophyll
ChlZ_{D2}	A chlorophyll close to PSII reaction center
CP24	An antenna protein of PSII
CP26	An antenna protein of PSII
CP29	An antenna protein of PSII
CP43	A core protein of PSII (intrinsic antenna)
CP47	A core protein of PSII (intrinsic antenna)
Cyt b6f	Cytochrome b ₆ /f complex
D1	A core protein of PSII
D2	A core protein of PSII
D_2O	Deuterium oxide (heavy water); $^2\text{H}_2\text{O}$

DCMU	3-(3,4-dichlorophenyl)-1,1-dimethylurea (a PSII herbicide)
Deg1	A protease important in PSII repair
EF-G	A cyanobacterial translation elongation factor
EF-Tu	A cyanobacterial translation elongation factor
EPR	Electron paramagnetic resonance
F ₀	Minimal fluorescence yield (Q _A oxidized), dark-acclimated sample
F ₀ '	Minimal fluorescence yield (Q _A oxidized), light-acclimated sample
F _A	A 4Fe-4S cluster of PSI
F _B	A 4Fe-4S cluster of PSI
Fd	Ferredoxin
F _M	Maximal fluorescence yield (Q _A reduced), dark-acclimated sample
F _M '	Maximal fluorescence yield (Q _A reduced), light-acclimated sample
FNR	Ferredoxin NADP ⁺ reductase
FR	Far-red ($\lambda \geq 680$ nm)
FtsH	A protease important in PSII repair
F _V	Variable fluorescence (F _M -F ₀)
F _V /F _M	Variable to maximal fluorescence (yield of PSII photochemistry)
F _X	A 4Fe-4S cluster of PSI
FWHM	Full width at half maximum
H ₂ O ₂	Hydrogen peroxide
HO•	Hydroxyl radical
k _{PI}	Rate constant of photoinhibition of PSII
LED	Light emitting diode
LHC	Light harvesting complex
LHCI	Light harvesting complex of PSI
LHCII	Light harvesting complex of PSII
LHCSR	A protein important in NPQ in algae and moss
MIMS	Membrane inlet mass spectroscopy
MS	Mass spectrometer/spectroscopy
NADP ⁺	Nicotinamide adenine dinucleotide phosphate (oxidized)
NADPH	Nicotinamide adenine dinucleotide phosphate (reduced)
NIR	Near-infrared (~750–2500 nm)
NO•	Nitric oxide (a radical)
NPQ	Non-photochemical quenching (of chlorophyll fluorescence)
O ₂ • ⁻	Superoxide anion (a radical)
OCP	Orange carotenoid protein (important in cyanobacterial NPQ)
OEC	Oxygen evolving complex of PSII
OJIP	Fluorescence rise from initial (O) to maximal (P) level, via J and I steps
P680	Reaction center chlorophylls of PSII (P _{D1} , P _{D2} , Chl _{D1} and Chl _{D2})
P680 ⁺	Oxidized reaction center of PSII (P _{D1} ⁺)

P700	Reaction center chlorophylls of PSI
PAM	Pulse amplitude modulated
PBCP	PSII core phosphatase
PC	Plastocyanin
P _{D1}	A PSII reaction center chlorophyll
P _{D2}	A PSII reaction center chlorophyll
PGR5	A protein important for PSI cyclic electron transfer
PGRL1	A protein important for PSI cyclic electron transfer
Pheo	Pheophytin
Pheo _{D1}	Redox active pheophytin of PSII
Pheo _{D2}	A pheophytin close to PSII reaction center
PhQ	Phylloquinone
Pi	Inorganic phosphate
PPFD	Photosynthetic photon flux density
PQ	Oxidized plastoquinone
PQH ₂	Reduced plastoquinone; plastoquinol
PsbS	A protein important in NPQ in higher plants
PsaA	A core protein of PSI
PsaB	A core protein of PSI
PSI	Photosystem I
PSII	Photosystem II
Q _A	Primary quinone acceptor of PSII
Q _B	Secondary quinone acceptor of PSII
qL	Photochemical quenching of fluorescence (lake model)
qE	Energy-dependent quenching of excitation energy
qP	Photochemical quenching of fluorescence (puddle model)
ROS	Reactive oxygen species
SD	Standard deviation
SOSG	Singlet oxygen sensor green (a ¹ O ₂ sensor)
STN7	State transition kinase (STT7 in <i>C. reinhardtii</i>)
STN8	A kinase involved in phosphorylating PSII core
TAP38	A phosphatase involved in state transitions (PPH1 in <i>C. reinhardtii</i>)
TEMP	2,2,6,6-tetramethylpiperidine (a ¹ O ₂ sensor)
TEMPO	1-oxyl of TEMP (reaction product of TEMP and ¹ O ₂)
TyrZ	Redox active tyrosine residue of PSII; Y _Z
(U)HPLC	(Ultra) high-performance liquid chromatography
UV-A	Ultraviolet radiation (315–400 nm)
UV-B	Ultraviolet radiation (280–315 nm)
UV-C	Ultraviolet radiation (200–280 nm)
VIS	Visible light (~400–700 nm)

List of original publications

This dissertation is based on the following original publications, which are referred to in the text by their Roman numerals:

- I Maarit Karonen*, Heta Mattila*, Ping Huang, Fikret Mamedov, Stenbjörn Styring, Esa Tyystjärvi. A tandem mass spectrometric method for singlet oxygen measurement. *Photochemistry and Photobiology*, 2014; 90: 965–971.
- II Heta Mattila, Sergey Khorobrykh, Esa Tyystjärvi. Singlet oxygen derives from ambient dissolved gas, not from nascent oxygen produced by Photosystem II. Manuscript.
- III Heta Mattila, Sujata Mishra, Taina Tyystjärvi, Esa Tyystjärvi. Misses of oxygen evolving complex play a crucial role in photoinhibition of Photosystem II. Manuscript.
- IV Heta Mattila*, Sergey Khorobrykh*, Marja Hakala-Yatkin, Vesa Havurinne, Iiris Kuusisto, Taras Antal, Taina Tyystjärvi, Esa Tyystjärvi. Action spectrum of the redox state of the plastoquinone pool defines its function in plant acclimation. *The Plant Journal*, 2020; doi: 10.1111/tpj.14983.

*Equal contribution.

The original publications have been reproduced with the permission of the copyright holders.

Other recent publications related to the topic:

Mattila H*, Mishra KB*, Kuusisto I, Mishra A, Novotná K, Šebela D, Tyystjärvi E (2020) Effects of low temperature on photoinhibition and singlet oxygen production in four natural accessions of *Arabidopsis*. *Planta* 252: 19

Khorobrykh S, Havurinne V, Mattila H, Tyystjärvi E (2020) Oxygen and ROS in photosynthesis. *Plants* 9: 91

Havurinne V, Mattila H, Antinluoma M, Tyystjärvi E (2019) Unresolved quenching mechanisms of chlorophyll fluorescence may invalidate multiple turnover saturating pulse analyses of photosynthetic electron transfer in microalgae. *Physiol Plant* 166: 365–379

Mattila H, Khorobrykh S, Havurinne V, Tyystjärvi E (2015) Reactive oxygen species: Reactions and detection from photosynthetic tissues. *J Photochem Photobiol B* 152: 176–214

1 Introduction

Some 3.7–2.4 billion years ago, a (pre)cyanobacterium, now extinct, achieved the ability to extract electrons from water with the energy of light (Fischer et al. 2016a, Martin et al. 2018; oxygenic photosynthesis might have evolved also much before the phylum Cyanobacteria, see e.g. Cardona et al. 2019). Light and water are plentiful on Earth, and the (eventual) rise in atmospheric oxygen, due to water splitting, enabled efficient respiration. Consequently, the emergence of oxygenic photosynthesis which, to the present knowledge, has occurred only once in the evolution of life, increased the primary production of Earth by a factor of ~30, as estimated by Raven (2009). Today, (oxygenic) photosynthesis supports almost all life on Earth (Mccollom and Shock 1997, Raven 2009). According to the widely accepted endosymbiosis hypothesis (Mereschkowsky 1905), chloroplasts of algae and plants were once free-living cyanobacteria but became engulfed into an ancient eukaryote and transformed into cell organelles (for a review and a recent phylogenetic analysis, respectively, see Gavelis and Gile 2018, Moore et al. 2019). Chloroplasts of all but one (*Paulinella chromatophora*; Marin et al. 2005, Delaye et al. 2016) photosynthetic eukaryotes known to science originate from the same primary endosymbiosis event.

1.1 Photosynthetic electron transport chain

Membrane embedded protein complexes and lipid- and water-soluble electron carriers of photosynthetic electron transport chain convert light energy to chemical form (Fig. 1); to the universal carriers of energy (adenosine triphosphate; ATP) and reducing power (nicotinamide adenine dinucleotide phosphate; NADPH). The thylakoid membrane system is, in the case of plants and algae, separated from the rest of the cell by the double membrane (called envelope) of chloroplasts. In land plants (and in some green macroalgae), thylakoids are organized to appressed, or stacked, (grana thylakoids) and non-appressed (stroma thylakoids) domains (see Fig. 1B). In the stroma (or, in the case of cyanobacteria, in the cytosol) ATP and NADPH are used by the Calvin-Benson cycle to fix atmospheric CO₂ into sugars.




Figure 1. Photosynthesis in plant chloroplasts. A. Light microscopy images showing cells and, B. electron microscopy images showing a chloroplast and few mitochondria, of an outdoors-grown dandelion (*Taraxacum officinale*). C. A schematic representation of photosynthetic electron transfer chain and ATP synthesis. Linear electron (e^-) transfer from water to NADP^+ , via photosystem II (PSII), plastoquinone (PQ) pool, cytochrome b_6/f complex (Cyt b_6f), plastocyanin (PC), photosystem I (PSI), ferredoxin (Fd) and Fd NADP^+ reductase (FNR), is indicated with a continuous arrow. Cyclic electron transfer from Fd to PQ pool is indicated with a dashed arrow. $h\nu$ (energy of a photon; h is the Planck's constant and ν is frequency) indicates absorption of light energy at PSII and PSI. ATP production from adenosine diphosphate (ADP) and inorganic phosphate (P_i) by ATP synthase (ATPase), powered by an electrochemical gradient over the thylakoid membrane created by water splitting and proton (H^+) translocation from stroma to lumen, is indicated by another continuous arrow. Here, a rate of 4 protons/an ATP produced is assumed, based on measurements by van Walraven et al. (1996). D. A general equation of oxygenic photosynthesis, assuming 100 % quantum efficiency. Microscopy images: Olli Virtanen, Molecular Plant Biology, and the Laboratory of electron microscopy, Faculty of medicine, University of Turku.

1.1.1 Energy transfer: (not simply) hop-hop-hop-hop-hop

In oxygenic photosynthesis, light energy is harvested by chlorophylls (Chl; all photosynthetic organisms) and phycobilins (cyanobacteria and some algae) and to a lesser degree by carotenoids (Hofmann et al. 1996, Caffarri et al. 2001, Balevičius

et al. 2017). To increase light absorption (cross-section), photosynthetic organisms have light harvesting complexes (LHC), or antennae, which contain a high number of tightly packed pigment molecules. After absorption of visible light (a photon with wavelength of ~400–700 nm) by a pigment, fast non-radiative excitation energy transfer to the respective reaction center (of either PSII or PSI) occurs (see e.g. Wientjes et al. 2011, Kreisbeck and Aspuru-Guzik 2016). It is not clear whether the rate limiting step is charge separation in the reaction center (trap limited model) or energy transfer to the reaction center, possibly after initial fast equilibrium in the antenna (transfer-to-trap limited model), in the membrane embedded chlorophyll binding antennae of plants and some algae (for discussion, see Renger and Schlodder 2011, Croce and van Amerongen 2013). Excitation transfer from the large, membrane attached protein-pigment complex, phycobilisome, in cyanobacteria and some algae (for a review, see Watanabe and Ikeuchi 2013), to a PSII reaction center is rather slow (~100 ps; Suter et al. 1984). Often, excitation does not reside on a single pigment molecule, but electronic interactions of pigments close to each other lead to delocalized excitations (i.e. several pigment molecules coherently share the excitation), called excitons. Whether excitation transfer is sufficiently described by semi-classical theories (e.g. generalized Förster; Sumi 1999) or whether quantum effects significantly affect excitation transfer *in vivo*, is debated (Engel et al. 2007, Fassioli et al. 2014, Jumper et al. 2018).

1.1.2 Structure and function of photosystem II

PSII core consists of the reaction center proteins D1 and D2, which bind the redox cofactors needed for charge separation, of the inner (or intrinsic) antenna proteins CP47 and CP43, which bind 16 and 13 chlorophylls, respectively, and 3 carotenoids each, and of multiple smaller subunits (for a 1.9 Å X-ray structure, see Umena et al. 2011). In plants, PSII mostly resides in grana and grana margins (Andersson and Anderson 1980), functions as a dimer and is usually surrounded by two strongly and two moderately bound trimeric LHCs (consisting of Lhcb1–3 proteins in different combinations) and by monomeric antenna proteins (CP24, CP26 and CP29), two of each (Caffarri et al. 2009). Bigger complexes, involving also PSI, may exist *in vivo* (e.g. Järvi et al. 2011; for a review, see Rantala et al. 2020a).

P680, the reaction center of PSII, may be considered to consist of four chlorophyll *a* molecules (P_{D1} , P_{D2} , Chl_{D1} and Chl_{D2} ; where P stands for pigment and subscripts indicate whether the chlorophylls are bound to D1 or D2). Light energy transfer from LHCII (light harvesting antenna of PSII) or direct light absorption by P680 leads to its excitation, delocalized over at least two of the chlorophylls (Durrant et al. 1992) and maybe also at the primary electron acceptor of PSII (a pheophytin *a* molecule called Pheo_{D1}; Durrant et al. 1995, Holzwarth et al. 2006). Fast charge

separation follows (Fig. 2A–B). At least two pathways function; the primary electron donor is either Chl_{D1} or P_{D1} (Diner et al. 2001, Novoderezhkin et al. 2007, Romero et al. 2010, Renger and Schlodder 2011). Irrespective of the primary donor, the hole (P680^+) is localized mainly at P_{D1} (Diner et al. 2001, Nagao et al. 2017), though Pavlou et al. (2018) suggested that in far-red light it might reside at Chl_{D1} . Romero et al. (2017) reviewed the proposed charge separation pathways as follows (where δ denotes charge transfer character of excitations or excitons, indicated by *):

1. $(\text{Chl}_{D1}^{\delta+}\text{Pheo}_{D1}^{\delta-})^* \rightarrow \text{Chl}_{D1}^+\text{Pheo}_{D1}^- \rightarrow \text{P}_{D1}^+\text{Pheo}_{D1}^-$
2. $(\text{P}_{D1}\text{P}_{D2}\text{Chl}_{D1})^* \rightarrow (\text{P}_{D2}^{\delta+}\text{P}_{D1}^{\delta-})^* \rightarrow \text{P}_{D1}^+\text{Chl}_{D1}^- \rightarrow \text{P}_{D1}^+\text{Pheo}_{D1}^-$




Figure 2. Electron transfer in photosystem II. A. Redox cofactors of PSII. The atoms in the oxygen evolving complex (OEC) are colored as follows: calcium with green, oxygen red and manganese purple. B. A schematic representation of electron transfer routes in PSII. Forward electron transfer is indicated with black continuous arrows and recombination (back-reactions) with gray dashed arrows. The red continuous arrow highlights the recombination reaction that may lead to singlet oxygen ($^1\text{O}_2$) production. C. The Kok cycle of OEC (according to Klauss et al. 2012, Siegbahn 2013, Kern et al. 2018). S-states (S_{0-4}) indicate oxidation of the Mn ions. Time constants are from Meyer et al. (1989), Schatz et al. (1987), Takahashi et al. (1987), Vasil'ev et al. (1996), Hastings et al. (1992), de Wijn and van Gorkom (2001) and Groot et al. (2005).

After the primary charge separation, the charge is stabilized at Q_A (a PQ molecule tightly bound to the D2 protein). Q_A^- donates the electron to a second PQ molecule, Q_B (residing in the D1 protein). After subsequent light absorption and charge separation, another electron is donated to Q_B^- . Q_B^{2-} is reduced sequentially by two protons from the stroma (most probably before and after the second electron donation from Q_A^-) and then released into thylakoid membrane (described in detail in Guskov et al. 2009, Saito et al. 2013, Zobnina et al. 2017, Yao et al. 2018).

$P680^+$ extracts an electron from a tyrosine residue of PSII (TyrZ; also called Y_Z), producing the neutral radical TyrZ• (a proton is transferred to a nearby hydrogen-bonded histidine residue; Barry and Babcock 1987, Hays et al. 1998, Junge et al. 2002). TyrZ• oxidizes the water splitting and oxygen evolving Mn_4CaO_5 complex (OEC). In the Kok cycle (Kok et al. 1970; Fig. 2C), four electrons and four protons are extracted from two water molecules with the energy of four photons (i.e. four charge separations), leading to evolution of a molecule of dioxygen (O_2). In its most reduced state (S_0), three of the Mn ions of OEC are (most probably) Mn^{+3} and one is Mn^{+4} . Each flash of light further oxidizes a Mn ion ($Mn^{+3} \rightarrow Mn^{+4}$), advancing the S-state ($S_0 \rightarrow S_1 \rightarrow S_2 \rightarrow S_3$). After the fourth flash, a transient S_4 (possibly involving a Mn^{5+}) is produced, O_2 is spontaneously released and OEC returns to S_0 . Alternating electron donation to TyrZ• and proton release into lumen occur simultaneously with oxidation of the Mn ions (Fig. 2C; Krauss et al. 2012). However, the exact mechanism, specifically formation of the O-O bond, is still under discussion (see e.g. Siegbahn 2013, Kupitz et al. 2014, Kern et al. 2018, Suga et al. 2019).

1.1.3 The mediators: plastoquinone pool, cytochrome b_6/f complex and plastocyanin

The lipid-soluble plastoquinone (or plastoquinol) Q_BH_2 (hereafter PQH_2) is oxidized back to PQ in the bifurcated Q-cycle of the Cyt b_6f complex (Mitchell 1976). One electron from PQH_2 is donated to the luminal soluble electron carrier protein PC, via a 2Fe-2S cluster and cytochrome f subunit of the Cyt b_6f complex, and the two protons are released into the lumen. The second electron is donated to another PQ molecule (protonation of which doubles the number of protons pumped through the thylakoid membrane) via two hemes (or possibly via three; Lavergne 1983, de Lavalette et al. 2009) of the cytochrome b_6 subunit of the Cyt b_6f (Mulkidjanian 2010, Laisk et al. 2016).

Those PQ molecules that mediate electron transfer from PSII to Cyt b_6f comprise the photochemically active PQ pool. In spinach, the size of the pool has been estimated to range from four to 12 PQ molecules/PSII (Graan and Ort 1984, Joliot et al. 1992, Kruk and Strzałka 1999). Plants contain PQ also in plastoglobuli

(chloroplasts localized globules surrounded by a lipid monolayer) and in the chloroplast membrane (Kruk and Karpinski 2006).

Translocation of protons over the thylakoid membrane by the photosynthetic electron transport chain creates an electrochemical gradient, which ATPase uses to catalyze ATP formation from ADP and inorganic phosphate (Fig. 1C). Lumenal pH has been measured to range from 7.5 (low light) to 5.7 (high light) in plants *in vivo* (Takizawa et al. 2007). Besides the pH gradient, a significant part of the proton motive force derives from the electric component (Cruz et al. 2001). In addition to light intensity, the proton motive force is modulated by changes in the conductivity of the ATPase (Kanazawa and Kramer 2002) and via anion and cation transport through the thylakoid membrane (for reviews, see Armbruster et al. 2017, Spetea et al. 2017).

1.1.4 Structure and function of photosystem I

It is generally accepted that PSII and PSI have a common ancestry (possibly more closely related to the present-day PSI; see e.g. Orf et al. 2018). Consequently, PSII and PSI resemble each other. However, the PsaA and PsaB proteins of PSI core comprise both “reaction center” and “inner antenna”. In plants, PSI is surrounded by four antenna proteins (Lhca1–4; collectively called LHCI), organized like a half-moon (for a recent X-ray structure of plant PSI with 2.6 Å resolution, see Mazor et al. 2017). The PSI complex binds over 150 chlorophylls and 30 carotenoids and consists of more than 15 subunits. Also part of the LHCII may serve PSI (see section 1.4.1). In plants, PSI resides in stroma thylakoids (Andersson and Anderson 1980).

Contrary to PSII, where primary charge separation occurs only between the cofactors bound to the D1 protein, electron transfer in PSI may proceed either in A (via cofactors bound to the PsaA protein) or B (via cofactors bound to the PsaB protein) branch (Guergova-Kuras et al. 2001, Santabarbara et al. 2015). However, if one or another of the branches is favored, is unclear (for a review, see Santabarbara et al. 2010). Both branches bind three chlorophyll *a* molecules and a phylloquinone (PhQ). The reaction center of PSI (P700) is formed by a pair of chlorophylls (one in each branch), oriented chlorin rings parallel to each other. After light absorption, primary charge separation may occur between P700 and a chlorophyll called A_0 (either A_0 in A branch or A_0 in B branch). However, it has also been suggested that charge separation can occur in both the branches: between a so-called accessory chlorophyll (which would therefore not be accessory) and the A_0 chlorophyll of each branch (Müller et al. 2003, Ptushenko et al. 2008, Giera et al. 2010). After the primary charge separation, the electron is donated to one of the PhQs. The two branches converge at a 4Fe-4S cluster (named F_X) after which the electron is donated via two additional 4Fe-4S clusters (F_A and F_B) to Fd, a soluble protein containing a

2Fe-2S cluster. After two subsequent electron donations from two reduced Fds, the flavoenzyme FNR catalyzes two electron reduction of NADP⁺ (NADP⁺ + 2e⁻ + H⁺ → NADPH; for a review, see Mulo and Medina 2017). P700⁺ is reduced by PC.

1.2 Singlet oxygen

In an atmosphere with 21 % oxygen, energy and electron transfer reactions inevitably lead to production of reactive oxygen species (ROS). ROS can be defined, for example, as oxygen-containing compounds that are more reactive towards biomolecules than the ground state O₂. Some relevant ROS include hydroxyl radical (HO•), superoxide (O₂^{•-}), hydrogen peroxide (H₂O₂), different peroxides (e.g. lipid peroxides), nitric oxide (NO•), ozone (O₃) and singlet oxygen.

1.2.1 Very shortly about the chemistry of singlet oxygen

The two outermost electrons of ground state molecular dioxygen have equal spins and therefore O₂ is a triplet (³O₂ or formally ³Σ_g⁻O₂). The outermost electrons of ³O₂ (hereafter O₂) occupy their own orbitals (Fig. 3), as two electrons can reside on the same orbital only if their spins are opposite (Pauli exclusion principle). Because biomolecules are commonly singlets, this restricts the reactivity of O₂; reactions of O₂ can occur via one-electron reduction or after spin inversion (with input of energy) in either of the reactants. Spin inversion of one of the electrons of O₂ produces a singlet form of O₂. In the case of ¹Σ_g⁺O₂, the electrons reside on two different orbitals, but this form rapidly decays to ¹Δ_gO₂, where the two electrons reside on the same orbital (Fig. 3). The latter form is commonly abbreviated as ¹O₂.




Figure 3. Different spin configurations of molecular dioxygen (O₂). Ground state triplet O₂ (³Σ_g⁻O₂) can, with addition of energy (hν), be converted to a singlet form (¹Σ_g⁺O₂ or ¹Δ_gO₂). The two outermost electrons of ³Σ_g⁻O₂ and ¹Σ_g⁺O₂ reside on two different molecular orbitals whereas the electrons of ¹Δ_gO₂ reside on a single orbital. Orbitals are represented by circles (names of the orbitals are written on the left). The arrows represent electrons (with spins “up” or “down”). Here, the “double” bond of ³Σ_g⁻O₂ is approximated as a single bond and two “half” bonds, whereas ¹Δ_gO₂ forms a true double bond.

Due to the decreased spin demands, compared to O_2 , 1O_2 rapidly reacts with many compounds, including proteins (e.g. histidine, cysteine, methionine, tryptophan and tyrosine), lipids (specifically unsaturated fatty acids) and nucleic acids (for reviews, see Mattila et al. 2015, di Mascio et al. 2019). Lifetime of 1O_2 is a few μs in water (Merkel and Kearns 1972) and 20–30 μs in lipid micelles (Lee and Rodgers 1983) but has not been measured in photosynthetic organisms *in vivo*. Consequently, the diffusion distance is not known (Mattila et al. 2015, Arellano and Naqvi 2016). Measurements from mammalian and plant systems suggest that little 1O_2 could diffuse out of the cells (Moan 1990, Krasnovsky 1994, Prasad et al. 2017).

1.2.2 Singlet oxygen production by chlorophylls

Intersystem crossing (spin inversion of one of the outermost electrons) converts excited singlet chlorophyll ($^1Chl^*$) to triplet form (3Chl), which may donate the energy to ground state O_2 producing 1O_2 :

1. $^1Chl + \text{light energy} \rightarrow ^1Chl^* \rightarrow ^3Chl$
2. $^3Chl + O_2 \rightarrow ^1Chl + ^1O_2$

Intersystem crossing occurs with over 60 % yield in dilute chlorophyll solutions (Bowers and Porter 1967) making free chlorophylls efficient producers of 1O_2 (e.g. Krasnovsky et al. 1993). Intersystem crossing may produce triplet chlorophylls (and subsequently 1O_2) also in light harvesting antennae.

In the reaction centers, on the other hand, triplet chlorophyll is produced by charge recombination reactions. In a recombination, instead of forward electron transfer, the electron returns to P680 or P700 (see Fig. 2B for PSII). Recombination of the primary radical pair of PSII ($P680^+Pheo_{D1}^-$) can re-excite P680 or produce singlet or triplet reaction center chlorophyll (3P680). If electron transfer to Q_A is not possible, recombination of the primary pair occurs in a few nanoseconds (Schlödter and Brettel 1988). In PSII centers that lack Q_A 3P680 yield is high, 20–30 % per excited reaction center (Takahashi et al. 1987, Durrant et al. 1990). Recombination of electrons at Q_A^- or Q_B^- is slow (1–1.5 and 30–35 ns, respectively) as both reactions are very endergonic (Vermaas et al. 1985, Vass et al. 1990). Usually, TyrZ has had enough time to reduce $P680^+$ and to advance the S-state of OEC, and therefore the S-state is reverted during the recombination: $S_{2/3}Q_{A/B}^- \rightarrow S_{1/2}Q_{A/B}$ (S_1 is stable in darkness). In addition to the recombination route re-exciting P680 (excitonic route) and the route leading to either ground state or triplet P680 via $Pheo_{D1}^-$ (indirect route), $S_{2/3}Q_A^-$ can recombine via direct tunneling (without reproducing the primary radical pair; Rappaport and Lavergne 2009). In PSII centers with an inactive OEC, Q_A^- recombines with $P680^+$ with millisecond kinetics (e.g. de Wijn and van Gorkom

2002a). At $-269\text{ }^{\circ}\text{C}$, $^3\text{P680}$ is located at Chl_{D1} , but at room temperature $\sim 30\%$ of the triplet was suggested to reside at P_{D1} (Diner et al. 2001, Noguchi et al. 2001).

Formation of the triplet reaction center chlorophyll of PSI ($^3\text{P700}$), via recombination of the radical pair $\text{P700}^+\text{PhQ}^-$, occurs in illuminated PSI complexes and chloroplasts (Frank et al. 1979, Rutherford and Mullet 1981). Both $^3\text{P680}$ and $^3\text{P700}$ can be detected based on their electron paramagnetic resonance (EPR) signals (e.g. Rutherford and Mullet 1981).

1.2.3 Antenna or reaction center?

Intersystem crossing takes $\sim 5\text{ ns}$ (Shipman 1979), and transfer of excitation energy from antennae to a reaction center competes with it. In addition, antennae contain carotenoids that efficiently quench both singlet excited and triplet chlorophylls (Peterman et al 1995; see also section 1.2.5), dissipating the energy safely as heat. Spectroscopic studies (Mozzo et al. 2008) showed that certain lutein (a xanthophyll carotenoid) pigments in LHCII quenched 95 % of triplet chlorophylls. Carotenoids can detoxify also $^1\text{O}_2$, produced by the remaining triplets, via physical quenching (energy transfer) or chemical scavenging (oxidation). These considerations have usually led to the assumption that not the antennae but the reaction centers (specifically P680) are the major sites of $^1\text{O}_2$ production in plants, algae and cyanobacteria (for a contradicting view, see Santabarbara et al. 2007). Calculations by Rehman et al. (2013) show that $^3\text{P680}$ accumulation closely resembles measured $^1\text{O}_2$ production. Accordingly, Ramel et al. (2012a) reported that β -carotene (present in PSII and PSI core), but not xanthophylls (in antennae), is oxidized during illumination. Similarly, during a 10 h illumination, no turnover of chlorophyll *b* was observed (Beisel et al. 2010). Chlorophyll *a* turns over in PSII reaction centers but not in LHCs or in PSI (Feierabend and Dehne 1996, Beisel et al. 2010). On the contrary, chlorophylls in LHCII have high turnover in etiolated seedlings upon light exposure (Feierabend and Dehne 1996), suggesting that antennae may be more significant sources of $^1\text{O}_2$ (only) during their biosynthesis.

Direct quenching of triplet chlorophylls by carotenoids diminishes $^1\text{O}_2$ production also in the reaction centers of non-oxygenic bacteria (deWinter and Boxer 1999, Liu et al. 2005) and possibly in the reaction center of PSI (Cazzaniga et al. 2012). The quenching requires that the emission and absorption spectra of the donor and acceptor overlap as well as an overlap between their electron orbitals (Dexter 1953) and therefore cannot occur in PSII reaction center where the closest two β -carotenes are too far, 20–30 Å from $^3\text{P680}$ (Umena et al. 2011). A carotenoid situated nearer would be irreversibly oxidised by P680^+ (formed prior to the triplet), rendering the carotenoid useless (Telfer and Barber 1995). Therefore, if a recombination reaction produces $^3\text{P680}$, it has little other options than to react with

ground state O_2 , producing 1O_2 . β -carotenes in PSII core might still be important quenchers of 1O_2 , as mutants with fewer β -carotene molecules per reaction center suffer from damage to PSII (Cazzaniga et al. 2012; see also section 1.3). It has also been suggested that when Q_A is reduced, 3P680 would localize over Chl_{D2} and could be quenched by the carotenoid Car_{D2} (Martínez-Junza et al. 2008); 3P680 decays faster when Q_A^- is present (van Mieghem et al. 1995).

Previously it was thought that PSI does not produce 1O_2 (Hideg and Vass 1995). Later, maybe due to more sensitive detection methods, 1O_2 production by isolated PSI complexes was observed (Cazzaniga et al. 2012, Takagi et al. 2016), at a rate of about 1/10th of the rate of 1O_2 production by PSII preparations. It has been speculated that donor and/or acceptor side limitation of PSI may enhance 3P700 formation *in vivo* (Rutherford et al. 2012). Based on the above-mentioned measurements, it cannot be judged, however, if the origin of 1O_2 is the antenna of PSI or 3P700 . Based on the presence of oxidized amino acid residues in PSI core and antenna, detected by Kale et al. (2020), both sites may generate 1O_2 . The lifetime of 3P700 is not shortened by the presence of O_2 (Sétif et al. 1981), contrary to what happens to 3P680 (Durrant et al. 1990). Therefore, 3P700 may be shielded from O_2 , or a carotenoid close by very efficiently quenches 3P700 . PSI specific illumination could also not induce oxidation of carotenoids (Ramel et al. 2012b). In addition, it has been reasoned that recombination reactions in PSI occur via the A branch, where a recombination route that does not produce 1O_2 is favored (Rutherford et al. 2012). Taken together, PSI does produce some 1O_2 but little of it may derive from 3P700 .

Cyt b6f complex contains a chlorophyll molecule (for a recent cryogenic electron microscopy structure with 3.6 Å resolution, see Malone et al. 2019). Some evidence has been presented that the chlorophyll produces 1O_2 in the light (Sang et al. 2010), damaging the complex itself (Taylor et al. 2018). It has been suggested, however, that intact Cyt b6f produces little 1O_2 (Ma et al. 2009).

1.2.4 Additional sources of singlet oxygen in plant cells

Apart from photosensitization by chlorophylls, lipid peroxidation is, perhaps, the most relevant route leading to 1O_2 production in (photosynthetic) organisms. Oxidation of lipids leads to formation of lipid radicals, tetroxides and triplet species. Decomposition of the tetroxide may directly produce 1O_2 (Russel mechanism) or the triplet may donate energy to O_2 producing 1O_2 . Once initiated, lipid peroxidation and 1O_2 production can continue for a long time (for reviews, see Miyamoto et al. 2007, Pospíšil et al. 2019). Oxidation of lipids (and, to a smaller degree, of proteins), possibly by $TyrZ^\bullet$, $P680^+$, 1O_2 or HO^\bullet , in PSII preparations with nonfunctional OEC has been shown to coincide with 1O_2 production (Hideg and Vass 1993, Pospíšil et al. 2007, Khorobrykh et al. 2011, Yadav and Pospíšil 2012). In addition,

lipoxygenase enzymes, present in plants, algae and cyanobacteria, catalyze insertion of O₂ into lipids, producing lipid peroxides (for a review, see Andreou and Feussner 2009). Some plant species also synthesize defense compounds that produce ¹O₂ in the light (for reviews, see Tyystjärvi 2004, Flors and Nonell 2006).

1.2.5 Non-photochemical quenching and other protective measures against singlet oxygen

Non-photochemical quenching (NPQ) of chlorophyll fluorescence reflects dissipation of light energy as heat. The so-called energy-dependent quenching (qE) comprises most of regulated NPQ; in plants, acidification of lumen induces protonation of the PsbS protein (Li et al. 2000) and conversion of violaxanthin to zeaxanthin (Bilger and Björkman 1994). The exact mechanism of qE is unclear; reorganization of LHCII and interactions within chlorophylls or between chlorophylls and xanthophylls have been proposed (e.g. Holt et al. 2005, Bode et al. 2009, Müller et al. 2010, Johnson et al. 2011). In any case, NPQ decreases the amount of singlet excited chlorophylls, and consequently triplet chlorophylls, in antennae (Carbonera et al. 2012), leading to a decrease in ¹O₂ production (Dall'Osto et al. 2012, Roach and Krieger-Liszkay 2012, Girolomoni et al. 2017). Whether NPQ affects PSI (Ballottari et al. 2014) or not (Tian et al. 2017) is debated, but also PSI complexes produce less ¹O₂ when isolated from light-acclimated plants compared to those isolated from dark-acclimated plants (Dall'Osto et al. 2012). Instead of PsbS, in the alga (e.g. *Chlamydomonas reinhardtii*) NPQ depends on the LHCSR protein (Peers et al. 2009) and in cyanobacteria on the OCP. In cyanobacteria, the site of the quenching is the phycobilisome core rather than a chlorophyll containing antenna (Wilson et al. 2006).

Accumulation of zeaxanthin can induce NPQ also independently of lumen acidification or the PsbS protein (Dall'Osto et al. 2005, Verhoeven 2014). A newly identified lipocalin regulated NPQ mechanism does not require zeaxanthin, lumen acidification or PsbS (Brooks 2012, Malnoë et al. 2018). In addition, damage to PSII units (section 1.3) causes NPQ. State transitions (section 1.4.1) and chloroplast movements (Cazzaniga et al. 2013, Wilson and Ruban 2020) may affect NPQ measurements but do not regulate thermal dissipation in PSII.

In addition to NPQ, the rate of ¹O₂ production can be affected by modulation of recombination reactions. Some cyanobacteria have high and low light forms of the D1 protein (Kós et al. 2008). Redox properties of the Q_A/Q_A⁻ or Pheo/Pheo⁻ pairs in the high light form of D1 favor direct charge recombinations, thereby decreasing ¹O₂ production (Tichý et al. 2003, Vass and Cser 2009, Vinyard et al. 2013). In plants, exposure to coldness or high light can decrease the redox gap between Q_A and Q_B,

again favoring the direct recombination (Janda et al. 2000, Ivanov et al. 2006, 2008), possibly leading to a decrease in $^1\text{O}_2$ production (Ramel et al. 2012a).

Plants and other photosynthetic organisms contain multiple compounds that can quench or scavenge $^1\text{O}_2$ (for a review, see Pinnola and Bassi 2018). Lipid-soluble carotenoids (discussed above) and tocopherols (see e.g. Kruk et al. 2005, Rastogi et al. 2014) are good quenchers of $^1\text{O}_2$. In addition, lipid-soluble isoprene (Affek and Yakir 2002) and flavonoids, and water-soluble flavonoid glycosides (Majer et al. 2014), ascorbate and glutathione have been reported to offer protection against $^1\text{O}_2$.

PQH₂ molecules react with $^1\text{O}_2$ both *in vitro* and *in vivo* (Gruszka et al. 2008, Kruk and Trebst 2008, Khorobrykh et al. 2015, Ferretti et al. 2018). In plants, the photochemically active PQ pool was shown to become oxidized during high light, presumably due to a reaction with $^1\text{O}_2$ (Kruk and Szymańska 2012). Longer illumination decreased also the total amount of PQ molecules (Ksas et al. 2015, Ferretti et al. 2018). Continued high light stress increased the size of the non-photochemically active PQ pool, supporting the view that *in vivo* PQ serves also as an antioxidant (Ksas et al. 2015).

1.2.6 Where, when and why to bother?

$^1\text{O}_2$ is thought to be a major agent of damage to photosynthetic organisms. During illumination, lipid peroxidation in leaves was shown to be mainly due to $^1\text{O}_2$ (Triantaphylidès et al. 2008) and the majority of the genes induced in *Arabidopsis thaliana* cell cultures by high light were identified as responsive to $^1\text{O}_2$ (González-Pérez et al. 2011). $^1\text{O}_2$ has been reported to have capacity to damage several photosynthetic protein complexes, including PSII and PSI (e.g. Hideg et al. 2007, Takagi et al. 2016; see also sections 1.3.2 and 1.3.3). Conditions producing $^1\text{O}_2$ induce the protective mechanisms discussed in the previous section. $^1\text{O}_2$ also directly initiates signaling pathways leading either to high light acclimation or programmed cell death (for a review, see e.g. Crawford et al. 2018).

Under which conditions is $^1\text{O}_2$ produced? Even though discussion about $^1\text{O}_2$ commonly relates to high light, $^1\text{O}_2$ -oxidized lipids and carotenoids have been observed even in low light (Triantaphylides et al. 2008, Ramel et al. 2012a). $^1\text{O}_2$ production increases linearly with increasing intensity of light, both *in vitro* (Fufezan et al. 2002) and *in vivo* (Rehman et al. 2013). It is also often stated that reduction of the PQ pool increases $^1\text{O}_2$ production (by decreasing forward electron transfer and promoting recombination reactions). However, caution should be used, as the rate of $^1\text{O}_2$ production is not only determined by the recombination rate but also by the type of the recombination (e.g. Fufezan et al. 2002). Furthermore, only rarely measurements of $^1\text{O}_2$ are presented. Chemical inhibition of CO₂ fixation leads to a ~30 % increase in $^1\text{O}_2$ production in a dinoflagellate alga *Symbiodinium* (Rehman et

al. 2016a). In an *A. thaliana* mutant (*chlorinal*), $^1\text{O}_2$ production increases two to three fold when CO_2 concentration decreases to 50 ppm (Ramel et al. 2013); unfortunately, $^1\text{O}_2$ was measured with singlet oxygen sensor green (SOSG) in white light, when the sensor itself may produce $^1\text{O}_2$ (Ragás et al. 2009).

Accumulation of free chlorophylls inside cells would pose a risk for photosynthetic organisms (as well as for herbivorous protists; Kashiyama et al. 2012), as demonstrated in mutants with impaired control of chlorophyll synthesis (Przybyla et al. 2008). $^1\text{O}_2$ production by chlorophyll catabolites may be relevant also *in vivo*, e.g. in pathogen attack (Mur et al. 2010, Hörtensteiner and Kräutler 2011, Tarahi Tabrizi et al. 2016), but these processes are not well understood. In addition, it has been suggested that $^1\text{O}_2$ is produced during repair of damaged PSII units (Feierabend and Dehne 1996, Wang et al. 2016a; section 1.3.3). Removal of OEC favors recombination reactions that do not produce $^1\text{O}_2$ (Spetea et al. 1997, Brinkert et al. 2016; see also section 5.2.4) and therefore PSII core may not produce $^1\text{O}_2$ during repair. On the other hand, detachment of some of the chlorophylls bound to PSII core or antennae could promote $^1\text{O}_2$ production. However, to my knowledge, no direct evidence exists.

How significant are sources other than chlorophyll in $^1\text{O}_2$ production? In general, $^1\text{O}_2$ production is much higher in the light than in darkness (Hideg et al. 1994) and in photosynthetic tissues than in non-photosynthetic ones (Fig. 4A). Experiments with added electron donors or metal chelates suggest that lipid peroxidation is a significant source of $^1\text{O}_2$ in PSII preparations with inactive OEC (Pathak et al. 2017). *In vivo*, this may become relevant during stresses that inactivate PSII, such as high temperature (Yamashita et al. 2008, Rehman et al. 2016a). However, Spetea et al. (1997) measured no $^1\text{O}_2$ production in Tris-washed (OEC removed) PSII. Also enzymatic lipid peroxidation may lead to significant $^1\text{O}_2$ production under specific stresses. Prasad et al. (2017) showed that a chloroplast localized lipoxygenase induces $^1\text{O}_2$ production upon wounding. $^1\text{O}_2$ production, probably due to lipoxygenases (Chen and Fluhr 2018), was observed also inside roots after wounding (Mor et al. 2014). In addition, $\text{HO}\cdot$ induces lipid peroxidation upon pathogen attack (Zoeller et al. 2012).




Figure 4. Singlet oxygen and photoinhibition. A. $^1\text{O}_2$ production in white (containing only undifferentiated plastids) or green (containing functional chloroplasts) leaf sections of *Hordeum vulgare* (cv. *Variegata*). Leaf discs were vacuum-infiltrated, incubated overnight in 200 μM SOSG and fluorescence (excitation wavelength 500 nm) was measured before and after a 90 min red ($\lambda > 650$ nm) light illumination (photosynthetic photon flux density, PPFD, 2000 $\mu\text{mol m}^{-2} \text{s}^{-1}$). Curves are averaged from at least three replications. Wavelength regions corresponding to SOSG and chlorophyll fluorescence are indicated. An increase in SOSG fluorescence corresponds to $^1\text{O}_2$ production and a decrease in chlorophyll fluorescence to NPQ formation and photoinhibition. B. Photoinhibition in heavy water (deuterium oxide; D_2O). Excised leaves of *A. thaliana* were incubated overnight with petioles in water (control) or in D_2O , in the absence or presence of 0.4 mg/ml lincomycin and illuminated in white light (PPFD 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$). After 0–120 min of illumination, PSII yield was quantified by measuring the fluorescence parameter F_v/F_M after 30 min in dark. Error bars show standard deviations (SD) from five biological replications. Lines show a fit to the first-order reaction equation (see Materials and methods). Significances of differences between the last time points were calculated with Student's t-test. Measurements by Pooneh Sotoudehnia (A) and own work (B), Molecular Plant Biology, University of Turku.

1.3 Photoinhibition

Here, the term photoinhibition of PSII is used to describe light-induced irreversible damage to PSII that leads to loss of charge separation and O_2 evolution activities of PSII and to proteolytic degradation of the reaction center protein D1. In order to recover from photoinhibition, the D1 protein is replaced in the so-called repair cycle (section 1.3.3). PSII photoinhibition occurs in all oxygenic photosynthesis (with a possible exception of a desert alga *Chlorella ohadii*; Treves et al. 2016), from very low light to intensities several times higher than required to saturate photosynthesis. The rate of the damage is directly proportional to the intensity of light (Tyystjärvi and Aro 1996). Inactive PSII units accumulate (net photoinhibition), if repair reactions cannot keep up with the rate of the damage; this may occur e.g. in high light.

1.3.1 Photodamage to photosystem II in ultraviolet radiation

Ultraviolet (UV) radiation (200–400 nm) causes photoinhibition of PSII with much higher quantum yield than visible light (Jones and Kok 1966a), damaging mostly OEC, while the quinones and TyrZ are also affected (Renger et al. 1989, Vass et al. 1996). Loss of the reaction center activity (i.e. charge separation ability) of PSII coincides with the loss of the D1 protein (Hakala et al. 2005). The reason for the damage is probably absorption of UV radiation by Mn^{+3} or Mn^{+4} ions of OEC (Ruettinger et al. 2000, Szilárd et al. 2007), possibly leading to release of a Mn ion (Hakala et al. 2005). The exact mechanism is not clear. UV-B (280–315 nm) does not induce 1O_2 production, but formation of $HO\bullet$ was observed (Hideg and Vass 1996). The $HO\bullet$ scavenger propyl gallate also partly protected PSII from UV-B induced photoinhibition *in vitro* (Lazarova et al. 2014).

UV-A (315–400 nm) damages PSII with a lower yield than UV-B or UV-C (200–280 nm), but sunlight on the surface of Earth contains much more UV-A than UV-B or UV-C. Therefore, UV-A has a high capacity to cause damage under natural conditions (Sarvikas et al. 2006); 35 % of sunlight-induced photoinhibition in indoors-grown pumpkin leaves was caused by UV-A (Hakala-Yatkin et al. 2010). Many organisms have evolved protective mechanisms, such as compounds screening UV radiation (see Rozema et al. 2002, Pescheck et al. 2014, Tilbrook et al. 2016). Indeed, removal of UV-A from sunlight did not affect photoinhibition in outdoors-grown pumpkin leaves as, due to the presence of protective compounds, the UV-A radiation did not reach thylakoid membranes (Hakala-Yatkin et al. 2010).

1.3.2 How does visible light damage photosystem II?

The mechanism of photoinhibition of PSII under visible light is debated (for reviews, see Vass 2012, Tyystjärvi 2013, Nishiyama and Murata 2014). The main mechanisms discussed in recent literature may be roughly divided as 1O_2 or donor side hypotheses. According to the latter, the damage to PSII is caused by light absorption of the Mn ions of OEC, similarly as under UV radiation (Hakala et al. 2005, Ohnishi et al. 2005). Alternatively, long-lived $P680^+$ or $TyrZ\bullet$, observed in OEC-less PSII, may extract electrons from their surroundings (Jegerschöld et al. 1990, Anderson et al. 1998). In the 1O_2 (or recombination) hypotheses, PSII is inactivated by oxidation of the D1 protein by 1O_2 produced via recombination reactions (Keren et al. 1997, Vass 2012). In addition, 1O_2 produced in antennae (Santabarbara et al. 2002) or other ROS (e.g. Song et al. 2006) have been proposed to cause photoinhibition of PSII.

Several lines of evidence support the involvement of recombination reactions and 1O_2 in photoinhibition of PSII. 1O_2 can damage PSII and induce degradation of the D1 protein *in vitro* (Durrant et al. 1990, Hideg et al. 2007). 1O_2 production and

photoinhibition have similar responses to intensity (Rehman et al. 2013) and quality (Jung and Kim 1990) of light, however, Jung and Kim (1990) used a $^1\text{O}_2$ detection method (imidazole and p-nitrosodimethylaniline) that inactivates PSII (Stemler and Jursinic 1983). Recombination reactions modulated to decrease or increase $^1\text{O}_2$ production have been shown to decrease or increase, respectively, the rate of the damage to PSII (Fufezan et al. 2002, Cser and Vass 2007, Rehman et al. 2013, Davis et al. 2016, Treves et al. 2016). In addition, quenchers and scavengers of $^1\text{O}_2$, such as carotenoids, protect PSII (e.g. Barényi and Krause 1985, Cazzaniga et al. 2012, Hakkila et al. 2014), even though the protection is not always observed (Tschiersch and Ohmann 1993, Nishiyama et al. 2001, Hakala-Yatkin et al. 2011). Dissipation of excess light energy decreases photoinhibition of PSII (e.g. Sarvikas et al. 2006), however, to a degree that may be too small to fully support the recombination hypothesis (Tyystjärvi et al. 2005). On the other hand, it was argued that conventional NPQ parameters do not correctly estimate the protective effect (Lambrev et al. 2012). Measurements by Dall'Osto et al. (2012) suggest that an NPQ deficient mutant (*npq1*) produces about two times as much $^1\text{O}_2$ as the wild type. Photoinhibition proceeded, according to Sarvikas et al. (2006), 5–20% faster in the mutant than in the wild type.

Experimental results support also $^1\text{O}_2$ -independent mechanisms. Photoinhibition proceeds under anaerobic conditions (Satoh 1970, Chaturvedi et al. 1992, Hakala et al. 2005, Fan et al. 2016), where no $^1\text{O}_2$ is produced (Hideg et al. 1994). The lifetime of $^1\text{O}_2$ is lengthened by ~ 10 times in D_2O , but photoinhibition *in vivo* does not proceed faster in D_2O than in water (Fig. 4B). The action spectrum of photoinhibition of PSII (reviewed by Zavafer et al. 2015a) resembles the absorption spectra of Mn-clusters (Hakala et al. 2005; for an artificial Mn-cluster closely resembling that of PSII, see Zhang et al. 2015) and is dramatically different in OEC-less PSII (Hakala et al. 2005). However, peaks at orange and red lights, observed in some of the measured action spectra, support the idea that photoinhibition is at least partially driven by light absorbed by PSII antenna (e.g. Santabarbara et al. 2002, Tyystjärvi et al. 2002, Havurinne and Tyystjärvi 2017). In the Mn-hypothesis (Hakala et al. 2005), also called two-step mechanism (Ohnishi et al. 2005), after initial inactivation of OEC, P680^+ was assumed to cause additional damage. If OEC could recover from the initial damage (without the repair cycle) and an additional damage, via a P680 related mechanism (or via $^1\text{O}_2$), would be needed to solidify the inactivation, a deviation from first order kinetics should be observed. However, photoinhibition of PSII proceeds with first order kinetics (e.g. Sarvikas et al. 2010). Therefore, to comply with the photoinhibition kinetics, the initial damage must already irreversibly inactivate PSII and any secondary damage should inactivate another, still functional PSII unit (Tyystjärvi 2013).

The molecular nature of the event inactivating PSII is not clear, but damage to P680, Phe_{D1}, the quinones (Cleland 1986, Ohad et al. 1990, Hou 2014) or OEC have been suggested. Kale et al. (2017) identified several oxidative modifications in the D1 and D2 proteins with mass spectroscopy. After 15 min of illumination, most oxidized amino acids were found in the vicinity of OEC whereas after 30 min of illumination modifications close to P680 appeared (Kale et al. 2017). This agrees with the finding that, during photoinhibition, the reaction center activity of PSII is less affected than the water splitting capacity (Chaturvedi et al. 1992, Hakala et al. 2005, Zavafer et al. 2015b). Several oxidized histidine and tryptophan residues (Kale et al. 2017) may speak for the involvement of ¹O₂, however, the authors proposed that HO• produced after H₂O₂ formation during incomplete water oxidation by OEC and O₂•⁻ formed by O₂ reduction by Phe_{D1}⁻ and/or Q_A⁻ caused the damage.

1.3.3 The repair cycle

Repair of damaged PSII units (for reviews, see Järvi et al. 2015, Theis and Schroda 2016) starts with monomerization of the PSII dimer and in plants with migration of PSII to grana margins or to stroma thylakoids (Yamamoto et al. 1981, Baena-González et al. 1999, Aro et al. 2005, Herbstová et al. 2012, Nagarajan and Burnap 2014). Specific zones for PSII repair may exist also in cyanobacteria (Sacharz et al. 2015). In plants, these steps are, at least partly, controlled by phosphorylation and dephosphorylation of the D1 (and D2) protein (Koivuniemi et al. 1995). PSII is partially disassembled (Christopher and Mullet 1994, Weisz et al. 2019), the D1 protein is degraded by Deg1 and FtsH proteases (Ohad et al. 1984, Lindahl et al. 2000, Silva et al 2003, Kapri-Pardes et al. 2007, Krynická et al. 2015, Knopf and Adam 2018) and a new D1 protein is co-translationally inserted into the PSII complex (Zhang et al. 1999, Zhang et al. 2001). Finally, PSII is reassembled (Bowyer et al. 1992, Rokka et al. 2005).

The repair cycle is sensitive to ROS: H₂O₂, ¹O₂ and O₂•⁻ all inhibit synthesis of the D1 protein (but did not affect the rate of the damage; Nishiyama et al. 2001, 2004, Jimbo et al. 2013). In *Synechocystis*, inactivation of the translation machinery is due to oxidation of cysteine residues in two elongation factors, EF-Tu and EF-G (Kojima et al. 2009, Yutthanasirikul et al. 2016). In *A. thaliana*, high light stress may transiently inactivate also the FtsH protease (Zaltsman et al. 2005, Kato et al. 2018). It was recently shown that the enzyme converting zeaxanthin back to violaxanthin upon NPQ relaxation is inactivated during photoinhibition, which possibly protects PSII repair (Bethmann et al. 2019).

As damage proceeds simultaneously with repair, it is often useful to separate the processes in research. In plants and algae, the D1 protein is coded by the chloroplast genome, and inhibitors of bacterial translation, most commonly lincomycin or

chloramphenicol (of these two, lincomycin may be preferred as it is shown not to block nuclear translation and chloramphenicol produces $O_2^{\bullet-}$; Okada et al. 1991, Mulo et al. 2003, Rehman et al. 2016b, Kodru et al. 2020), can be used to block the repair (for an example, see Fig. 4B).

1.3.4 Photoinhibition of photosystem I

PSI too can be damaged in light, specifically at low temperatures (Terashima et al. 1994) and under high (e.g. Huang et al. 2015) or fluctuating light (Tsuyama and Kobayashi 2008, Suorsa et al. 2012). In the absence of (other) electron acceptors, PSI donates electrons to O_2 , producing $O_2^{\bullet-}$ (Asada et al. 1974; for the site of the production, see Kale et al. 2020 and discussion in Khorobrykh et al. 2020a). $O_2^{\bullet-}$, or other ROS derived from $O_2^{\bullet-}$ (H_2O_2 via dismutation and HO^{\bullet} via dismutation and Fenton chemistry), may damage PSI (Takahashi and Asada 1988, Sonoike 1996, Tjus et al. 1998, Grieco et al. 2012), possibly the Fe-S clusters (Inoue 1986, Tiwari et al. 2016). In addition, based on e.g. protective effects of scavengers of 1O_2 , a role for 1O_2 has been suggested (Rajagopal et al. 2005, Cazzaniga et al. 2012, Takagi et al. 2016, Kale et al. 2020). As PSI photoinhibition requires O_2 (and electrons from PSII), it does not occur under anaerobicity (Sato 1970). Contrary to PSII, UV radiation does not damage PSI (Jones and Kok 1966b). Recovery from PSI photoinhibition is costly (Zhang and Scheller 2004, Lima-Melo et al. 2019).

It has been suggested that the function of photoinhibition of PSII is to protect PSI (Sonoike 2011, Tikkanen et al. 2014). So, is the extent of PSII repair regulated (the extent of the PSII damage increases linearly with light, give or take some modulation by the above described protective measures such as NPQ)? Replacement of a cysteine residue in the cyanobacterial EF-Tu with an amino acid not sensitive to ROS rendered EF-Tu more tolerant against high light stress and improved PSII repair (Jimbo et al. 2018). Thus, the susceptibility of the translation machinery to oxidative stress may not be a necessity of its function but a regulation mechanism. However, in *Synechocystis*, transcription for the messenger RNA of the D1 protein increases in high light (Mohamed and Jansson 1989) following the amount of PSII photoinhibition (Tyystjärvi et al. 2002). Also the amount of EF-Tu increases in high light, leading to decreased net photoinhibition (Jimbo et al. 2019). Exposure to high light leads to increased amount and/or activity of FtsH in *C. reinhardtii* (Wang et al. 2017) and in plants (Sinvány-Villalobo et al. 2004), enhancing the repair of PSII (Tyystjärvi et al. 1992, Aro et al. 1994, Serôdio et al. 2017). Therefore, at least in continuous high light, it seems more important to keep PSII functional, rather than to limit the electron flow to PSI (by means of PSII photoinhibition; see also section 1.4.2). Sustained low electron transfer rates of PSII (compared to those of PSI) were proposed to protect PSI during sunflecks in a shade plant (quantitative comparisons

are, however, not possible based on fluorescence data only, as presented by Sun et al. (2020). Measurements of PSI functionality from *C. ohadii*, which can grow under extreme light intensities while PSII seems not to get inhibited (Treves et al. 2016), or from the cyanobacterial mutants in which the normally sensitive translation is made more tolerant against ROS (Jimbo et al. 2018), may be of interest.

1.4 Responses to changing quantity and quality of light

Photosynthetic organisms have evolved multiple mechanisms to cope with constantly changing natural conditions (for an interesting example of a plant experiencing fluctuating light intensities, see Townsend et al. 2018). Synthesis of light screening compounds (e.g. against UV radiation) can reduce the amount of irradiation received by the photosystems. Plants (and other photosynthetic organisms) can optimize light capture by leaf and chloroplast movements and by reorganizing the thylakoid membrane and photosynthetic machinery, for example by modulating antenna sizes (e.g. Kasahara et al. 2002, Khatoon et al. 2009, Betterle 2009, Zhao et al. 2020). If light is, however, absorbed in excess, NPQ can dissipate it as heat (section 1.2.5). Some of the alternative electron transfer pathways may be considered as dissipative, too. Excess light can be also partly utilized by enhancing sinks (e.g. Seemann et al. 1987). If produced in excess, ROS can be detoxified (see section 1.2.5 for $^1\text{O}_2$). Finally, damaged cell components can be repaired or replaced (see e.g. Li et al. 2018, Nakamura et al. 2018). Some of these mechanisms are discussed in more detail below.

1.4.1 Balancing the photosystems: roles for plastoquinone redox state

PQ molecules mediate electron transfer from PSII to PSI and, consequently, in low light, the redox state of the PQ pool depends on the relative electron transfer rates of the two photosystems. In high light, electron transfer through Cyt b6f complex becomes the limiting step (Stiehl and Witt 1969; section 1.4.2), which leads to reduction of the PQ pool (note that $^1\text{O}_2$ produced by high light can also oxidize PQH_2 ; section 1.2.5).

State transitions change excitation distribution between PSII and PSI. The STN7 kinase (STT7 in *C. reinhardtii*) is activated by reduction (occupancy of a PQH_2 molecule at the luminal Q_o site of the Cyt b6f complex) of the PQ pool (Vener et al. 1995, Zito et al. 1999). How exactly the luminal redox message is transferred to the STN7 residing on the stromal side of the thylakoid membrane is unclear (see e.g. Vladkova 2016, Dumas et al. 2017). In any case, LHCI is phosphorylated (Depège

et al. 2003, Bellaﬁore et al. 2005, Pietrzykowska et al. 2014), which leads to an increase in the absorption cross-section of PSI and/or to a decrease in the absorption cross-section of PSII (state 1 → state 2). Which LHCII (those bound to PSII super-complex, or free, “extra” LHCII, those in grana margins, or all) migrate to PSI, and to which degree (if at all), is not clear (e.g. Delosme et al. 1996, Wientjes et al. 2013a, Grieco et al. 2015, Ünlü et al. 2015, Bos et al. 2019). Oxidation of the PQ pool activates the TAP38 phosphatase (PPH1 in *C. reinhardtii*, where also PBCP may be involved; Cariti et al. 2020), which leads to dephosphorylation of LHCII and back to state 1.

In plants, state transitions optimize light absorption under low light (state 2 is not induced in high light even though the PQ pool may be reduced, probably due to inactivation of STN7 by thioredoxins; Rintamäki et al. 2000, Ancín et al. 2019) or fluctuating light (Bellaﬁore et al. 2005, Wagner et al. 2008, Tikkanen et al. 2010) and after PSI photoinhibition (Rantala et al. 2020b). The amplitude of state transitions can be much bigger in *C. reinhardtii* than in plants (see e.g. Delosme et al. 1996, Drop et al. 2014), CP29 and CP26 are involved (Kargul et al. 2005) and a part of LHCII may disconnect from any photosystem in state 2 (Nagy et al. 2014). *C. reinhardtii* mutants incapable of state transitions suffer in high light, possibly due to increased production of H₂O₂ (Allorent et al. 2013). Redox state of the PQ pool has been shown to regulate state transitions also in cyanobacteria (Mullineaux and Allen 1990, Mao et al. 2002) but the mechanism and the contribution of the phycobilisome are not resolved (e.g. Ranjbar Choubeh et al. 2018; for a review, see Kirilovsky 2015). In addition, it was recently shown that neither the Cyt b6f complex nor phosphorylation events are involved in cyanobacterial state transitions (Calzadilla et al. 2019).

In addition, the redox state of the PQ pool has been shown to regulate the expression of chloroplast and nuclear genes. For example, LHCII transcription was shown to be affected in the green alga *Dunaliella tertiolecta* (Escoubas et al. 1995, Chen et al. 2004) and in plants (Yang et al. 2001). In cyanobacteria, the stoichiometry of the photosystems was suggested to be under the control of PQ redox state (Allen et al. 1989). However, wider scale transcription studies do not show a clear set of genes regulated by the redox state of the PQ (see e.g. Piippo et al. 2006, Bräutigam et al. 2009, Pesaresi et al. 2009, Bode et al. 2016). Signaling cascades are not well known, but an involvement of the STN7 kinase has been proposed (Pesaresi et al. 2009, Berger et al. 2014, Petrillo et al. 2014). Recently it was shown that self-phosphorylating activity of the chloroplast sensor kinase is inhibited by PQH₂ (Ibrahim et al. 2020). The kinase represses transcription of some chloroplast genes and was proposed to transfer the PQ pool redox signal to gene expression changes (Puthiyaveetil et al. 2008, Ibrahim et al. 2020). In addition, H₂O₂ produced by PQH₂ (Khorobrykh et al. 2015; for a review, see Khorobrykh et al. 2020a) has been

proposed to act as a signaling molecule, regulating e.g. stomatal closure and PSII antenna size (Borisova-Mubarakshina et al. 2015, Wang et al. 2016b).

1.4.2 Photosynthetic control

Oxidation of PQH₂ by the Cyt b6f complex may take several milliseconds (Stiehl and Witt 1969, Laisk et al. 2016), making it the slowest step of the photosynthetic electron transport chain (for discussion, see Hasan and Cramer 2012, Tikhonov 2018). Lumen acidification below pH ~6.5 further slows down oxidation of PQH₂ at the Cyt b6f (Tikhonov et al. 1984, Hope et al. 1994). Other mechanisms decreasing the activity of Cyt b6f may also exist (Heimann et al. 1998, Laisk et al. 2005). This limitation of the electron transfer rate by lumen acidification (commonly due to high light) is termed “photosynthetic control”. Photosynthetic control keeps PSI sufficiently oxidized preventing photoinhibition of PSI (Grieco et al. 2012, Suorsa et al. 2012).

1.4.3 Cyclic pathways and other alternative routes

In addition to the linear electron transfer, several alternative routes exist. A cyclic electron transfer (i.e. separate from charge recombinations) within PSII is assumed to function: cytochrome b559 may reduce P680⁺, possibly via the β -carotene Car_{D2} and/or the chlorophyll ChlZ_{D2} (Knaff and Arnon 1969, Buser et al. 1992, Shinopoulos et al. 2014). Cytochrome b559 is then reduced by electrons from Q_B⁻ or PQ pool (for a review, see Shinopoulos and Brudvig 2012). Simultaneous proton pumping from Q_BH₂ to lumen has been proposed (Ananyev et al. 2016, 2017) even though, to my understanding, no convincing evidence exists. The pathway protects PSII either by dissipating excess energy when linear electron transfer and NPQ cannot keep the PQ pool sufficiently oxidized, or by preventing accumulation of oxidizing Car_{D2}⁺ (Telfer et al. 1991, Barber and De Las Rivas 1993, Miyake and Okamura 2003, Faller et al. 2005, Feikema et al. 2006, von Sydow et al. 2016).

In contrast to the cyclic electron transfer within PSII, cyclic electron transfer from PSI (most probably via reduced Fd) back to the PQ pool conserves energy in the form of ATP (but not NADPH; see Fig. 1C). Two pathways are known to function; the PGR5 and PGRL1 dependent route (Munekage et al. 2002, DalCorso et al. 2008, Hertle et al. 2013) and the NAD(P)H dehydrogenase (or NADH dehydrogenase-like) complex dependent route (e.g. Shikanai et al. 1998). In plants, the PGR5 route seems to be more important (Munekage et al. 2004).

These cyclic pathways may balance the ATP/NADPH ratio. Based on theoretical predictions, 4.7 protons need to go through the ATPase of chloroplasts to produce an ATP molecule (e.g. Rastogi and Girvin 1999, Seelert et al. 2000), however,

measurements rather suggest ~ 4 (van Walraven et al. 1996, Petersen et al. 2012). Assuming the latter, linear electron transport would produce an ATP/NADPH ratio of $3/2$ ($=1.5$), which exactly matches the ATP/NADPH need of CO_2 fixation reactions. Accordingly, Laisk et al. (2005) did not observe much cyclic electron transfer under low light, suggesting little “ATP deficiency”. In addition to CO_2 fixation, ATP is needed for other chloroplast functions, such as nitrogen metabolism and protein import and synthesis. It has been suggested that cyclic electron transfer is important also in supplying ATP for the repair cycle of PSII (Endo et al. 1999, Huang et al. 2018a, Murata and Nishiyama 2018).

Cyclic electron transfer may significantly contribute to the total electron transfer during stress conditions or in the case of sink limitations (e.g. Harbinson and Foyer 1991) and during dark to light transitions (Joliot and Joliot 2002). Importantly, cyclic electron transfer induces lumen acidification, proposed to protect both PSII and PSI from photoinhibition by inducing NPQ and photosynthetic control, respectively (Munekage et al. 2002, DalCorso et al. 2008, Suorsa et al. 2012, Yamori and Shikanai 2016). Accordingly, PSI is damaged in mutants incapable of cyclic electron transfer (Munekage et al. 2004, Suorsa et al. 2012). However, the sensitivity of the *pgr5* mutant (Suorsa et al. 2012) may not be due to the inactivated cyclic electron transfer *per se* but because of impairments in Cyt b6f complex (Buchert et al. 2020) or proton conductivity of the ATPase (Avenson et al. 2005). Upon high light, acidification of lumen is accelerated also by ion channels and decreased proton conductivity of the ATPase (Armbruster et al. 2017, Huang et al. 2018b).

O_2 reduction by PSI or plastid terminal oxidase (for a review, see Nawrocki et al. 2015) may at least in some conditions contribute to oxidation of the PQ pool (Asada 2000, Li et al. 2016, Kambakam et al. 2016, Rantala et al. 2020b). In the dark, NADH dehydrogenase-like complex may reduce the PQ pool (Burrows et al. 1998). In cyanobacteria, some algae and gymnosperm plants, flavodiiron proteins function as efficient electron acceptors of PSI (for a review, see Alboresi et al. 2019).

2 Aims of the study

The present work intends to find answers to basic research questions, related to photosynthetic reactions and their regulation in plants and cyanobacteria, which have so far remained unresolved. More specifically, I aimed:

1. To develop detection methods for $^1\text{O}_2$, and to measure $^1\text{O}_2$ from photosynthetic organisms in order to understand the origin and role of $^1\text{O}_2$ in photosynthesis, especially as a damaging agent,
2. To resolve the mechanism(s) of photoinhibition of PSII, and
3. By directly measuring the redox state of the photochemically active PQ pool from plants, to better understand how the redox state of the PQ pool responds to different light conditions.

3 Materials and methods

3.1 Organisms and growth conditions

The main organisms used in the present work were wild types of the plants *A. thaliana* and *Cucurbita maxima* (pumpkin) and the cyanobacterium *Synechocystis* sp. PCC 6803. Plants were grown under PPFD of 100–200 $\mu\text{mol m}^{-2} \text{s}^{-1}$ at ~ 20 °C with day/night rhythms of 8 h/16 h for *A. thaliana* and 16 h/8 h for *C. maxima*. *Synechocystis* was grown in BG-11 (Rippka et al. 1979; buffered to pH 7.5 with 20 mM Hepes-KOH) as batch cultures with mixing under constant PPFD of 40 $\mu\text{mol m}^{-2} \text{s}^{-1}$ at 32 °C. Thylakoid membranes were isolated from leaves as described in Hakala et al. (2005) and stored at -75 °C until use.

3.1.1 Light sources

“White” lights may greatly differ from each other. Spectra of growth lights for *A. thaliana* (growth chamber) and *C. maxima* (greenhouse), as well as spectra of the Sun and several high light sources used here are presented in Fig. 5.




Figure 5. Spectra of different light sources. A. Photon number spectra, normalized to equal number of photons at 400–800 nm, from Osram Powerstar HQI-BT lamps (a Weiss Gallenkamp growth chamber), a halogen lamp (Greenhouse), the Sun (measured on 27.2.2020 on a cloudless day in Turku, Finland), an LED (light emitting diode) based sunlight simulator (SLHolland) and a high-pressure xenon lamp (Sciencetech Inc). B. The scale of the Y-axis is reduced for better visualization of the continuous spectra.

Monochromatic illumination was obtained with different LEDs (LED Fedy, China) or defined by filters (e.g. Corion, Newport Corporation), UV radiation with VL-8.LC and VL-8.M lamps (Vilber Lourmat, France) and short flashes with a Nd:YAG laser (532 nm, 4 ns; Continuum, USA) or an FX-200 xenon lamp (3 μ s; EG&G, USA). Intensity and spectra were measured with wavelength-calibrated light sensors, LI-189 (LiCor, USA) and STS-VIS (Ocean Optics).

3.2 Chlorophyll *a* fluorescence and P700 signal

Photosynthesis can be studied by chlorophyll *a* fluorescence; fluorescence competes with charge separation and heat dissipation (in PSII). Pulse amplitude modulated (PAM) fluorometry was used to probe the quantum yield of PSII photochemistry (F_V/F_M), electron transfer rate, NPQ, photochemical quenching (qP or qL; Maxwell and Johnson 2000, Kramer et al. 2004) and other PSII parameters. Recombination reactions were measured with thermoluminescence: a thylakoid sample was given a Xenon flash at -10 °C and fluorescence yield was recorded while heating the sample at a constant rate (0.56 °C/s). Absorption cross-sections of PSII and PSI were studied by measuring fluorescence spectra at -196 °C (77 K; in liquid N₂): a dilute thylakoid sample (to prevent re-absorption of fluorescence resulting in an over-estimation of light absorption by PSI) was excited with monochromatic light with a wavelength shorter than those of chlorophyll fluorescence.

Contribution of PSI to variable fluorescence at room temperature is small (Franck et al. 2002). Instead, redox states of P700 and PC were followed by measuring absorbance changes at specific near-infrared (NIR) wavelengths. NIR signals recorded by DUAL-KLAS-NIR fluoro/spectrometer (Walz GmbH) were empirically deconvoluted by Schreiber and Klughammer (2016).

3.3 Quantification of photoinhibition

Photoinhibition of PSII was measured by the fluorescence parameter F_V/F_M (after dark incubation to relax NPQ) or by light-saturated rate of O₂ evolution of thylakoids or *Synechocystis* cells in the presence of artificial electron acceptors. In the case of *Synechocystis* cells or intact leaves, the repair cycle was blocked with lincomycin. In the absence of repair, the rate constant of photoinhibition of PSII (k_{PI}) was calculated by fitting the loss of PSII activity to the first-order reaction equation ($y=A*e^{-k_{PI}*t}$, where y is F_V/F_M or O₂ evolution and t is time. A was fixed to a control value of F_V/F_M or PSII oxygen evolution). In isolated thylakoid membranes, also the rate of PSII inactivation in darkness was measured, and this rate constant was subtracted from the final values.

3.4 Measurements of singlet oxygen

$^1\text{O}_2$ was measured from thylakoid membranes with 2,2,6,6-tetramethylpiperidine (TEMP) or histidine. The reaction of TEMP with $^1\text{O}_2$ is specific to $^1\text{O}_2$ (Zang et al. 1995) and produces 2,2,6,6-tetramethylpiperidine 1-oxyl (TEMPO), which can be detected with EPR (Lion et al. 1976). Here, a method to quantify TEMPO with ultra high-performance liquid chromatography ((U)HPLC; Acquity UPLC, Waters Corporation, USA) combined with an electrospray ionization triple quadrupole mass spectrometer (Xevo TQ, Waters Corporation) was developed. The column used was Acquity UPLC BEH Phenyl (Waters Corporation). Acetonitrile and 0.1 % formic acid were used as solvents. The rate of the reaction of TEMP with $^1\text{O}_2$ increases with pH (Houba-Herlin et al. 1982), and therefore an alkaline pH 8.7 was used.

Histidine reacts with $^1\text{O}_2$, but not with H_2O_2 or $\text{O}_2^{\bullet-}$ (Rehman et al. 2013). In a biological system, $^1\text{O}_2$ may be deactivated by collisions with water molecules ($\text{O}_2 \rightarrow ^1\text{O}_2 \rightarrow \text{O}_2$), other quenchers and potential reaction partners (e.g. Merkel and Kearns 1972); in the presence of histidine, less of $^1\text{O}_2$ is deactivated by collisions with water. Therefore, a decrease in O_2 concentration in the presence of added histidine reflects $^1\text{O}_2$ production by the sample (Telfer et al. 1994, Rehman et al. 2013). O_2 concentration was recorded with an O_2 electrode (Hansatech Instruments Ltd, UK) or an optical sensor (Firesting, PyroScience GmbH). Isolated thylakoid membranes consume O_2 in the absence of histidine (mainly due to O_2 reduction by PSI; Asada et al. 1974) and thus measurements in the presence and absence of histidine are needed. Different stable isotopes of O_2 (^{16}O and ^{18}O ; ^{18}O is rare in the nature) were distinguished with membrane inlet mass spectroscopy (MIMS); gases diffused from a sample chamber through a PTFE membrane via vacuum lines to Prima PRO Process Mass Spectrometer (Thermo Scientific™). The dye Rose Bengal was used to chemically produce $^1\text{O}_2$ in light with a known yield (Redmond and Gamlin 1999).

3.5 Measurements of plastoquinone

Total amount and the redox state of the PQ pool were measured with HPLC (Agilent; Kruk and Karpinski 2006) as described in Khorobrykh et al. (2020b), from plant leaves. Shortly, leaves were fast (~10 s) ground in ice-cold dry ethyl acetate under treatment conditions, the extract was dried under N_2 flow, dissolved to a solution of methanol and hexane and divided into two parts; the other part was immediately assayed with HPLC. All PQ in the other part was reduced by freshly made NaBH_4 (2 mM). Based on these measurements, the ratio of PQH_2 to all PQ in the sample can be calculated. To obtain fully oxidized photochemically active PQ pool as a reference, leaves were illuminated for 4 min with 690 nm light (PPFD 50 $\mu\text{mol m}^{-2} \text{s}^{-1}$), and dark-acclimated leaves were illuminated for 30 s with high light (PPFD 2000 $\mu\text{mol m}^{-2} \text{s}^{-1}$), to obtain fully reduced photochemically active PQ pool.

4 Overview of the results

4.1 Mass spectroscopy could be used to quantify singlet oxygen

A method, combining liquid chromatography with a triple quadrupole mass spectrometer (UHPLC-MS), was developed to quantify TEMPO, the reaction product of TEMP and $^1\text{O}_2$. A high concentration of TEMP (here 20 mM was used) is required to detect $^1\text{O}_2$ from photosynthetic material. Therefore, to protect the mass spectrometer, samples were diluted 50-fold prior to the analysis. As the concentration of the reaction product, TEMPO, is much smaller than that of TEMP, further dilution should be avoided. Reliable quantification of TEMPO was ensured by directing only TEMPO to mass spectrometer; otherwise the UHPLC flow (including TEMP) went to waste. With the UHPLC-MS method, TEMPO was first quantified from solutions with known concentrations of TEMP. The response of the signal to the amount of TEMP was linear (Fig. 4 in Paper I). TEMPO could be detected also from samples where $^1\text{O}_2$ was produced by illuminating Rose Bengal or isolated pumpkin (*C. maxima*) thylakoid membranes in the presence of TEMP (Fig. 4 in Paper I). The results correlated well ($R^2 = 0.83\text{--}0.99$; Fig. 7 in Paper I) with quantifications of the same samples with a traditionally used EPR method. The rate of the reaction between TEMP and $^1\text{O}_2$ was calculated based on how much the Rose Bengal solution, which absorbed 2.6 % of the light, produced TEMP. It could be then calculated that thylakoids produced $3.7 \times 10^{-7} \text{ } ^1\text{O}_2 \text{ chlorophyll}^{-1} \text{ s}^{-1}$.

The newly developed UHPLC-MS method allows fast measurements (one sample takes a bit over seven min) of multiple samples. However, long storage, even at $-75 \text{ } ^\circ\text{C}$ under an inert gas, is not advisable, as the signal decreased up to $\sim 30 \text{ } \%$ during the storage (Fig. 6 in Paper I).

4.2 Nascent oxygen produced by photosystem II was not converted to singlet oxygen

If histidine is added to a sample, it efficiently reacts with $^1\text{O}_2$ (mostly originating from gaseous O_2) and thereby decreases the O_2 concentration of the sample; the more $^1\text{O}_2$ is produced, the faster is the decrease (Telfer et al. 1994, Rehman et al. 2013;

see Materials and methods for details). Consequently, $^1\text{O}_2$ production can be measured by recording O_2 concentration in the presence and absence of histidine. Here, oxygen ($^{16}\text{O}_2$) dissolved in the buffer was replaced with a heavier isotope ($^{18}\text{O}_2$) and concentrations of the two isotopes were monitored with MIMS. With this method, it is possible to differentiate oxygen evolution ($\text{H}_2^{16}\text{O} \rightarrow ^{16}\text{O}_2$; water splitting by PSII) from oxygen consumption ($^{18}\text{O}_2 \rightarrow ^{18}\text{O}_2^{\bullet-}$; mostly due to electron donation by PSI). Illumination of isolated thylakoid membranes of pumpkin in the presence of histidine accelerated consumption of $^{18}\text{O}_2$ (compared to the control without added histidine) but did not significantly affect the rate of $^{16}\text{O}_2$ evolution (Fig. 6A; Fig. 1 in Paper II). The result indicates that the $^1\text{O}_2$ histidine reacted with derived from dissolved ambient gas ($^{18}\text{O}_2$), but the nascent $^{16}\text{O}_2$ produced by PSII was not converted to $^1\text{O}_2$ (Fig. 6B). Very similar results were obtained with two different initial oxygen ($^{16}\text{O}_2 + ^{18}\text{O}_2$) concentrations ($\sim 280 \mu\text{M}$ in Fig. 6 and $150 \mu\text{M}$ in Paper II; air-saturated concentration of oxygen at 25°C in fresh water is $253 \mu\text{M}$). In the present experimental setting, all $^{16}\text{O}_2$ could not be replaced with $^{18}\text{O}_2$; before switching on the light, 15 to 25 % of the ambient oxygen was of the lighter isotope $^{16}\text{O}_2$. Theoretically, this would lead to a small decrease ($\sim 5\%$ of the observed $^{16}\text{O}_2$ evolution) in apparent oxygen evolution in the presence of histidine, very similarly as observed (Fig. 6A; assuming that all $^1\text{O}_2$ derived from ambient dissolved gas and not from nascent oxygen).




Figure 6. Origin of singlet oxygen in photosystem II. A. Concentrations of two oxygen isotopes ($^{16}\text{O}_2$ and $^{18}\text{O}_2$) upon a high light (PPFD $3000 \mu\text{mol m}^{-2} \text{s}^{-1}$) illumination at 25°C in isolated thylakoids of pumpkin in the presence and absence of 20 mM histidine, measured with MIMS. The difference between samples with and without histidine represents production of $^1\text{O}_2$. Prior to the measurements, dissolved oxygen in the sample was removed and replaced with $^{18}\text{O}_2$. The curves are normalized to the respective oxygen concentrations at the time point zero. The data represent averages from at least three replications. B. A schematic representation showing the proposed origin of $^1\text{O}_2$; $^1\text{O}_2$, produced in the light by PSII, derives from ambient dissolved gas ($^{18}\text{O}_2$ in this case), not from nascent oxygen ($^{16}\text{O}_2$) originating from splitting of water (H_2^{16}O) by PSII.

Next, to test whether active or inactive PSII is needed for $^1\text{O}_2$ production, leaves and isolated thylakoid membranes of pumpkin were illuminated for different times to inactivate different proportion of PSII units, after which $^1\text{O}_2$ production was measured with the histidine method. In the case of leaves, thylakoid membranes were isolated immediately after the illumination treatment and prior to $^1\text{O}_2$ measurements. The results show that the capacity of PSII to evolve O_2 (or the amount of PSII photoinhibition) did not affect the rate of $^1\text{O}_2$ production (Fig. 2 in Paper II).

4.3 Photoinhibition of photosystem II

4.3.1 $\text{S}_2\text{Q}_\text{A}^-$ recombination reactions cannot be the reason for photoinhibition caused by visible light or laser pulses

To resolve the mechanism of photoinhibition of PSII in visible light, the temperature dependence of the damaging reaction at 3–40 °C was measured. In white light, the rate constant of photoinhibition, measured *in vitro* from isolated pumpkin thylakoid membranes and *in vivo* in the presence of lincomycin from pumpkin leaves and *Synechocystis* cells, slowly increased with increasing temperature (Fig. 1A in Paper III). Similar response in all the three experimental materials show that the observed positive temperature dependence is a universal feature of PSII photoinhibition.

The temperature dependence was measured also under UV radiation (UV-A, UV-B and UV-C), blue light ($\lambda < 450$ nm) and red light ($\lambda > 650$ nm). Photoinhibition depended on temperature rather similarly in all the cases, however, slightly steeper temperature dependence was observed under visible light than under the tested UV wavelengths (Fig. 1E–F in Paper III). In addition, the temperature dependence steepened towards longer wavelengths also under visible light (Fig. 1E–F in Paper III). The results indicate that more than one dominant photoinhibition mechanism functions under visible light.

$^1\text{O}_2$ produced by the slow recombination reactions between the S-states of OEC and the quinone acceptors ($\text{S}_{2/3}\text{Q}_{\text{A/B}}^- \rightarrow \text{S}_{1/2}\text{Q}_{\text{A/B}}$) have been proposed to cause photoinhibition of PSII (e.g. Vass and Cser 2009). Therefore, the rate of $^1\text{O}_2$ production in pumpkin thylakoids in white light was measured with the histidine method; the rate of $^1\text{O}_2$ production increased with temperature very similarly as the rate constant of photoinhibition (Fig. 2A in Paper III). Next, the temperature dependences of $\text{S}_2\text{Q}_\text{A}^-$ recombination reactions were measured by thermoluminescence (Fig. S4 in Paper III). Compared to photoinhibition and $^1\text{O}_2$ production, the rate constants of the $\text{S}_2\text{Q}_\text{A}^-$ recombination reactions very strongly depended on the temperature at 0–40 °C (Fig. 1C in Paper III). *In vivo*, the rate of

the charge recombination depends on the rate constant of the recombination, but also on the amount of the substrate (Q_A^-); in pumpkin leaves Q_A was slightly less reduced at higher temperatures (Fig. 1B in Paper III). However, the temperature response of the rate of the direct $S_2Q_A^-$ recombination reaction in pumpkin leaves, obtained by multiplying the rate constant by the fluorescence parameter $1-qL$ (reflecting the amount of Q_A^-), did not resemble the observed temperature dependence of photoinhibition nor that of 1O_2 production (Fig. 1C in Paper III).

Based on the fact that the efficiency of short laser pulses to damage PSII increases with increasing pulse interval (Fig. 1D in Paper III), it has been suggested that the $S_{2/3}Q_{A/B}^-$ recombination reactions cause laser pulse photoinhibition (as the longer the interval the higher the probability of the recombination; Keren et al. 1997). However, photoinhibition of PSII induced by 4 ns flashes did not have any observable temperature dependence (Fig. 1D in Paper III).

To conclude, the $S_2Q_A^-$ recombination reactions cannot be the main reason for photoinhibition of PSII or 1O_2 production.

4.3.2 Singlet oxygen did not affect photoinhibition

To understand the role of 1O_2 in photoinhibition, several experiments affecting the rate of production and quenching of 1O_2 were conducted. Quenchers of 1O_2 did not protect PSII during illumination of thylakoid membranes (Fig. 2B in Paper III). Both a water-soluble quencher (histidine) and a lipid-soluble quencher (α -tocopherol) were tested, as water-soluble quenchers are expected to react only with 1O_2 that already has escaped PSII and therefore may not be anymore very dangerous to PSII (Telfer et al. 1994, Stamatakis et al. 2016).

According to Hideg et al. (1994) 1O_2 is not produced under anaerobic conditions. However, photoinhibition proceeded under anaerobicity, and the rate was even faster than under aerobic conditions (Fig. 2C in Paper III). In addition, photoinhibition under anaerobic conditions depended on temperature and wavelength of illumination similarly as aerobic photoinhibition (Fig. 2D in Paper III). Previously it has been shown that a reciprocity between duration and intensity of illumination on the amount of PSII damage holds for anaerobic photoinhibition, as it holds for aerobic photoinhibition (Park et al. 1997, Serôdio et al. 2017). These data suggest that the same mechanism(s) causes photoinhibition of PSII both in the presence and absence of oxygen, indicating that 1O_2 cannot be the sole inducer of the damage.

Here, anaerobicity was achieved by constantly flushing the sample with nitrogen gas, and low O_2 concentrations may have persisted due to water splitting of PSII (though, in the absence of O_2 that functions as the only terminal acceptor in isolated thylakoid membranes, PSII O_2 evolution markedly diminishes). To eliminate this possibility, the experiments were repeated in the presence of 3-(3,4-dichlorophenyl)-

1,1-dimethylurea (DCMU), which binds to the Q_B site of PSII, blocking O_2 evolution. Anaerobicity did not protect PSII from photoinhibition with DCMU either (Fig. 2C in Paper III). Curiously, photoinhibition in the presence of DCMU did not have any clear temperature dependence.

4.3.3 Photoinhibition proceeded even at $-78.5\text{ }^\circ\text{C}$

To probe the role of the recombination of the primary pair ($P680^+PheO_{D1}^-$) in photoinhibition of PSII, pumpkin thylakoids were frozen and illuminated at $-78.5\text{ }^\circ\text{C}$. At this temperature, the only recombination reaction capable of producing 1O_2 is that of $P680^+PheO_{D1}^-$ (Takahashi et al. 1987). Even at $-78.5\text{ }^\circ\text{C}$, PSII was photoinhibited, both under visible light and UV radiation (Fig. 4 in Paper III). Similarly to photoinhibition at physiological temperatures, degradation of the D1 protein was induced by illumination at $-78.5\text{ }^\circ\text{C}$ and took place after melting the sample (Fig. 4B in Paper III). However, contrary to photoinhibition at physiological temperatures, anaerobicity almost completely abolished the damage to PSII (Fig. 4A in Paper III).

4.4 Redox state of the plastoquinone pool

4.4.1 F_0 rise revealed visible light wavelengths favoring photosystem II or photosystem I

When an illumination of moderate intensity ceases, chlorophyll *a* fluorescence yield decreases to a dark level (F_0), but after a few seconds a small bump, called F_0 rise, is often observed (Fig. 7A). The increase in fluorescence level has been proposed to be due to reduction of the PQ pool by the NADH dehydrogenase-like complex (Endo et al. 1997). However, fluorescence probes the redox state of Q_A , not that of the PQ pool, and the big difference in redox potentials between Q_A and PQH_2 (e.g. De Causmaecker et al. 2019) makes direct reduction of Q_A by a plastoquinol molecule an unlikely event. Accordingly, an attempt to record F_0 rise in total darkness (the measuring beam of the PAM fluorometer was switched on every five seconds, only for a short time) resulted in no rise in fluorescence level (Fig. 1 in Paper IV). If, on the top of the chopped measuring beam, low intensity "activating" light was applied, F_0 rise was restored (Fig. 7A; Fig. 1 in Paper IV). Even with activating light, F_0 rise was not observed in a mutant lacking the NADH dehydrogenase-like complex (Fig. 1B in Paper IV).




Figure 7. Activated F_0 rise in leaves of *Nicotiana tabacum*. A. The measurement protocol. Leaves were illuminated with white preillumination light (from a KL-1500 halogen illuminator), weak activating light and saturating pulses (SP; upward arrows), and chlorophyll *a* fluorescence was measured with a PAM fluorometer. Durations (by horizontal lines) and intensities of illumination are indicated. At the end of the preillumination, the measuring beam of the fluorometer was switched to a chopped mode (on 0.6 s for every five seconds), to record a post-illumination rise in the fluorescence level (F_0 rise; indicated with a downward arrow). The wavelength of the activating light is indicated next to the F_0 rise. PSII or PSI indicates whether the activating light favors PSII or PSI, respectively. B. A model explaining the wavelength-dependence of the appearance of an F_0 rise. If activating light ($h\nu$ (ex.)) excites PSII more efficiently than PSI, accumulation of PQH₂ (by components of PSI cyclic electron flow) slows down re-oxidation of Q_A⁻, thereby increasing fluorescence ($h\nu$ (fluor.)). Activating light favoring PSI, oxidizes the PQ pool, leading to fast re-oxidation of Q_A⁻ and low fluorescence. C–H. Inserts showing the post-illumination fluorescence transient. The wavelength of the activating light is indicated. The curves have been averaged from three replications.

Therefore, an F_0 rise occurs when re-oxidation of Q_A⁻ (reduced by the measuring beam of a fluorometer or by the activating light) slows down when the PQ pool is

reduced. Consequently, activated F_0 rise should be sensitive to the electron transfer balance between PSII and PSI, as activating light favoring PSII over PSI would efficiently reduce Q_A (giving rise to an F_0 rise), whereas activating light favoring PSI would suppress the F_0 rise because PSI would oxidize the PQ pool (Fig. 7B).

Next, the activated F_0 rise method was used to screen visible light wavelengths that would favor PSII or PSI; an action spectrum of F_0 rise in light-acclimated *N. tabacum* leaves was measured with monochromatic illumination for the activating light, with 10 nm intervals. Three wavelength ranges favoring PSII (460–500 nm, 560–570 nm and 650–660 nm) and four favoring PSI (420–440 nm, 520 nm, 630 nm and 680–690 nm) were found, irrespective if the PPFD of the activating light was 0.5 (data not shown), 2.5 (Fig. 2A in Paper IV) or 10 (Fig. 7) $\mu\text{mol m}^{-2} \text{s}^{-1}$. Separate measurements of fluorescence yield, P700 oxidation and 77 K fluorescence (Figs. 2, 4 and 6 in Paper IV) confirmed the PSI or PSII nature of the wavelengths.

4.4.2 The photochemically active plastoquinone pool was reduced by three and oxidized by four visible light wavelengths

As any fluorescence method, including the activated F_0 rise, probes PQ pool indirectly, PQ was next extracted from light acclimated *A. thaliana* leaves and the redox state of the photochemically active PQ pool was measured with HPLC (Fig. 8A). The photochemically active PQ (i.e. PQ molecules taking part in electron transfer from PSII to PSI; defined here as those PQ molecules that can be reduced by high light and oxidized by far-red light) comprised ~50 % of all PQ + PQH₂ of the leaves (Fig. 8B).

Illumination for four min with moderate intensity (PPFD 50 $\mu\text{mol m}^{-2} \text{s}^{-1}$) of light favoring PSII (470 nm, 560 nm or 660 nm) reduced 80–90 % of the photochemically active PQ pool (hereafter PQ pool), and illumination with 440 nm, 520 nm or 690 nm PSI light led to oxidation of 90–100 % of the pool (Fig. 3 in Paper IV). Illumination with 630 nm light reduced 40–60 % of the PQ pool (Fig. 3 in Paper IV). After three h of darkness, ~30 % of the photoactive pool PQ pool was reduced (Fig. 8B; Fig. 3 in Paper IV).

4.4.3 The lights favoring photosystem II or photosystem I induced related physiological changes

Illumination (PPFD 50 $\mu\text{mol m}^{-2} \text{s}^{-1}$) of plant leaves with wavelengths favoring PSII (560 nm or 660 nm) induced phosphorylation of LHCII and illumination with wavelengths favoring PSI (430 nm or 520 nm) induced dephosphorylation (Fig. 10B in Paper IV). Occurrence of state transitions was confirmed with 77 K fluorescence

measurements; illumination with PSII wavelengths (470 nm, 560 nm or 660 nm) led to a decrease in the absorption cross-section of PSII relative to that of PSI, and illumination with PSI wavelengths (420 nm, 520 nm or 680 nm) to an increase (Fig. 10 in Paper IV). At the beginning of the illumination 440 nm, 520 nm, 630 nm or 690 nm light oxidized P700 and PC (oxidation was persistent towards the end of the four min illumination only in 690 nm), whereas in 470 nm, 560 nm and 660 nm light both P700 and PC were almost completely reduced (Fig. 6 in Paper IV). Similarly as in the fluorescence experiments (Fig. 7F; Figs. 2, 4 and 6 in Paper IV), 630 nm light behaved as a PSI light in the PC and P700 measurements. However, 630 nm induced more LHCII phosphorylation than the 430 nm and 520 nm PSI lights, and 77 K measurements revealed state transitions that were somewhere between those induced by extreme PSII or PSI wavelengths, in accordance with the observed 40–60 % reduction of the PQ pool after 630 nm illumination (Figs. 3 and 10 in Paper IV).




Figure 8. Redox state of the plastoquinone pool in leaves of *A. thaliana* measured with high performance liquid chromatography. A. Representative chromatograms from a leaf extract, divided into two parts, one in the absence (black dashed line) and the other in the presence (grey continuous line) of NaBH_4 (used to reduce all PQ in the sample). The insert shows an enlarged view of the PQH_2 peak. B. Percentage of reduced PQ (of total PQ + PQH_2) after 30 s high light illumination (PPFD $2000 \mu\text{mol m}^{-2} \text{s}^{-1}$ on dark-acclimated leaves; HL), four min illumination with 690 nm (PPFD $50 \mu\text{mol m}^{-2} \text{s}^{-1}$ on light acclimated leaves; FR) and three h in darkness (Sample). Proportions of oxidized and reduced PQ in non-photochemically active (non-photoactive) and photochemically active (photoactive) PQ pools for the dark sample are indicated. Error bars show SD, calculated from three biological replications. C. The setting for treatments under sunlight.

4.4.4 Redox state of the plastoquinone pool in white light

The experiments so far were carried out with monochromatic light; however, natural light is always polychromatic. Therefore, the redox state of the PQ pool was measured after illumination with various sources of white light. All the white light sources used here, at the PPFD of $50 \mu\text{mol m}^{-2} \text{s}^{-1}$, oxidized more than 50 % of the photochemically active PQ pool. Next, an empirical model, based on the observed

effects of 25 monochromatic and polychromatic light sources on the redox state of the PQ pool, calculating how much a single wavelength of light affects the reduction or oxidation of the PQ pool, was developed (Fig. 7 in Paper IV). The analysis showed that green and red PSI wavelengths, in addition to far-red, often contributed significantly to oxidation of the PQ pool (Fig. 8 in Paper IV). Green and red PSII wavelengths were often equally important in reducing the PQ pool, and in natural sunlight also blue PSII wavelengths contributed significantly to the reduction (Fig. 8 in Paper IV).

In addition to the wavelength of light, intensity also affects the redox state of the PQ pool. Next, intensity responses of a PSII light (470 nm), a PSI light (440 nm) and natural sunlight (Fig. 8C) were measured. 50 % of the PQ pool was already reduced at the PPFD of $\sim 25 \mu\text{mol m}^{-2} \text{s}^{-1}$ of the PSII light, whereas the intensity response of the PSI light was much weaker; at the PPFD of $200 \mu\text{mol m}^{-2} \text{s}^{-1}$, only 30 % of the PQ was reduced (Fig. 9A in Paper IV). Similarly to all the other studied white light sources, sunlight slightly favored PSI and a gradual reduction of the PQ pool with increasing intensity, faster than in 440 nm PSI light but clearly slower than in 470 nm PSII light, was observed (Fig. 9A in Paper IV).

Finally, the rate of the change in the PQ pool redox state was measured. Alternating one min of high light (PPFD $500 \mu\text{mol m}^{-2} \text{s}^{-1}$) with four min of low light (PPFD $45 \mu\text{mol m}^{-2} \text{s}^{-1}$) resulted in a similarly alternating redox state of the PQ pool (Fig. 11A in Paper IV). Faster changes were probed with the activated F_0 rise fluorescence method. It was found that 10 s was enough to induce or remove an F_0 rise (Fig. 11B in Paper IV), suggesting that the redox state of the PQ pool can change very rapidly, in a few seconds.

4.4.5 Validation of the liquid chromatography method

Here, PQ was extracted by grinding a leaf in a mortar, which takes 10 to 15 s. To ensure that the method reflects the redox state of the PQ pool *in vivo*, the grinding was performed under treatment conditions. However, the pestle may cause a shadow during grinding, and therefore, we repeated a set of experiments with a transparent mortar, illuminated from the bottom. The results were almost identical with those obtained with the ceramic mortar (Table S1 in Paper IV), indicating that even though grinding of the whole leaf takes some time, individual chloroplasts are destroyed immediately. In addition, far-red illumination ($\lambda \geq 700$) resulted in similar oxidation of the PQ pool (Fig. 7A in Paper IV) as the 690 nm light used here as a control in most of the experiments.

5 Discussion

5.1 Improved detection methods for singlet oxygen are needed

It seems clear that photosynthetic materials produce significant amounts of $^1\text{O}_2$ (see e.g. Telfer et al. 1994, Rehman et al. 2013, Hakkila et al. 2014). Here, the rate of $^1\text{O}_2$ production by pumpkin thylakoid membranes at the PPFD of $3000 \mu\text{mol m}^{-2} \text{s}^{-1}$ was roughly 15 % of the maximum electron transfer rate in the same thylakoids (i.e. for every ~ 6 electrons transferred by PSII, a molecule of $^1\text{O}_2$ was born; Fig. 6; Papers I and II). Clearly, $^1\text{O}_2$ is an important molecule (in good and bad) for photosynthetic organisms. Unfortunately, very different $^1\text{O}_2$ yields, differing by four orders of magnitude, are presented in the literature (Paper I, Telfer et al. 1994, Tomo et al. 2012), indicating that to fully understand the role of $^1\text{O}_2$, better detection methods are still needed.

Here, the detection methods for $^1\text{O}_2$, based on TEMP and histidine, were developed further (Papers I and II). Some limitations, however, remain. For example, both methods are difficult to use for leaves *in vivo* (for the use of TEMP with leaves, see Mor et al. 2014). SOSG (Fig. 3), which has been widely used *in vivo* (e.g. Ramel et al. 2013, Ballottari et al. 2014), on the other hand, may be used only with red light (Ragás et al. 2009). Furthermore, to achieve even distribution of SOSG into leaf material may be challenging (Prasad et al. 2018).

5.2 Singlet oxygen and photoinhibition

Even though more than 60 years of research and vast amount of data, no consensus exists about the molecular mechanism of photoinhibition of PSII, mainly because none of the present hypotheses completely explain all the published results. Here, an attempt is made.

5.2.1 Photosystem II is protected against nascent oxygen

OEC lays only $\sim 20 \text{ \AA}$ away from P680 (Umena et al. 2011), possibly exposing the PSII reaction center to high O_2 concentrations. Nonetheless, $^1\text{O}_2$ measurements

with histidine combined with MIMS (Fig. 6) show that the $^1\text{O}_2$ produced by isolated plant thylakoid membranes during illumination derives from ambient dissolved O_2 , not from the nascent O_2 produced by the water splitting reactions of OEC. Of course, (practically) all atmospheric oxygen on Earth originates in O_2 evolution by PSII. The present results, however, indicate that the nascent O_2 does not immediately (in the seconds to minutes timescale of the present experiments) travel to (triplet) P680 to become converted to $^1\text{O}_2$. It should be noted that with the present signal to noise ratio, it cannot be stated with absolute certainty that none of the nascent O_2 molecules diffused to $^3\text{P680}$ and became $^1\text{O}_2$. However, clearly the majority, if not all, of the observed $^1\text{O}_2$ derived from ambient dissolved O_2 .

The results (Fig. 6 and Fig. 2 in Paper II) show that the ability to evolve O_2 *as such* does not render PSII vulnerable to oxidative damage (it may be added that the reactions leading to O_2 evolution are energetically demanding and incomplete water oxidation, for example, may induce damage and/or production of ROS as suggested e.g. by Kale et al. 2017). How is PSII reaction center protected from nascent O_2 ? Both experimental data and structural analysis suggest that channels exist in PSII to divert nascent O_2 out of OEC (as well as to bring protons out and water in), possibly into the lumen (e.g. Murray and Barber 2007, Gabdulkhakov et al. 2009, Frankel et al. 2012, Vassiliev et al. 2013). As proposed already by Anderson (2001), the channel(s) seem to efficiently divert nascent O_2 out of the reach of the PSII reaction center. The fact that nascent O_2 is not converted to $^1\text{O}_2$ would have been even more important during the evolution of oxygenic photosynthesis, when all the protective measures against $^1\text{O}_2$ of extant organisms were not yet fully functional (for discussion, see Fischer et al. 2016b, Cardona et al. 2019). Furthermore, it has been calculated that nascent O_2 quickly equilibrates with the environment (Kihara et al. 2014), further diminishing $^1\text{O}_2$ production in the early anaerobic (or microaerobic) conditions of Earth.

The rate of $^1\text{O}_2$ production did not depend on the O_2 evolution capacity of PSII (Fig. 2 in Paper II). The observation agrees with the previous measurements showing that the cumulative amount of $^1\text{O}_2$ production increases linearly with illumination time (even though the high light intensities used are expected to lead to some degree of photoinhibition of PSII) (Nishiyama et al. 2004, Alboresi et al. 2011). High light should also induce NPQ, which should decrease $^1\text{O}_2$ production (see e.g. Dall'Osto et al. 2012). Accordingly, some experiments show that the rate of $^1\text{O}_2$ production is fastest at the beginning of the illumination (Poulson and Thai 2015), but even in this case, linear increase in $^1\text{O}_2$ with time was observed at later time points.

Hideg et al. (1998), on the other hand, reported a correlation of *in vivo* $^1\text{O}_2$ production and the amount of inactive PSII units (with the D1 protein not yet degraded), up to ~60 min of illumination, after which the correlation was lost. The discrepancy with the present results may be partly explained due to increased $^1\text{O}_2$

production by lincomycin treatment (used by Hideg et al. 1998), as it has been suggested that impaired repair reactions of PSII lead to increased $^1\text{O}_2$ production (Wang et al. 2016a).

5.2.2 Photoinhibition and misses of oxygen evolving complex: an unexpected resemblance

Temperature dependences of PSII photoinhibition, $^1\text{O}_2$ production and the slow recombination reactions between OEC and the quinone acceptors of PSII were measured, in order to understand the mechanism(s) behind photoinhibition. Clearly, $\text{S}_{2/3}\text{Q}_{\text{A/B}}^- \rightarrow \text{S}_{1/2}\text{Q}_{\text{A/B}}$ recombination reactions cannot explain the slow increase from 5 to 35 °C either in the $^1\text{O}_2$ production rate or in the rate constant of photoinhibition of PSII (Figs. 1 and 2 in Paper III). Recombination of the primary pair, on the other hand, occurs even at cryogenic temperatures (Takahashi et al. 1987), and should therefore have a negligible temperature response at physiological temperatures.

On the contrary, the temperature dependence of misses of PSII, measured by Isgandarova et al. (2003) from spinach thylakoids, is almost identical to the temperature dependence of photoinhibition measured here in white light from pumpkin thylakoids (Fig. 2A in Paper III). A “miss” is a failure of a photon to induce stable charge separation and to advance the S-state of OEC. Misses are rather common (occurring usually with 5 to 10 % probability, per flash), dampening the famous period of four oscillation in O_2 evolution, induced by single turnover saturating flashes, of PSII (Kok et al. 1970). It is not clear whether misses are induced by charge recombination reactions or if the recombination occurs because an S-state of OEC fails to advance, leading to an inability of PSII to reduce P680^+ (or $\text{TyrZ}\cdot$) (e.g. de Wijn and van Gorkom 2002b, Han et al. 2012). In the latter case, misses would be an intrinsic property of OEC, possibly due to the demanding chemistry of water splitting (Han et al. 2012). The finding that the “miss-recombination” takes only 100–200 μs (Reinman and Mathis 1981, de Wijn and van Gorkom 2002a) suggests that the origin of misses lay within OEC (as water splitting with its millisecond kinetics would be outcompeted by the much faster recombination). In addition, miss frequency is not affected by the redox properties of $\text{Q}_\text{A}/\text{Q}_\text{A}^-$ pair (Han et al. 2012), but depends on the S-state of OEC, being highest in either $\text{S}_2 \rightarrow \text{S}_3$ or $\text{S}_3 \rightarrow \text{S}_0$ transition (de Wijn and van Gorkom 2002b, Han et al. 2012, Suzuki et al. 2012, Pham and Messinger 2016).

5.2.3 Long-lived P680^+ damages photosystem II

The similarity of the temperature dependences of PSII photoinhibition, $^1\text{O}_2$ production and misses obviously suggest a connection between these phenomena.

Also literature data support a connection between $^1\text{O}_2$ and photoinhibition (e.g. Jung and Kim 1990, Fischer et al. 2006, Rehman et al. 2013). Does $^1\text{O}_2$, produced mainly by the miss-recombination, inactivate PSII? The present experiments do not agree with $^1\text{O}_2$ being the major agent of the photoinhibitory damage to PSII. Quenchers of $^1\text{O}_2$ or anaerobicity did not protect PSII from damage (Fig. 2 in Paper III), in line with previous studies (e.g. Satoh 1970, Chaturvedi et al. 1992, Nishiyama et al. 2001, 2004, Hakala et al. 2005). Based on the results presented here (Fig. 6; Paper II), PSII is not expected to produce $^1\text{O}_2$ in the absence of ambient dissolved O_2 . Accordingly, Hideg et al. (1994) could not detect $^1\text{O}_2$ under anaerobic illumination of isolated spinach thylakoids.

The data presented here strongly suggest that $^1\text{O}_2$ production in thylakoids mainly originates in the recombination reaction occurring after a miss. However, if a miss occurs, but no recombination reaction follows (especially probable if the electron has already travelled to Q_B), long-lived P680^+ (or TyrZ^\bullet) is produced. P680^+ is very oxidizing and could potentially extract an electron from its surroundings, oxidizing a vital component of PSII and inactivating PSII. Here, a model is proposed where a miss induces both photoinhibition and production of $^1\text{O}_2$, however, $^1\text{O}_2$ is not the major damaging agent, but a side product of the recombination reaction that removes the dangerous P680^+ (Fig. 9A–B).

The results where increased redox potential of the $\text{Q}_\text{A}/\text{Q}_\text{A}^-$ pair coincides with decreased rates of photoinhibition and $^1\text{O}_2$ production (due to increased rate of those recombination reactions that do not produce $^1\text{O}_2$) have been explained to indicate that $^1\text{O}_2$ is the main damaging agent in photoinhibition (Fufezan et al. 2002, Vass 2012, Treves et al. 2016). However, a decrease in the redox gap between Q_A and Q_B is expected also to increase the overall rate of recombination reactions (as also observed by Roach et al. 2013 and Takegawa et al. 2019), at the expense of forward electron transfer. This would decrease the lifetime of the miss induced P680^+ , protecting PSII.

5.2.4 Direct light absorption of the manganese ions contributes to photoinhibition also in visible light

UV radiation damages PSII by direct light absorption of the Mn ions of OEC. Temperature dependences of photoinhibition caused by UV radiation and visible light resembled each other (Fig. 1F in Paper III). However, the slowly increasing activation energy of photoinhibition from UV radiation to blue light and further to red light (Fig. 2E in Paper III) suggest that direct light absorption by OEC functions also in visible light but becomes gradually less important towards longer wavelengths. Accordingly, release of Mn ions from OEC occurs faster at blue (460 nm) than in red (660 nm) light (Zavafer et al. 2017).




Figure 9. A photoinhibition hypothesis. A. When OEC of PSII is functional, forward electron transfer dominates over recombination reactions. B. When OEC is transiently inactive (a miss), the lifetime of $P680^+$ increases. A recombination reaction removes $P680^+$ but may produce 1O_2 . If the recombination does not occur (more probable if the electron has gone to Q_B), $P680^+$ lives long enough to oxidize a vital component of PSII. C. UV radiation, but also visible light (VIS) with decreasing efficiency towards longer wavelengths, can directly damage OEC. When OEC is unable to donate an electron, further damage may arise due to $P680^+$ or 1O_2 , however, a conformational change has been proposed to occur, increasing probabilities of such recombination reactions that do not repopulate $P680^+Pheo^-$ and thus do not produce 1O_2 . Damage by $P680^+$ or 1O_2 does not induce further photoinhibition, as PSII is already irreversibly inactivated, unless (in the case of 1O_2) the damage is directed to another, still functional, PSII unit.

Originally, $TyrZ^\bullet$ and $P680^+$ were suggested to cause damage in PSII centers without functional OEC (Jegerschöld et al. 1990, Chen et al. 1992, Anderson et al. 1998). After OEC is directly damaged by UV radiation or visible light, does $P680^+$ or 1O_2 , produced via a recombination reaction, cause further damage (see also Hakala et al. 2005, Ohnishi et al. 2005)? A less negative, compared to active PSII, value for the redox potential of the Q_A/Q_A^- pair has been measured in PSII units with non-functional (Cl-depleted PSII) or missing Mn-cluster (during PSII assembly or due to chemical removal; Johnson et al. 1995, Krieger and Rutherford 1997, Rova et al. 1998, Ido et al. 2011, Brinkert et al. 2016), favoring a recombination that does not produce 1O_2 (Spetea et al. 1997, Brinkert et al. 2016) and also shortening the lifetime of $P680^+$. However, 680 nm light damages the reaction center activity with high yield, if OEC has been inactivated by UV radiation prior the 680 nm illumination (Ohnishi et al. 2005). Anaerobicity slightly protects the reaction center activity of PSII, but not the water splitting activity (Chaturvedi et al. 1992), suggesting a role for a ROS in damaging PSII after (light-induced) inactivation of OEC. Electron acceptors are also able to protect PSII reaction center from photodamage (Chen et

al. 1992, Krieger and Rutherford 1997, Zavafer et al. 2015b). The data suggest that long-lived P680⁺ or ¹O₂ or both are able to damage PSII reaction center after inactivation of OEC (note that the PSII unit is already irreversibly damaged, net photoinhibition increases only if the damage targets a nearby still functional PSII unit; Fig. 9C).

5.2.5 Why do carotenoids protect photosystem II?

The photoinhibition hypothesis presented above can explain most of the results presented in the literature (protection by NPQ, evidence for involvement of the antennae, general lack of effect by ¹O₂ quenchers, anaerobic photoinhibition and the effects of Q_A redox changes). However, protection against PSII damage by carotenoids (e.g. Hakkila et al. 2014), is difficult to explain with the present model. If P680⁺ accumulates, photobleaching of a carotenoid and an accessory chlorophyll has been observed to occur (Telfer et al. 1991). However, whether cyclic electron transfer within PSII via cytochrome b559 and the β-carotene Car_{D2} is fast enough to significantly protect against photoinhibition (Thompson and Brudwig 1988, Buser et al. 1992, Barber and De Las Rivas 1993), remains to be solved. It has been proposed, however, that *C. ohadii* tolerates (and grows in) extreme high light intensities by utilizing (yet to be identified form of) cyclic electron flow within PSII (Ananyev et al. 2017).

Contrary to photoinhibition at 5–35 °C, photoinhibition at -78.5 °C was practically abolished by removal of oxygen, suggesting an involvement of ROS. Recombination of the primary pair occurs readily even at 10 K (-263 °C), producing ³P680 with 80 % yield (Takahashi et al. 1987). In a purple bacterium (*Rhodobacter sphaeroides*) a strong external magnetic field decreases ¹O₂ production by the recombination of the primary pair (Liu et al. 2005). However, in plants the rate of the damaging reaction is not slowed down in the presense of a magnetic field, even though the repair cycle appears to function faster during illumination in a strong magnetic field than in the weak ambient field (Hakala-Yatkin et al. 2011), speaking against an important role of ¹O₂ produced by the recombination of the primary pair in photoinhibition of PSII at physiological temperatures. These data suggest that ¹O₂ may be more dangerous when its diffusion is limited, as in a frozen state. Alternatively, if the amount of ¹O₂ is very high, damage may be inevitable, as observed by Hideg et al. (2007). Similarly, lethal amounts of H₂O₂ accelerated PSII damage (in addition to slowing down the repair reactions) in a *Synechococcus* species, but a H₂O₂ concentration that could be tolerated by the cells did not affect the rate of the damage to PSII (Blot et al. 2011). A protecting effect of NPQ against PSII photoinhibition was also observed in elevated O₂ but not in an N₂ atmosphere (Roach et al. 2020).

5.2.6 Sustained quenching of fluorescence may affect photoinhibition measurements

Previously, Lazarova et al. (2014) reported a temperature response of UV-B induced photoinhibition between 4 and 22 °C that was rather similar to what was observed here (Fig. 1E in Paper III). The reports about the temperature dependence of visible-light-induced photoinhibition, on the other hand, are variable. Photoinhibition was reported to decrease with decreasing temperature in pumpkin thylakoids (Tyystjärvi et al. 1994, Tyystjärvi 2013), similarly as here. Also in cyanobacteria, the rate of the damage slowed down (Ueno et al. 2016), or did not change much (Allakhverdiev and Murata 2004, Vonshak and Novoplansky 2008), with temperature. On the contrary, Tsonev and Hikosaka (2003) and Kornyejev et al. (2003) observed higher rates of damage at lower temperatures in the leaves of *Chenopodium album* and *Gossypium hirsutum*. In the latter two studies, photoinhibition was quantified by chlorophyll fluorescence and, therefore, oxygen evolution and fluorescence assays were compared here (Fig. S3A in Paper III). In pumpkin leaves, the decline in both the fluorescence parameter F_V/F_M and oxygen evolution capacity of PSII depended similarly on temperature. However, if the 30 min dark incubation after the illumination and prior to the fluorescence measurement was conducted at 5 °C, instead of 22 °C, photoinhibition appeared faster (Fig. S3B in Paper III). Tsonev and Hikosaka (2003) performed the dark incubation at the illumination temperature. Thus, it may be possible that the discrepancy between some literature and the present results originate in low-temperature-induced quenching of chlorophyll fluorescence, which does not relax during a typical dark acclimation time (usually 20 to 60 min). Indeed, a form of chlorophyll fluorescence quenching, which does not relax fast, has been observed to develop in *A. thaliana* in high light after short cold treatments (Malnoë et al. 2018). However, to reveal the molecular players behind the slow reversion of the decline in F_V/F_M in darkness (presumably due to slow relaxation of NPQ) in pumpkin leaves at low temperatures, more research would be needed.

Differences in observed temperature responses of photoinhibition in the literature may also reflect real differences in the rates of damage to PSII, as the miss frequency may depend on temperature differently in different species. Isgandarova et al. (2003) measured more misses at low than at high temperature in the thermophilic cyanobacterium *Synechococcus elongatus*. The observed low rate of photoinhibition at their optimal growth temperature in *Arthrospira platensis* (Vonshak and Novoplansky 2008) may be due to small number of misses at that temperature. On the contrary, in *Synechocystis* here, photoinhibition proceeded relatively fast at their growth temperature (Fig. 1A in Paper III).

5.3 Redox state of the plastoquinone pool senses both the imbalances between the photosystems and the intensity of sunlight

The redox state of the PQ pool regulates gene expression and photoacclimation responses (section 1.4.1), but direct measurements of the sizes of the photochemically active PQ pool and the pool of the rest of the PQ molecules or of their redox states are surprisingly rare (so far!). Here, the redox state of the PQ pool under various illumination conditions was measured with HPLC from *A. thaliana* leaves.

5.3.1 Delicate balance between the photosystems

Three wavelength ranges of visible light were found to favor PSII and four wavelength ranges were found to favor PSI (Fig. 2A in Paper IV). Also previous measurements show that PSII and PSI absorb visible light differently (Canaani and Malkin 1984, Wientjes et al. 2013b, Laisk et al. 2014). The most obvious difference between the photosystems is that PSI + LHCI binds less chlorophyll *b* than PSII + LHCII (see e.g. Caffarri et al. 2014, Mazor et al. 2017). Accordingly, the PSII wavelengths of 460–500 nm and 660 nm coincide with absorption peaks of chlorophyll *b*, and at the PSI wavelengths below 450 nm and above 680 nm chlorophyll *a* absorbs better than chlorophyll *b* (Frigaard et al. 1996, Cinque et al. 2000). Indeed, comparison of the present results with chlorophyll *b* to *a* absorption ratio in methanol (calculated based on the data from Frigaard et al. 1996) reveals peaks and deeps at very similar positions as those observed in the F_0 rise spectrum (reflecting PSII/PSI absorption; Fig. 2B in Paper IV). In the F_0 rise spectrum the peaks above 550 nm are red-shifted compared to the spectrum of chlorophyll *b* to *a* absorption ratio, most probably due to the protein environments of both pigments (Thomas and Van Der Wal 1963, Cinque et al. 2000).

Illumination at the peak and deep wavelengths of the F_0 rise spectrum almost completely reduced or oxidized, respectively, the PQ pool (Figs. 2A and 3 in Paper IV). A relatively low intensity of light (PPFD 50 $\mu\text{mol m}^{-2} \text{s}^{-1}$) was used to prevent photosynthetic control (i.e. slowing down of the electron transfer rate of the Cyt b6f complex due to lumen acidification; Tikhonov et al. 1984). 630 nm behaved as PSI light in fluorescence and absorbance measurements but reduced ~50 % of the PQ pool (Figs. 2 and 3 in Paper IV) suggesting that 630 nm light may have steeper light intensity response than the other PSI wavelengths tested here (see Fig. 9A in Paper IV).

The absorption differences between PSII and PSI were very small (Figs. 2, 4 and 5 in Paper IV), even though large effects were exerted on the PQ pool. The data indicate that in the course of evolution, absorption properties of the two

photosystems have been so well balanced that small differences such as state transitions can readjust the balance (it seems that under almost any conditions, LHCI serves as an antenna also for PSI; Wientjes et al. 2013b, Grieco et al. 2015). All polychromatic white light sources tested here, including, importantly, the Sun, oxidized the PQ pool at the PPFD of 30–50 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 9A in Paper IV). Oxidation of the PQ pool by low intensity sunlight in combination with the mild intensity response enable plants to sense the intensity of sunlight via PQ redox state.

Enrichment of far-red light in canopy shade has been believed to oxidize the PQ pool. However, low intensity white light already favors PSI, oxidizing the PQ pool (Fig. 7A in Paper IV). Accordingly, photosynthetic parameters measured from plants grown under either neutral or canopy shade were rather similar (Pons and de Jong van Berkel 2004). In addition, green PSII light is also proportionally increased under canopy (Fig. 9B in Paper IV; a spectrum of light under canopy is available, for example, also in a maize field in Hirth et al. 2013). In field grown maize, canopy light induced more phosphorylation of LHCI than artificial light enriched in far-red (Hirth et al. 2013), which may suggest that the green PSII light in canopy shade antagonizes the effects of far-red light. Indeed, green light, especially when other wavelengths of light are lacking, can be used to drive photosynthesis (Zhang et al. 2011). On the other hand, green light (as well as far-red light) induces shade avoidance signaling (for a review, see Smith et al. 2017).

5.3.2 Implications for state transitions

The redox state of the PQ pool correlated well with state transitions (quantified as the ratio of PSI to PSII fluorescence at 77 K), after short illumination with selected wavelengths favoring either PSII or PSI (Fig. 10C in Paper IV). The relationship was curvilinear, and state 2 was reached already after moderate reduction (~30 % reduction of the PQ pool was needed to induce 50 % of the maximal state transition). This is in agreement with the observation that only a single PQH₂ molecule at the Q_o site of Cyt b6f is needed to activate the STN7 kinase (Zito et al. 1999), and with the previous research where state 2 has been reached with illumination with “PSII light” which is predicted to reduce 30–80 % of the PQ pool (Table S2 and Fig. S5 in Paper IV; Tullberg et al. 2000, Pesaresi et al. 2002, Adamiec et al. 2008, Wagner et al. 2008).

Contrary to what sometimes is assumed, photochemically active PQ pool is not completely oxidized in the dark in *A. thaliana* (Kruk and Karpinski 2006, Szechyńska-Hebda et al. 2010, Yoshida et al. 2010); here, about 26 % of the PQ pool was reduced after 2 to 3 h of darkness (Fig. 8). In the dark, the PQ pool is probably reduced by NADH dehydrogenase-like complex (Burrows et al. 1998). The light state, defined through the PSI/PSII fluorescence ratio at 77 K, diligently followed

the redox state of the PQ pool also in darkness (Fig. 10C in Paper IV). The STN7 kinase is able to phosphorylate LHCII in darkness (though, for the full activity, *flux* of electrons through the Cyt b6f may be needed; Rintamäki et al. 2000, Hou et al. 2003, Tikkanen et al. 2010). Accordingly, usually (but not always; see e.g. Rintamäki et al. 2000), a small amount of phosphorylated LHCII is detected in darkness, as well as the accumulation of the “state transition complex” consisting of LHCII, LHCI and PSI (e.g. Fristedt et al. 2010, Tikkanen et al. 2010). Different LHCII phosphorylation levels observed may be due to different dark acclimation times used in the different studies, as the light history as well as the age of the plants have been shown to affect LHCII phosphorylation (Trotta et al. 2016, Schwarz et al. 2018, Longoni et al. 2019). On the contrary, during far-red illumination, STN7 is degraded (Willig et al. 2011, Trotta et al. 2016). In the light as well as in darkness, a threonine residue of STN7 is phosphorylated, which seems to protect STN7 from degradation (Trotta et al. 2016).

During low intensity illumination with monochromatic light, the phosphorylation level of LHCII linearly correlated with the extent of state transitions (Fig. 10A). When moving to white lights of different intensities, the situation becomes more complicated. Even though the phosphorylation level of LHCII decreases from low to high light (presumably due to inactivation of STN7; Rintamäki et al. 2000, possibly also due to oxidation of PQH₂ by ¹O₂; Kruk and Szymańska 2012), almost no state transitions were observed (Tikkanen et al. 2010). The lack of state transitions was observed under sunlight (Wientjes et al. 2013b) and in field-grown leaves (Hirth et al. 2013). Also here, illumination with sunlight of different intensities did not lead to different light states (Fig. 10B). However, sometimes more considerable state transitions, upon changes in light intensity, following rather nicely LHCII phosphorylation, have been reported (Pesaresi et al. 2009, Wood et al. 2018). Interestingly, here, the light state in the samples taken directly from growth chamber followed LHCII phosphorylation (Fig. 10A). It is not yet completely clear, why, despite changes in phosphorylation of LHCII and in the accumulation of PSI-LHCII, state transitions are not (sometimes) observed. It was proposed that state transitions are modulated also by PSII core phosphorylation, which under different intensities of white light mirrors LHCII phosphorylation (Tikkanen et al. 2010, Mekala et al. 2015). Also, why does STN7 phosphorylate LHCII under low light (e.g. Rintamäki et al. 2000, Tikkanen et al. 2010) when the PQ pool should be rather oxidized (Figs. 7 and 9 in Paper IV)? One possibility is that plants compensate the imbalance between electron transfer rates of PSII and PSI (see e.g. Wagner et al. 2008), which after a while is expected to lead to a more optimal oxidation state of the PQ pool. At least in mutants lacking active STN7, also STN8 (PSII core kinase) can to a degree phosphorylate LHCII (Longoni et al. 2019).

Acetylation may also be important in state transitions (Koskela et al. 2018). Clearly, the mechanistics of state transitions are not yet fully understood.




Figure 10. State transitions and phosphorylation of LHCII in *A. thaliana* leaves. A. Phosphorylation of LHCII (relative to the level of the 560 nm sample) after 8-min illumination with indicated wavelengths (PPFD $50 \mu\text{mol m}^{-2} \text{s}^{-1}$) or in growth light (GL) plotted against the ratio of PSI (735 nm) to PSII (685 nm) fluorescence, measured at 77 K from thylakoids isolated from leaves illuminated with the indicated lights for 1 h or taken directly from growth light. Data are from Fig. 10 in Paper IV. B. The ratio of PSI to PSII fluorescence, measured at 77 K from thylakoids isolated from leaves illuminated with sunlight of indicated intensity (PPFD 30, 100 or $430 \mu\text{mol m}^{-2} \text{s}^{-1}$) or far-red (FR; PPFD $50 \mu\text{mol m}^{-2} \text{s}^{-1}$) or taken directly from growth light. Fluorescence was excited with 470 nm light. Error bars show standard error of the mean (A) or SD (B) from at least three biological replications.

Expectedly, the PSI to PSII fluorescence ratio was higher when fluorescence was excited with light favoring PSI compared to excitation with a PSII light (Figs. 2 and 10D in Paper IV). Accordingly, if values of PSI to PSII fluorescence from literature are compared, the effect of the excitation wavelength should be kept in mind. In addition, the finding that a larger apparent amplitude of state transitions (i.e. larger relative difference in PSII to PSI fluorescence levels between states 1 and 2) was observed when fluorescence was excited with a PSII light, rather than with a PSI light (Fig. 10D in Paper IV, see also calculations in Paper IV), supports the view that during state 1 to 2 transition a proportion of LHCII serving PSII moves to serve PSI (e.g. Bassi et al. 1988, Iwai et al. 2008).

5.3.3 Redox state of the plastoquinone pool can and should be measured

So far direct HPLC measurements have revealed important roles for PQ at least in plants' tolerance to high light (Kruk and Szymańska 2012, Ksas et al. 2015, Ferretti et al. 2018, Pralon et al. 2019; see also section 1.2.5), heat (Pshybytko et al. 2008) and salinity (Wiczarz et al. 2018). In addition, Yoshida et al. (2010) showed that

besides the PQ pool, illumination affects the redox state of the ubiquinone pool (carriers of electrons from complex I or II to complex III in mitochondria) in *A. thaliana*. Worries about the redox state of the PQ pool possibly changing during the required extraction prior HPLC analysis, therefore not reflecting the situation *in vivo*, may be a reason for the relatively small number of published HPLC measurements. However, the redox state of the PQ pool, measured with HPLC, correlated well with redox states of other components of the photosynthetic electron transport chain (Figs. 4–6 in Paper IV) and acclimation responses known to be regulated by the redox state of the PQ pool (Fig. 10 in Paper IV). We also confirmed that grinding a leaf in a ceramic mortar did not affect the results (Table S1 in Paper IV); once in an organic solvent, PQH₂ is rather stable (Kruk and Strzałka 1999). In addition, comparison with literature shows that the method gives rather consistent results (about e.g. the size of the photoactive pool and the redox state in darkness; Kruk and Karpinski 2006, Yoshida et al. 2010, Szechyńska-Hebda et al. 2010, Ksas et al. 2015).

However, there are important points to note in order to achieve successful PQ measurements with HPLC. Extraction of the sample should be done under treatment conditions, as fast as possible and right before the HPLC analysis. Solutions of NaBH₄ (used to reduce PQH₂) cannot be stored. In addition, good controls should be performed for 100 % oxidation and reduction of the photochemically active pool (also the non- photochemically active PQ pool consists of both reduced and oxidized fractions; Fig. 8). These matters may explain why Schuurmans et al. (2014) failed to observe changes in the redox state of the PQ pool under different conditions in *Synechocystis*, contrary to Khorobrykh et al. (2020b).

Compared to HPLC, fluorescence measurements are fast and (seemingly; see e.g. Schansker et al. 2014) simple, and therefore, fluorescence methods have often been used to estimate (even stronger words are sometimes used in the literature) the redox state of the PQ pool. A fluorescence parameter developed to measure the proportion of closed reaction centers (i.e. amount of Q_A⁻), 1-qP (in general, and also particularly in this case, instead of qP, qL that takes into account connectivity between PSII units, may be preferable; Kramer et al. 2004), was found to correlate well with HPLC measurements of the redox state of PQ pool, during different time points of high light illumination after darkness (Yoshida et al. 2010). On the contrary, under monochromatic light of moderate intensity, as used here, the correlation was rather poor ($R^2 = 0.74$; Fig. S3 in Paper IV). Fluorescence parameters reflect the reduction state of Q_A, and it is not surprising that the reduction of Q_A does not necessarily follow the reduction of PQ; the difference between the redox potentials of the Q_A/Q_A⁻ and Q_B/Q_B⁻ pairs have been measured to be over 200 mV (in intact PSII), and between Q_B⁻/PQH₂ and PQ/PQH₂ ~70 mV (Golbeck and Kok 1979, Krieger et al. 1995, Brinkert et al. 2016, Kato et al. 2016, De Causmaecker et al. 2019), making it hard for an electron from PQH₂ to climb to Q_A. Absence of an F₀

rise in the absence of light (using a chopped measuring beam of the fluorometer; Fig. in Paper IV), also shows that PQH₂ does not directly reduce Q_A. In addition, the redox state of the PQ pool in darkness cannot be estimated based on these fluorescence parameters (both qP and qL are by definition 0 in the dark).

Tóth et al. (2007) developed a proxy for the redox state of the photochemically active PQ pool, based on fluorescence rise curve (called OJIP rise), which may be more reliable than qP or qL (however, if light acclimated leaves are used, or if severe PSII damage exists, control values for 100 % oxidation and reduction may not be possible to obtain). The method has not been compared, to my knowledge, with an HPLC method. Such a comparison may, however, be of importance, as the theory behind the method of Tóth et al. (2007) assumes, for example, that binding constants of PQ and PQH₂ to an empty Q_B site are the same (which is probably not the case; Zobnina et al. 2017, De Causmaecker et al. 2019).

To conclude, direct measurements of the redox state of the PQ pool are advisable, especially when the redox state is of importance for interpretation of the results.

6 Conclusions and future perspectives

For plants and all other photosynthetic organisms, light is a source of both energy and problems. To absorb light is easy but to safely convert it to useful forms of energy is not so. To split water, for example, it is required to generate the strongest of the known biological oxidants (P680⁺). How is the photosynthetic electron transport chain regulated and protected, damaged and repaired? How do plants and other photosynthetic organisms achieve the relatively high efficiencies in the environment that is all the time changing? These are fascinating topics to do research on.

In the present work, methodology was expanded (Papers I, II and IV) and then utilized to solve basic research questions in the field of photosynthesis (all Papers). One may hope that increased understanding could produce increased appreciation to our one-of-a-kind (probably) world, which is, at least for the present human population, irreplaceable and should therefore be taken great care of.

Acknowledgements

This work was conducted in the laboratory of Molecular Plant Biology. Financial support from Academy of Finland, Turku University Foundation, Doctoral Programme of Molecular Life Sciences, Väisälä Foundation, FinCEAL, The Finnish Concordia Fund, Emil Aaltonen Foundation and University of Turku is appreciated.

I thank Esa Tyystjärvi for teaching me how to do solid science and also for letting me quite freely play with whatever things I wanted to. I especially appreciate that you have always kept your door open for questions. For constructive criticism and comments, I wish to thank Saijaliisa Kangasjärvi, Yagut Allahverdiyev-Rinne and Albert Porcar-Castell, the members of my advisory committee, and the pre-examiners professor Shunichi Takahashi and assistant professor Alizée Malnoë. My sincere thanks go also to associate professor Pavel Pospíšil for agreeing to be my opponent and to Eevi Rintamäki to be the custos. Taina I would like to thank for being a very nice and supporting person. I want to thank also all my collaborators.

I have enjoyed my years with plant research. Thanks to Dimitar, Mariel, Sergey, Satu, Sujata, Susa, Tinja and all the other people I have worked with! I hope that this unit will continue to be one where help can always be found when needed. Thank you, Kurt, Tapio Mika, Anniina and all the other office and laboratory personnel, and sorry for the trouble... When asked what I liked most in my work, the coffee breaks, I once answered. For the Pharmacy coffee culture, I want to thank Daniel, Duncan, Hari, Janne, Juha, Pasi, Olli and everybody else visiting the coffee room. Thank you, Gayathri and Pooneh for sports and gossip (for serious discussions too). Special thanks to everybody organizing and attending Shitty Movie Nights.

I want to thank my family for support in all times, for fun discussions or activities and, importantly in this respect, for teaching me critical thinking since birth.

Last, I try to express my gratitude to Vesa, for peer-support during these years, for giving credit (too much, I have no doubt) to my scientific achievements and simply for being (and being wonderful).

9.11.2020

A handwritten signature in black ink, consisting of stylized, overlapping loops and a long horizontal stroke extending to the right.

List of references

- Adamiec M, Drath M, Jackowski G (2008) Redox state of plastoquinone pool regulates expression of *Arabidopsis thaliana* genes in response to elevated irradiance. *Acta Biochim Pol* 55: 161–173
- Affek HP, Yakir D (2002) Protection by isoprene against singlet oxygen in leaves. *Plant Physiol* 129: 269–277
- Alboresi A, Storti M, Cendron M, Morosinotto T (2019) Role and regulation of class-C flavodiiron proteins in photosynthetic organisms. *Biochem J* 476: 2487–2498
- Allakhverdiev S, Murata N (2004) Environmental stress inhibits the synthesis de novo of proteins involved in the photodamage-repair cycle of Photosystem II in *Synechocystis* sp. PCC 6803. *Biochim Biophys Acta Bioenerg* 1657: 23–32
- Allen JF, Mullineaux CW, Sanders CE, Melis A (1989) State transitions, photosystem stoichiometry adjustment and non-photochemical quenching in cyanobacterial cells acclimated to light absorbed by photosystem I or photosystem II. *Photosynth Res* 22: 157–166
- Allorent G, Tokutsu R, Roach T, Peers G, Cardol P, Girard-Bascou J, Seigneurin-Berny D, Petroustos D, Kuntz M, Breyton C, Franck F, Wollman FA, Niyogi KK, Krieger-Liszka A, Minagawa J, Finazzi G (2013) A dual strategy to cope with high light in *Chlamydomonas reinhardtii*. *Plant Cell* 25: 545–557
- Ananyev G, Gates C, Dismukes GC (2016) The oxygen quantum yield in diverse algae and cyanobacteria is controlled by partitioning of flux between linear and cyclic electron flow within photosystem II. *Biochim Biophys Acta Bioenerg* 1857: 1380–1391
- Ananyev G, Gates C, Kaplan A, Dismukes GC (2017) Photosystem II-cyclic electron flow powers exceptional photoprotection and record growth in the microalga *Chlorella ohadii*. *Biochim Biophys Acta Bioenerg* 1858: 873–883
- Ancin M, Fernández-San Millán A, Larraya L, Morales F, Veramendi J, Aranjuelo I, Farran I (2019) Overexpression of thioredoxin m in tobacco chloroplasts inhibits the protein kinase STN7 and alters photosynthetic performance. *J Exp Bot* 70: 1005–1016
- Anderson JM (2001) Does functional photosystem II complex have an oxygen channel? *FEBS Lett* 488: 1–4
- Anderson JM, Park YI, Chow WS (1998) Unifying model for the photoinactivation of Photosystem II in vivo under steady-state photosynthesis. *Photosynth Res* 56: 1–13
- Andersson B, Anderson JM (1980) Lateral heterogeneity in the distribution of chlorophyll-protein complexes of the thylakoid membranes of spinach chloroplasts. *Biochim Biophys Acta* 593: 427–440
- Andreou A, Feussner I (2009) Lipoxygenases – Structure and reaction mechanism. *Phytochemistry* 70: 1504–1510
- Arellano JB, Naqvi KR (2016) Endogenous singlet oxygen photosensitizers in plants. In: Nonell S, Flors C (eds) Singlet oxygen: Applications in biosciences and nanosciences, *The Royal Society of Chemistry*, pp. 239–269

- Armbruster U, Correa Galvis V, Kunz HH, Strand DD (2017) The regulation of the chloroplast proton motive force plays a key role for photosynthesis in fluctuating light. *Curr Opin Plant Biol* 37: 56–62
- Aro EM, McCaffery S, Anderson JM (1994) Recovery from photoinhibition in peas (*Pisum sativum* L.) acclimated to varying growth irradiances. Role of D1 protein turnover. *Plant Physiol* 104: 1033–1041
- Aro EM, Suorsa M, Rokka A, Allahverdiyeva Y, Paakkarinen V, Saleem A, Battchikova N, Rintamaki E (2005) Dynamics of photosystem II: a proteomic approach to thylakoid protein complexes. *J Exp Bot* 56: 347–356
- Asada K (2000) The water–water cycle as alternative photon and electron sinks. *Phil Trans R Soc Lond B* 355: 1419–1431
- Asada K, Kiso K, Yoshikawa (1974) Univalent reduction of molecular oxygen by spinach chloroplasts on illumination. *J Biol Chem* 249: 2175–2181
- Avenson TJ, Cruz JA, Kanazawa A, Kramer DM (2005) Regulating the proton budget of higher plant photosynthesis. *Proc Natl Acad Sci USA* 102: 9709–9713
- Baena-González E, Barbato R, Aro EM (1999) Role of phosphorylation in the repair cycle and oligomeric structure of photosystem II. *Planta* 208: 196–204
- Balevičius V Jr, Fox KF, Bricker WP, Jurinovich S, Prandi IG, Mennucci B, Dufy CDP (2017) Fine control of chlorophyll-carotenoid interactions defines the functionality of light-harvesting proteins in plants. *Sci Rep* 7: 13956
- Ballottari M, Alcocer MLP, D'Andrea C, Viola D, Ahn DK, Petrozza A, Polli D, Fleming GR, Cerullo G, Bassi R (2014) Regulation of photosystem I light harvesting by zeaxanthin. *Proc Natl Acad Sci USA* E2431–E2438
- Barber J, De Las Rivas J (1993) A functional model for the role of cytochrome b559 in the protection against donor and acceptor side photoinhibition. *Proc Natl Acad Sci USA* 90: 10942–10946
- Barényi B, Krause GH (1985) Inhibition of photosynthetic reactions by light. A study with isolated spinach chloroplasts. *Planta* 163: 218–226
- Barry BA, Babcock GT (1987) Tyrosine radicals are involved in the photosynthetic oxygen-evolving system. *Proc Natl Acad Sci USA* 84: 7099–7103
- Bassi R, Giacometti GM, Simpson DJ (1988) Changes in the organization of stroma membranes induced by in vivo state I-state 2 transition. *Biochim Biophys Acta* 935: 152–165
- Beisel KG, Jahnke S, Hofmann D, Köppchen S, Schurr U, Matsubara S (2010) Continuous turnover of carotenes and chlorophyll a in mature leaves of *Arabidopsis* revealed by ¹⁴C₂ pulse-chase labeling. *Plant Physiol* 152: 2188–2199
- Bellafiore S, Barneche F, Peltier G, Rochaix JD (2005) State transitions and light adaptation require chloroplast thylakoid protein kinase STN7. *Nature* 433: 892–895
- Berger H, Blifern-Klassen O, Ballottari M, Bassi R, Wobbe L, Kruse O (2014) Integration of carbon assimilation modes with photosynthetic light capture in the green alga *Chlamydomonas reinhardtii*. *Mol Plant* 7: 1545–1559
- Bethmann S, Melzer M, Schwarz N, Jahns P (2019) The zeaxanthin epoxidase is degraded along with the D1 protein during photoinhibition of photosystem II. *Plant Direct* 3: 1–13
- Betterle N, Ballottari M, Zorzan S, de Bianchi S, Cazzaniga S, Dall'Osto L, Morosinotto T, Bassi R (2009) Light-induced dissociation of an antenna hetero-oligomer is needed for non-photochemical quenching induction. *J Biol Chem* 284: 15255–15266
- Bilger W, Björkman O (1994) Relationships among violaxanthin deepoxidation, thylakoid membrane conformation, and nonphotochemical chlorophyll fluorescence quenching in leaves of cotton (*Gossypium-hirsutum* L.). *Planta* 193: 238–246
- Blot N, Mella-Flores D, Six C, Le Corguillé G, Boutte C, Peyrat A, Monnier A, Ratin M, Gourvil P, Campbell DA, Garczarek L (2011) Light history influences the response of the marine cyanobacterium *Synechococcus* sp. WH7803 to oxidative stress. *Plant Physiol* 156: 1934–1954

- Bode R, Ivanov AG, Hüner NPA (2016) Global transcriptome analyses provide evidence that chloroplast redox state contributes to intracellular as well as long-distance signalling in response to stress and acclimation in *Arabidopsis*. *Photosynth Res* 128: 287–312
- Bode S, Quentmeier CC, Liao PN, Hafi N, Barros T, Wilk L, Bittner F, Walla PJ (2009) On the regulation of photosynthesis by excitonic interactions between carotenoids and chlorophylls. *Proc Natl Acad Sci USA* 106: 12311–12316
- Borisova-Mubarakshina MM, Ivanov BN, Vetoshkina DV, Lubimov VY, Fedorchuk TP, Naydov IA, Kozuleva MA, Rudenko NN, Dall'Osto L, Cazzaniga S, Bassi R (2015) Long-term acclimatory response to excess excitation energy: Evidence for a role of hydrogen peroxide in the regulation of photosystem II antenna size. *J Exp Bot* 66: 7151–7164
- Bos P, Oosterwijk A, Koehorst R, Bader A, Philippi J, van Amerongen H, Wientjes E (2019) Digitonin-sensitive LHClI enlarges the antenna of Photosystem I in stroma lamellae of *Arabidopsis thaliana* after far-red and blue-light treatment. *Biochim Biophys Acta Bioenerg* 1860: 651–658
- Bowers PG, Porter G (1967) Quantum yields of triplet formation in solutions of chlorophyll. *Proc R Soc A* 296: 435–441
- Bowyer JR, Packer JC, McCormack BA, Whitelegge JP, Robinson C, Taylor MA (1992) Carboxyl-terminal processing of the D1 protein and photoactivation of watersplitting in photosystem II. Partial purification and characterization of the processing enzyme from *Scenedesmus obliquus* and *Pisum sativum*. *J Biol Chem* 267: 5424–5433
- Bräutigam K, Dietzel L, Kleine T, Ströher E, Wormuth D, Dietz KJ, Radke D, Wirtz M, Hell R, Dörmann P, Nunes-Nesi A, Schauer N, Fernie AR, Oliver SN, Geigenberger P, Leister D, Pfannschmidt T (2009) Dynamic plastid redox signals integrate gene expression and metabolism to induce distinct metabolic states in photosynthetic acclimation in *Arabidopsis*. *Plant Cell* 21: 2715–2732
- Brooks MD (2012) A suppressor of quenching regulates photosynthetic light harvesting. PhD dissertation, University of California, Berkeley
- Brinkert K, De Causmaecker S, Krieger-Liszkay A, Fantuzzi A, Rutherford WA (2016) Bicarbonate-induced redox tuning in Photosystem II for regulation and protection. *Proc Natl Acad Sci USA* 113: 12144–12149
- Buchert F, Mosebach L, Gäbelein P, Hippler M (2020) PGR5 is required for efficient Q cycle in the cytochrome b6f complex during cyclic electron flow. *Biochem J* 477: 1631–1650
- Burrows PA, Sazanov LA, Svab Z, Maliga P, Nixon PJ (1998) Identification of a functional respiratory complex in chloroplasts through analysis of tobacco mutants containing disrupted plastid *ndh* genes. *EMBO J* 17: 868–876
- Buser CA, Brudvig GW, Diner BA (1992) Photooxidation of cytochrome B559 in oxygen-evolving Photosystem II. *Biochemistry* 31:11449–11459
- Caffarri S, Croce R, Breton J, Bassi R (2001) The major antenna complex of Photosystem II has a xanthophyll binding site not involved in light harvesting. *J Biol Chem* 276: 35924–35933
- Caffarri S, Kouril R, Kereiche S, Boekema EJ, Croce R (2009) Functional architecture of higher plant photosystem II supercomplexes. *EMBO J* 28: 3052–3063
- Caffarri S, Tibiletti T, Jennings RC, Santabarbara S (2014) A comparison between plant Photosystem I and Photosystem II architecture and functioning. *Curr Protein Pept Sci* 15: 296–331
- Calzadilla PI, Zhan J, Sétif P, Lemaire C, Solymosi D, Battchikova N, Wang Q, Kirilovsky D (2019) the cytochrome b6f complex is not involved in cyanobacterial state transitions. *Plant Cell* 31: 911–931
- Canaani O, Malkin S (1984) Distribution of light excitation in an intact leaf between the two photosystems of photosynthesis. *Biochim Biophys Acta* 766: 513–524
- Carbonera D, Gerotto C, Posocco B, Giacometti GM, Morosinotto T (2012) NPQ activation reduces chlorophyll triplet state formation in the moss *Physcomitrella patens*. *Biochim Biophys Acta Bioenerg* 1817: 1608–1615

- Cardona T, Sánchez-Baracaldo P, Rutherford AW, Larkum AW (2019) Early Archean origin of Photosystem II. *Geobiology* 17: 127–150
- Cariti F, Chazaux M, Lefebvre-Legendre L, Longoni P, Ghysels B, Johnson X, Goldschmidt-Clermont M (2020) Regulation of light harvesting in *Chlamydomonas reinhardtii* two protein phosphatases are involved in state transitions. *Plant Physiol* 183: 1749–1764
- Cazzaniga S, Bressan M, Carbonera D, Agostini A, Dall'Osto L (2016) Differential roles of carotenoids and xanthophylls in Photosystem I Photoprotection. *Biochemistry* 55: 3636–3649
- Cazzaniga S, Dall'Osto L, Kong SG, Wada M, Bassi R (2013) Interaction between avoidance of photon absorption, excess energy dissipation and zeaxanthin synthesis against photooxidative stress in *Arabidopsis*. *Plant J* 76: 568–579
- Cazzaniga S, Li Z, Niyogi KK, Bassi R, Dall'Osto L (2012) The *Arabidopsis* szl1 mutant reveals a critical role of b-carotene in Photosystem I photoprotection. *Plant Physiol* 159: 1745–1758
- Chaturvedi R, Singh M, Sane PV (1992) Photoinactivation of photosynthetic electron transport under anaerobic and aerobic conditions in isolated thylakoids of spinach. *Z Naturforsch* 47: 63–68
- Chen GX, Kazimir J, Cheniae GM (1992) Photoinhibition of hydroxylamine-extracted photosystem II membranes: studies of the mechanism. *Biochemistry* 31: 11072–11083
- Chen T, Fluhr R (2018) Singlet oxygen plays an essential role in the root's response to osmotic stress. *Plant Physiol* 177: 1717–1727
- Chen YB, Durnford DG, Kobalzik M, Falkowski PQ (2004) Plastid regulation of lhcb1 transcription in the chlorophyte alga *Dunaliella tertiolecta*. *Plant Physiol* 136: 3737–3750
- Christopher DA, Mullet JE (1994) Separate photosensory pathways co-regulate blue light/ultraviolet-A-activated psbD-psbC transcription and light-induced D2 and CP43 degradation in barley (*Hordeum vulgare*) chloroplasts. *Plant Physiol* 104: 1119–1129
- Cinque G, Croce R, Bassi R (2000) Absorption spectra of chlorophyll a and b in Lhcb protein environment. *Photosynth Res* 64: 233–242
- Cleland RE, Melis A, Neale PJ (1986) Mechanism of photoinhibition: photochemical reaction center inactivation in system II of chloroplasts. *Photosynth Res* 9: 79–88
- Crawford T, Lehotai N, Strand Å (2018) The role of retrograde signals during plant stress responses. *J Exp Bot* 69: 2783–2795
- Croce R, van Amerongen H (2013) Light-harvesting in photosystem I. *Photosynth Res* 116: 153–166
- Cruz JA, Sacksteder CA, Kanazawa A, Kramer DM (2001) Contribution of electric field ($\Delta\psi$) to steady-state transthylakoid proton motive force (pmf) in vitro and in vivo. Control of pmf parsing into $\Delta\psi$ and ΔpH by ionic strength. *Biochemistry* 40: 1226–1237
- Cser K, Vass I (2007) Radiative and non-radiative charge recombination pathways in Photosystem II studied by thermoluminescence and chlorophyll fluorescence in the cyanobacterium *Synechocystis* 6803. *Biochim Biophys Acta Bioenerg* 1767: 233–243
- DalCorso G, Pesaresi P, Masiero S, Aseeva E, Schünemann D, Finazzi G, Joliot P, Barbato R, Leister D (2008) A complex containing PGRL1 and PGR5 is involved in the switch between linear and cyclic electron flow in *Arabidopsis*. *Cell* 132: 273–85
- Dall'Osto L, Caffarri S, Bassi R (2005) A mechanism of nonphotochemical energy dissipation, independent from PsbS, revealed by a conformational change in the antenna protein CP26. *Plant Cell* 17: 1217–1232
- Dall'Osto L, Holt NE, Kaligotla S, Fuciman M, Cazzaniga S, Carbonera D, Frank HA, Alric J, Bassi R (2012) Zeaxanthin protects plant photosynthesis by modulating chlorophyll triplet yield in specific light-harvesting antenna subunits. *J Biol Chem* 287: 41820–41834
- Davis GA, Kanazawa A, Schöttler MA, Kohzuma K, Froehlich JE, Rutherford AW, Satoh-Cruz M, Minhas D, Tietz S, Dhingra A, Kramer DM (2016) Limitations to photosynthesis by proton motive force-induced photosystem II photodamage. *eLife* 5: e16921
- De Causmaecker S, Douglass JS, Fantuzzi A, Nitschke W, Rutherford AW (2019) Energetics of the exchangeable quinone, Q(B), in Photosystem II. *Proc Natl Acad Sci USA* 116: 19458–19463

- de Lavalette ADL, Barucq L, Alric J, Rappaport F, Zito F (2009) Is the redox state of the *ci* heme of the cytochrome b6f complex dependent on the occupation and structure of the Q_i site and vice versa? *J Biol Chem* 284: 20822–20829
- de Wijn R, van Gorkom HJ (2001) Kinetics of electron transfer from QA to QB in Photosystem II. *Biochemistry* 40: 11912–11922
- de Wijn R, van Gorkom HJ (2002b) S-state dependence of the miss probability in Photosystem II. *Photosynth Res* 72: 217–222
- de Wijn R, van Gorkom HJ (2002a) The rate of charge recombination in Photosystem II. *Biochim Biophys Acta* 1553: 302–308
- Delaye L, Valadez-Cano C, Pérez-Zamorano B (2016) How really ancient is *Paulinella chromatophora*? *PLoS Curr* 8: ecurrrents.tol.e68a099364bb1a1e129a17b4e06b0c6b
- Delosme R, Olive J, Wollman FA (1996) Changes in light energy distribution upon state transitions: an in vivo photoacoustic study of the wild type and photosynthesis mutants from *Chlamydomonas reinhardtii*. *Biochim Biophys Acta* 1273: 150–158
- Depège N, Bellafiore S, Rochaix JD (2003) Role of chloroplast protein kinase Stt7 in LHCII phosphorylation and state transition in *Chlamydomonas*. *Science* 299: 157–1575
- deWinter A, Boxer SG (1999) The mechanism of triplet energy transfer from the special pair to the carotenoid in bacterial photosynthetic reaction centers. *J Phys Chem B* 103: 8786–8789
- Dexter DL (1953) Theory of sensitized luminescence in solids. *Chem Phys* 21: 836
- Di Mascio P, Martinez GR, Miyamoto S, Ronsein GE, Medeiros MHG, Cadet J (2019) Singlet molecular oxygen reactions with nucleic acids, lipids, and proteins. *Chem Rev* 119: 2043–2086
- Diner BA, Schlodder E, Nixon PJ, Coleman WJ, Rappaport F, Lavergne J, Vermaas WJM, Chisholm DA (2001) Site-directed mutations at D1-His198 and D2-His197 of Photosystem II in *Synechocystis* PCC 6803: Sites of primary charge separation and cation and triplet stabilization. *Biochemistry* 40: 9265–9281
- Drop B, Yadav KNS, Boekema EJ, Croce R (2014) Consequences of state transitions on the structural and functional organization of photosystem I in the green alga *Chlamydomonas reinhardtii*. *Plant J* 78: 181–191
- Dumas L, Zito F, Clangy S, Auroy P, Johnson X, Peltier G, Alric J (2017) A stromal region of cytochrome b6f subunit IV is involved in the activation of the Stt7 kinase in *Chlamydomonas*. *Proc Natl Acad Sci USA* 114: 12063–12068
- Durrant JR, Giorgi LB, Barber J, Klug DR, Porter G (1990) Characterization of triplet-states in isolated photosystem II reaction centres—oxygen quenching as a mechanism for photodamage. *Biochim Biophys Acta* 1017: 176–175
- Durrant JR, Hastings G, Joseph DM, Barber J, Porter G, Klug DR (1992) Subpicosecond equilibration of excitation energy in isolated photosystem II reaction centers. *Proc Natl Acad Sci USA* 89: 11632–11636
- Durrant JR, Klug DR, Kwa SL, van Grondelle R, Porter G, Dekker JP (1995) A multimer model for P680, the primary electron donor of photosystem II. *Proc Natl Acad Sci USA* 92: 4798–4802
- Endo T, Mi H, Shikanai T, Asada K (1997) Donation of electrons to plastoquinone by NADP(H) dehydrogenase and by ferredoxin-quinone reductase in spinach chloroplasts. *Plant Cell Physiol* 38: 1271–1277
- Endo T, Shikanai T, Takabayashi A, Asada K, Sato F (1999) The role of chloroplastic NAD(P)H dehydrogenase in photoprotection. *FEBS Lett* 457: 5–8
- Engel GS, Calhoun TR, Read EL, Ahn TK, Mančal T, Cheng Y-C, Blankenship RE, Fleming GR (2007) Evidence for wavelike energy transfer through quantum coherence in photosynthetic systems. *Nature* 446: 782–786
- Escoubas JM, Lomas M, LaRoche J, Falkowski PG (1995) Light intensity regulation of *cab* gene transcription is signaled by the redox state of the plastoquinone pool. *Proc Natl Acad Sci USA* 92: 10237–10241

- Faller P, Fufezan C, Rutherford AW (2005) Side-path electron donors: cytochrome b559, chlorophyll Z and β -carotene. In: Wydrzynski TJ, Satoh K (Eds), Photosystem II: The Light-Driven Water: Plastoquinone Oxidoreductase. Springer Publishers, Dordrecht, pp. 347–365
- Fan DY, Ye ZP, Wang SC, Chow W (2016) Multiple roles of oxygen in the photoinactivation and dynamic repair of Photosystem II in spinach leaves. *Photosynth Res* 127: 307–319
- Fassioli F, Dinshaw R, Arpin PC, Scholes GD (2014) Photosynthetic light harvesting: excitons and coherence. *J R Soc Interface* 11: 20130901
- Feierabend J, Dehne S (1996) Fate of the porphyrin cofactors during the light-dependent turnover of catalase and of the photosystem II reaction-center protein D1 in mature rye leaves. *Planta* 198: 413–422
- Feikema WO, Marosvölgyi MA, Lavaud J, van Gorkom HJ (2006) Cyclic electron transfer in photosystem II in the marine diatom *Phaeodactylum tricornutum*. *Biochim Biophys Acta* 1757: 829–834
- Ferretti U, Ciura J, Ksas B, Rac M, Sedlarova M, Kruk J, Havaux M, Pospisil P (2018) Chemical quenching of singlet oxygen by plastoquinols and their oxidation products in *Arabidopsis*. *Plant J* 95: 848–861
- Fischer BB, Eggen RIL, Trebst A, Krieger-Liszkay A (2006) The glutathione peroxidase homologous gene *Gpxh* in *Chlamydomonas reinhardtii* is upregulated by singlet oxygen produced in photosystem II. *Planta* 223: 583–590
- Fischer WW, Hemp J, Johnson JE (2016a) Evolution of oxygenic photosynthesis. *Annu Rev Earth Planet Sci* 44: 647–683
- Fischer WW, Hemp J, Valentine JS (2016b) How did life survive Earth's great oxygenation? *Curr Opin Chem Biol* 31: 166–178
- Flors C, Nonell S (2006) Light and singlet oxygen in plant defense against pathogens: phototoxic phenalenone phytoalexins. *Acc Chem Res* 39: 293–300
- Franck F, Juneau P, Popovic R (2002) Resolution of the Photosystem I and Photosystem II contributions to chlorophyll fluorescence of intact leaves at room temperature. *Biochim Biophys Acta* 1556: 239–246
- Frank HA, McClean MB, Sauer K (1979) Triplet states in photosystem I of spinach chloroplasts and subchloroplast particles. *Proc Nat Acad Sci USA* 76: 5124–5128
- Frankel LK, Sallans L, Limbach PA, Bricker TM (2012) Identification of oxidized amino acid residues in the vicinity of the Mn₄CaO₅ cluster of Photosystem II: Implications for the identification of oxygen channels within the photosystem. *Biochemistry* 51: 6371–6377
- Frigaard NU, Larsen KL, Cox RP (1996) Spectrochromatography of photosynthetic pigments as a fingerprinting technique for microbial phototrophs. *FEMS Microbiol Ecol* 20: 69–77
- Fristedt R, Granath P, Vener AV (2010) A protein phosphorylation threshold for functional stacking of plant photosynthetic membranes. *PLoS One* 5: e10963
- Fufezan C, Rutherford AW, Krieger-Liszkay A (2002) Singlet oxygen production in herbicide-treated Photosystem II. *FEBS Lett* 532: 407–410
- Gabdulkhakov A, Guskov A, Broser M, Kern J, Müh F, Saenger W, Zouni A (2009) Probing the accessibility of the Mn₄Ca cluster in Photosystem II: channels calculation, noble gas derivatization, and cocrystallization with DMSO. *Structure* 17: 1223–1234
- Gavelis GS, Gile GH (2018) How did cyanobacteria first embark on the path to becoming plastids?: lessons from protist symbioses. *FEMS Microbiol Lett* 365: fny209
- Giera W, Ramesh VM, Webber AN, van Stokkum I, van Grondelle R, Gibasiewicz K (2010) Effect of the P700 pre-oxidation and point mutations near A0 on the reversibility of the primary charge separation in Photosystem I from *Chlamydomonas reinhardtii*. *Biochim Biophys Acta Bioenerg* 1797: 106–112
- Girolomoni L, Ferrante P, Berteotti S, Giuliano G, Bassi R, Ballottari M (2017) The function of LHCBM4/6/8 antenna proteins in *Chlamydomonas reinhardtii*. *J Exp Bot* 68: 627–641

- Golbeck JH, Kok B (1979) Redox titration of electron acceptor Q and the plastoquinone pool in Photosystem II. *Biochim Biophys Acta Bioenerg* 547: 347–360
- González-Pérez S, Gutiérrez J, García-García F, Osuna D, Dopazo J, Lorenzo Ó, Revuelta JL, Arellano JB (2011) Early transcriptional defense responses in arabidopsis cell suspension culture under high-light conditions. *Plant Physiol* 156: 1439–1456
- Graan T, Ort DR (1984) Quantitation of the rapid electron donors to P700, the functional plastoquinone pool, and the ratio of the photosystems in spinach chloroplasts. *J Biol Chem* 259: 14003–14010
- Grieco M, Suorsa M, Jajoo A, Tikkanen M, Aro EM (2015) Light-harvesting II antenna trimers connect energetically the entire photosynthetic machinery — including both photosystems II and I. *Biochim Biophys Acta Bioenerg* 1847: 607–619
- Grieco M, Tikkanen M, Paakkari V, Kangasjarvi S, Aro EM (2012) Steady-state phosphorylation of light-harvesting complex II proteins preserves photosystem I under fluctuating white light. *Plant Physiol* 160: 1896–1910
- Groot ML, Pawlowicz NP, van Wilderen LJ, Breton J, van Stokkum IH, van Grondelle R (2005) Initial electron donor and acceptor in isolated Photosystem II reaction centers identified with femtosecond mid-IR spectroscopy. *Proc Natl Acad Sci USA* 102: 13087–13092
- Gruszka J, Pawlak A, Kruk J (2008) Tocochromanols, plastoquinol, and other biological prenyllipids as singlet oxygen quenchers – determination of singlet oxygen quenching rate constants and oxidation products. *Free Radic Biol Med* 45: 920–928
- Guergova-Kuras M, Boudreaux B, Joliet A, Joliet P, Redding K (2001) Evidence for two active branches for electron transfer in photosystem I. *Proc Natl Acad Sci USA* 98: 4437–4442
- Guskov A, Kern J, Gabdulkhakov A, Broser M, Zouni A, Saenger W (2009) Cyanobacterial photosystem II at 2.9-Å resolution and the role of quinones, lipids, channels and chloride. *Nat Struct Mol Biol* 16: 334–342
- Hakala M, Tuominen I, Keränen M, Tyystjärvi T, Tyystjärvi E (2005) Evidence for the role of the oxygen-evolving manganese complex in photoinhibition of photosystem II. *Biochim Biophys Acta* 1706: 68–80
- Hakala-Yatkin M, Mäntysaari M, Mattila H, Tyystjärvi E (2010) Contributions of visible and ultraviolet parts of sunlight to photoinhibition. *Plant Cell Physiol* 51: 1745–1753
- Hakala-Yatkin M, Sarvikas P, Paturi P, Mäntysaari M, Mattila H, Tyystjärvi T, Nedbal L, Tyystjärvi (2011) Magnetic field protects plants against high light by slowing down production of singlet oxygen. *Physiol Plant* 142: 26–34
- Hakkila K, Antal T, Rehman AU, Kurkela J, Wada H, Vass I, Tyystjärvi E, Tyystjärvi T (2014) Oxidative stress and photoinhibition can be separated in the cyanobacterium *Synechocystis* sp. PCC 6803. *Biochim Biophys Acta* 1837: 217–225
- Han G, Mamedov F, Styring S (2012) Misses during water oxidation in photosystem II are S state-dependent. *J Biol Chem* 287: 13422–13429
- Harbinson J, Foyer CH (1991) Relationships between the efficiencies of PS I and II and stromal redox state in CO₂-free air: evidence for cyclic electron flow in vivo. *Plant Physiol* 97: 41–49
- Hasan SS, Cramer WA (2012) On rate limitations of electron transfer in the photosynthetic cytochrome b6/f complex. *Phys Chem Chem Phys* 14: 13853–13860
- Hastings G, Durrant JR, Barber J, Porter G, Klug DR (1992) Observation of pheophytin reduction in photosystem two reaction centers using femtosecond transient absorption spectroscopy. *Biochemistry* 31: 7638–7647
- Havurinne V, Tyystjärvi E (2017) Action spectrum of photoinhibition in the diatom *Phaeodactylum tricornutum*. *Plant Cell Physiol* 58: 2217–2225
- Hays AMA, Vassiliev IR, Golbeck JH, Debus RJ (1998) Role of D1-His 190 in proton-coupled electron transfer reactions in photosystem II: a chemical complementation study. *Biochemistry* 37: 11352–11365
- Heimann S, Klughammer C, Schreiber U (1998) Two distinct states of the thylakoid bf complex. *FEBS Lett* 426: 126–130

- Herbstová M, Tietz S, Kinzel C, Turkina MV, Kirchoff H (2012) Architectural switch in plant photosynthetic membranes induced by light stress. *Proc Natl Acad Sci USA* 109: 20130–20135
- Hertle AP, Blunder T, Wunder T, Pesaresi P, Pribil M, Armbruster U, Leister D (2013) PGRL1 is the elusive ferredoxin-plastoquinone reductase in photosynthetic cyclic electron flow. *Mol Cell* 49: 511–523
- Hideg É, Kálai T, Hideg K, Vass I (1998) Photoinhibition of photosynthesis in vivo results in singlet oxygen production detection via nitroxide-induced fluorescence quenching in broad bean leaves. *Biochemistry* 37: 11405–11411
- Hideg É, Kós PB, Vass I (2007) Photosystem II damage induced by chemically generated singlet oxygen in tobacco leaves. *Physiol Plant* 131: 33–40
- Hideg É, Spetea C, Vass I (1994) Singlet oxygen production in thylakoid membranes during photoinhibition as detected by EPR spectroscopy. *Photosynth Res* 39: 191–199
- Hideg É, Vass I (1993) The 75°C thermoluminescence band of green tissues: Chemiluminescence from membrane-chlorophyll interaction. *Photochem Photobiol* 58: 280–283
- Hideg É, Vass I (1995) Singlet oxygen is not produced in photosystem I under photoinhibitory conditions. *Photochem Photobiol* 62: 949–952
- Hideg É, Vass I (1996) UV-B induced free radical production in plant leaves and isolated thylakoid membranes. *Plant Sci* 115: 251–260
- Hirth M, Dietzel L, Steiner S, Ludwig R, Weidenbach H, Pfalz J, Pfannschmidt T (2013) Photosynthetic acclimation responses of maize seedlings grown under artificial laboratory light gradients mimicking natural canopy conditions. *Front Sci* 4: 334
- Hofmann E, Wrench PM, Sharples FP, Hiller RG, Welte W, Diederichs K (1996) Structural basis of light harvesting by carotenoids: peridinin-chlorophyll-protein from *Amphidinium carterae*. *Science* 272: 1788–1791
- Holt NE, Zigmantas D, Valkunas L, Li XP, Niyogi KK, Fleming GR (2005) Carotenoid cation formation and the regulation of photosynthetic light harvesting. *Science* 307: 433–436
- Holzwarth AR, Müller MG, Reus M, Nowaczyk M, Sander J, Rögner M (2006) Kinetics and mechanism of electron transfer in intact photosystem II and in the isolated reaction center: pheophytin is the primary electron acceptor. *Proc Natl Acad Sci USA* 103: 6895–6900
- Hope AB, Valente P, Matthews DP (1994) Effects of pH on the kinetics of redox reactions in and around the cytochrome b_f complex in an isolated system. *Photosynth Res* 42: 111–120
- Hörtensteiner S, Kräutler B (2011) Chlorophyll breakdown in higher plants. *Biochim Biophys Acta* 1807: 977–988
- Hou HJM (2014) Unidirectional photodamage of pheophytin in photosynthesis. *Front Plant Sci* 4: 554
- Hou CX, Rintamäki E, Aro EM (2003) Ascorbate-mediated LHCII protein phosphorylation-LHCII kinase regulation in light and in darkness. *Biochemistry* 42: 5828–5836
- Houba-Herin N, Calberg-Bacq CM, Piette J, van de Vorst A (1982) Mechanisms for dye-mediated photodynamic action: Singlet oxygen production, deoxyguanosine oxidation and phage inactivation efficiencies. *Photochem Photobiol* 36: 297–306
- Huang W, Cai YF, Wang JH, Zhang SB (2018b) Chloroplastic ATP synthase plays an important role in the regulation of proton motive force in fluctuating light. *J Plant Physiol* 226: 40–47
- Huang W, Yang YJ, Zhang SB, Liu T (2018a) Cyclic electron flow around photosystem I promotes ATP synthesis possibly helping the rapid repair of photodamaged Photosystem II at low light. *Front Plant Sci* 9: 239
- Huang W, Zhang SB, Zhang JL, Hu H (2015) Photoinhibition of photosystem I under high light in the shade-established tropical tree species *Psychotria rubra*. *Front Plant Sci* 6: 801
- Ibrahim IM, Wu H, Ezhov R, Kayanja GE, Zakharov SD, Du Y, Tao WA, Pushkar Y, Cramer WA, Puthiyaveetil S (2020) An evolutionarily conserved iron-sulfur cluster underlies redox sensory function of the Chloroplast Sensor Kinase. *Commun Biol* 3: 13

- Ido K, Gross CM, Guerrero F, Sedoud A, Lai TL, Ifuku K, Rutherford AW, Krieger-Liszkay A (2011) High and low potential forms of the QA quinone electron acceptor in Photosystem II of *Thermosynechococcus elongatus* and spinach. *J Photochem Photobiol B* 104: 154–157
- Inoue K (1986) Photoinactivation sites of Photosystem I in isolated chloroplasts. *Plant Cell Physiol* 27: 961–968
- Isgandarova S, Renger G, Messinger J (2003) Functional differences of photosystem II from *Synechococcus elongatus* and spinach characterized by flash induced oxygen evolution patterns. *Biochemistry* 42: 8929–8938
- Ivanov AG, Sane PV, Hurry V, Öquist G, Huner NPA (2008) Photosystem II reaction centre quenching: mechanisms and physiological role. *Photosynth Res* 98: 565–574
- Ivanov AG, Sane PV, Krol M, Gray GR, Balseris A, Savitch LV, Öquist G, Huner NPA (2006) Acclimation to temperature and irradiance modulates PS II charge recombination. *FEBS Lett* 580: 2797–2802
- Iwai M, Takahashi Y, Minagawa J (2008) Molecular remodeling of photosystem II during state transitions in *Chlamydomonas reinhardtii*. *Plant Cell* 20: 2177–2189
- Janda T, Szalai G, Páldi E (2000) Thermoluminescence investigation of low temperature stress in maize. *Photosynthetica* 38: 635–639
- Järvi S, Suorsa M, Aro EM (2015) Photosystem II repair in plant chloroplasts — Regulation, assisting proteins and shared components with photosystem II biogenesis. *Biochim Biophys Acta* 1847: 900–909
- Järvi S, Suorsa M, Paakkarinen V, Aro EM (2011) Optimized native gel systems for separation of thylakoid protein complexes: novel super- and mega-complexes. *Biochem J* 439: 207–214
- Jegerschöld C, Virgin I, Styring S (1990) Light-dependent degradation of the D1 protein in Photosystem II is accelerated after inhibition of the water splitting reaction. *Biochemistry* 29: 6179–6185
- Jimbo H, Izuhara T, Hihara Y, Hisabori T, Nishiyama Y (2019) *Synechocystis* sp. PCC 6803 Light-inducible expression of translation factor EF-Tu during acclimation to strong light enhances the repair of photosystem II. *Proc Natl Acad Sci USA* 116: 21268–21273
- Jimbo H, Noda A, Hayashi H, Nagano T, Yumoto I, Orikasa Y, Okuyama H, Nishiyama Y (2013) Expression of a highly active catalase VktA in the cyanobacterium *Synechococcus elongatus* PCC 7942 alleviates the photoinhibition of photosystem II. *Photosynth Res* 117: 509–515
- Jimbo H, Yutthanasirikul R, Nagano T, Hisabori T, Hihara Y, Nishiyama Y (2018) Oxidation of translation factor EF-Tu inhibits the repair of Photosystem II. *Plant Physiol* 176: 2691–2699
- Johnson MP, Goral TK, Duffy CD, Brain AP, Mullineaux CW, Ruban AV (2011) Photoprotective energy dissipation involves the reorganization of photosystem II light-harvesting complexes in the grana membranes of spinach chloroplasts. *Plant Cell* 23: 1468–1679
- Johnson GN, Rutherford AW, Krieger A (1995) A change in the midpoint potential of the quinone QA in Photosystem II associated with photoactivation of oxygen evolution. *Biochim Biophys Acta Bioenerg* 1229: 202–207
- Joliot P, Joliot A (2002) Cyclic electron transfer in plant leaf. *Proc Natl Acad Sci USA* 99: 10209–10214
- Joliot P, Lavergne J, Beal D (1992) Plastoquinone compartmentation in chloroplasts. I. Evidence for domains with different rates of photo-reduction. *Biochim Biophys Acta* 1101: 1–12
- Jones LW, Kok B (1966a) Photoinhibition of chloroplast reactions. I. Kinetics and action spectra. *Plant Physiol* 41: 1037–1043
- Jones LW, Kok B (1966b) Photoinhibition of chloroplast reactions. II. Multiple effects. *Plant Physiol* 41: 1044–1049
- Jumper CC, Rafiq S, Wang S, Scholes GD (2018) From coherent to vibronic light harvesting in photosynthesis. *Curr Opin Chem Biol* 47: 39–46
- Jung J, Kim HS (1990) The chromophores as endogenous sensitizers involved in the photogeneration of singlet oxygen in spinach thylakoids. *Photochem Photobiol* 52: 1003–1009

- Junge W, Haumann H, Ahlbrink R, Mulkidjanian A, Clausen J (2002) Electrostatics and proton transfer in photosynthetic water oxidation. *Philos Trans R Soc Lond B* 357: 1407–1417
- Kale K, Hebert AE, Frankel LK, Sallans L, Bricker TM, Pospíšil P (2017) Amino acid oxidation of the D1 and D2 proteins by oxygen radicals during photoinhibition of Photosystem II. *Proc Natl Acad Sci USA* 114: 2988–29933
- Kale R, Sallans L, Frankel LK, Bricker TM (2020) Natively oxidized amino acid residues in the spinach PS I-LHC I supercomplex. *Photosynth Res* 143: 263–273
- Kambakam S, Bhattacharjee U, Petrich J, Rodermel S (2016) PTOX mediates novel pathways of electron transport in etioplasts of Arabidopsis. *Mol Plant* 9: 1240–1259
- Kanazawa A, Kramer DM (2002) In vivo modulation of nonphotochemical exciton quenching (NPQ) by regulation of the chloroplast ATP synthase. *Proc Natl Acad Sci USA* 99: 12789–12794
- Kapri-Pardes E, Naveh L, Adam Z (2007) The thylakoid lumen protease Deg1 is involved in the repair of photosystem II from photoinhibition in Arabidopsis. *Plant Cell* 19: 1039–1047
- Kargul J, Turkina MV, Nield J, Benson S, Vener AV, Barber J (2005) Light-harvesting complex II protein CP29 binds to photosystem I of *Chlamydomonas reinhardtii* under state 2 conditions. *FEBS J* 272: 4797–4806
- Kasahara M, Kagawa T, Oikawa K, Suetsugu N, Miyao M, Wada M (2002) Chloroplast avoidance movement reduces photodamage in plants. *Nature* 420: 829–832
- Kashiyama Y, Yokoyama A, Kinoshita Y, Shoji S, Miyashita H, Shiratori T, Suga H, Ishikawa K, Ishikawa A, Inouye I, Ishida K, Fujinuma D, Aoki K, Kobayashi M, Nomoto S, Mizoguchi T, Tamiaki H (2012) Ubiquity and quantitative significance of detoxification catabolism of chlorophyll associated with protistan herbivory. *Proc Natl Acad Sci USA* 109: 17328–17335
- Kato Y, Hyodo K, Sakamoto W (2018) The photosystem II repair cycle requires FtsH turnover through the EngA GTPase. *Plant Physiol* 178: 596–611
- Kato Y, Nagao R, Noguchi T (2016) Redox potential of the terminal quinone electron acceptor QB in photosystem II reveals the mechanism of electron transfer regulation. *Proc Natl Acad Sci USA* 113: 620–625
- Kato Y, Ohira A, Nagao R, Noguchi T (2019) Does the water-oxidizing Mn₄CaO₅ cluster regulate the redox potential of the primary quinone electron acceptor QA in photosystem II? A study by fourier transform infrared spectroelectrochemistry. *Biochim Biophys Acta Bioenerg* 1860: 148082
- Keren N, Berg A, Van Kan PJM, Levanon H, Ohad I (1997) Mechanism of photosystem II photoinactivation and D1 protein degradation at low light: The role of back electron flow. *Proc Natl Acad Sci USA* 94: 1579–1584
- Kern J et al. (2018) Structures of the intermediates of Kok's photosynthetic water oxidation clock. *Nature* 563: 421–425
- Khatoon M, Inagawa K, Pospíšil P, Yamashita A, Yoshioka M, Lundin B, Horie J, Morita N, Jajoo A, Yamamoto Y, Yamamoto Y (2009) Quality control of photosystem II: thylakoid unstacking is necessary to avoid further damage to the D1 protein and to facilitate D1 degradation under light stress in spinach thylakoids. *J Biol Chem* 284: 25343–25352
- Khorobrykh S, Tsurumaki T, Tanaka K, Tyystjärvi T, Tyystjärvi E (2020b) Measurement of the redox state of the plastoquinone pool in cyanobacteria. *FEBS Lett* 594: 367–375
- Khorobrykh SA, Havurinne V, Mattila H, Tyystjärvi E (2020a) Oxygen and ROS in photosynthesis. *Plants* 9: 91
- Khorobrykh SA, Khorobrykh AA, Yanykin DV, Ivanov BN, Klimov VV, Mano J (2011) Photoproduction of catalase-insensitive peroxides on the donor side of manganese-depleted Photosystem II: Evidence with a specific fluorescent probe. *Biochemistry* 50: 10658–10665
- Khorobrykh SA, M Karonen, E Tyystjärvi (2015) Experimental evidence suggesting that H₂O₂ is produced within the thylakoid membrane in a reaction between plastoquinol and singlet oxygen. *FEBS Lett* 589: 779–786
- Kihara S, Hartzler DA, Savikhin S (2014) Oxygen concentration inside a functioning photosynthetic cell. *Biophys J* 106: 1882–1889

- Kirilovsky D (2015) Modulating energy arriving at photochemical reaction centers: orange carotenoid protein-related photoprotection and state transitions. *Photosynth Res* 126: 3–17
- Klauss A, Haumann M, Dau H (2012) Alternating electron and proton transfer steps in photosynthetic water oxidation. *Proc Natl Acad Sci USA* 109: 16035–16040
- Knaff DB, Arnon DI (1969) Light-induced oxidation of a chloroplast b-type cytochrome at -189 °C. *Proc Natl Acad Sci USA* 63: 956–962
- Knopf RR, Adam Z (2018) Lumenal exposed regions of the D1 protein of PSII are long enough to be degraded by the chloroplast DegI protease. *Sci Rep* 8: 5230
- Kodru S, ur Rehman A, Vass I (2020) Chloramphenicol enhances Photosystem II photodamage in intact cells of the cyanobacterium *Synechocystis* PCC 6803. *Photosynth Res* 145: 227–235
- Koivuniemi A, Aro EM, Andersson B (1995) Degradation of the D1- and D2-proteins of photosystem II in higher plants is regulated by reversible phosphorylation. *Biochemistry* 34: 16022–16029
- Kojima K, Motohashi K, Morota T, Oshita M, Hisabori T, Hayashi H, Nishiyama Y (2009) Regulation of translation by the redox state of elongation factor G in the cyanobacterium *Synechocystis* sp. PCC 6803. *J Biol Chem* 284: 18685–18691
- Kok B, Forbush B, McGloin M (1970) Cooperation of charges in photosynthetic O₂ evolution. I. A linear four step mechanism. *Photochem Photobiol* 11: 457–475
- Kornyeyev D, Holaday S, Logan B (2003) Predicting the extent of Photosystem II photoinactivation using chlorophyll a fluorescence parameters measured during illumination. *Plant Cell Physiol* 44: 1064–1070
- Kós PB, Deák Z, Cheregi O, Vass I (2008) Differential regulation of *psbA* and *psbD* gene expression, and the role of the different D1 protein copies in the cyanobacterium *Thermosynechococcus elongatus* BP-1. *Biochim Biophys Acta* 1777: 74–83
- Koskela MM, Brünje A, Ivanauskaite A, Grabsztunowicz M, Lassowskat I, Neumann U, Dinh TV, Sindlinger J, Schwarzer D, Wirtz M, Tyystjärvi E, Finkemeier I, Mulo P (2018) Chloroplast acetyltransferase NSI is required for state transitions in *Arabidopsis thaliana*. *Plant Cell* 30: 1695–1709
- Kramer DM, Johnson G, Kiirats O, Edwards GE (2004) New fluorescence parameters for the determination of QA redox state and excitation energy fluxes. *Photosynth Res* 79: 209–218
- Krasnovsky Jr AA (1994) Singlet molecular oxygen: mechanisms of formation and deactivation pathways in the photosynthetic systems. *Biofizika* 39: 236–250
- Krasnovsky Jr AA, Cheng P, Blankenship RE, Moore TA, Gust D (1993) The photophysics of monomeric bacteriochlorophylls-C and bacteriochlorophylls-D and their derivatives - properties of the triplet-state and singlet oxygen photogeneration and quenching. *Photochem Photobiol* 57: 324–330
- Kreisbeck C, Aspuru-Guzik A (2016) Efficiency of energy funneling in the photosystem II supercomplex of higher plants. *Chem Sci* 7: 4174–4183
- Krieger A, Rutherford AW (1997) Comparison of chloride-depleted and calcium-depleted PSII: themidpoint potential of Q(A) and susceptibility to photodamage. *Biochim Biophys Acta* 1319: 91–98
- Krieger A, Rutherford AW, Johnson GN (1995) On the determination of redox midpoint potential of the primary quinone electron acceptor, QA, in photosystem II. *Biochim Biophys Acta* 1229: 193–201
- Kruk J, Hollander-Czytko H, Oettmeier W, Trebst A (2005) Tocopherol as singlet oxygen scavenger in photosystem II. *J Plant Physiol* 162: 749–757
- Kruk J, Karpinski S (2006) An HPLC-based method of estimation of the total redox state of plastoquinone in chloroplasts, the size of the photochemically active plastoquinone-pool and its redox state in thylakoids of *Arabidopsis*. *Biochim Biophys Acta* 1757: 1669–1675
- Kruk J, Strzałka K (1999) Dark reoxidation of the plastoquinone-pool is mediated by the low-potential form of cytochrome b-559 in spinach thylakoids. *Photosynth Res* 62: 273–279

- Kruk J, Szymańska R (2012) Singlet oxygen and non-photochemical quenching contribute to oxidation of the plastoquinone-pool under high light stress in Arabidopsis. *Biochim Biophys Acta* 1817: 705–710
- Kruk J, Trebst A (2008) Plastoquinol as a singlet oxygen scavenger in Photosystem II. *Biochim Biophys Acta* 1777: 154–162
- Krynická V, Shao S, Nixon PJ, Komenda J (2015) Accessibility controls selective degradation of photosystem II subunits by FtsH protease. *Nat Plants* 1: 15168
- Ksas B, Becuwe N, Chevalier A, Havaux M (2015) Plant tolerance to excess light energy and photooxidative damage relies on plastoquinone biosynthesis. *Sci Rep* 5: 10919
- Kupitz C et al. (2014) Serial time-resolved crystallography of photosystem II using a femtosecond X-ray laser. *Nature* 513: 261–265
- Kyle DJ, Ohad I, Arntzen CJ (1984) Membrane protein damage and repair: Selective loss of a quinone-protein function in chloroplast membranes. *Proc Natl Acad Sci USA* 81: 4070–4074
- Laisk A, Eichelmann H, Oja V, Peterson RB (2005) Control of cytochrome b6 f at low and high light intensity and cyclic electron transport in leaves. *Biochim Biophys Acta* 1708: 79–90
- Laisk A, Oja V, Eichelmann H (2016) Kinetics of plastoquinol oxidation by the Q-cycle in leaves. *Biochim Biophys Acta* 1857: 819–830
- Laisk A, Oja V, Eichelmann H, Dall'Osto L (2014) Action spectra of photosystems II and I and quantum yield of photosynthesis in leaves in State I. *Biochim Biophys Acta* 1837: 315–325
- Lambrev PH, Miloslavina Y, Jahns P, Holzwarth AE (2012) On the relationship between non-photochemical quenching and photoprotection of Photosystem II. *Biochim Biophys Acta* 1817: 760–769
- Lavergne J (1983) Membrane potential-dependent reduction of cytochrome b-6 in an algal mutant lacking Photosystem I centers. *Biochim Biophys Acta Bioenerg* 725: 25–33
- Lazarova D, Stanoeva D, Popova A, Vasilev D, Velitchkova M (2014) UV-B induced alteration of oxygen evolving reactions in pea thylakoid membranes as affected by scavengers of reactive oxygen species. *Biol Plantarum* 58: 319–327
- Lee PC, Rodgers MAJ (1983) Singlet molecular oxygen in micellar systems. 1. Distribution equilibria between hydrophobic and hydrophilic compartments. *J Phys Chem* 87: 4894–4898
- Li L, Aro EM, Millar AH (2018) Mechanisms of photodamage and protein turnover in photoinhibition. *Trends Plant Sci* 23: 667–676
- Li Q, Yao ZJ, Mi H, (2016) Alleviation of photoinhibition by co-ordination of chlororespiration and cyclic electron flow mediated by NDH under heat stressed condition in tobacco. *Front Plant Sci* 7: 285
- Li XP, Björkman O, Shih C, Grossman AR, Rosenquist M, Jansson S, Niyogi KK (2000) A pigment-binding protein essential for regulation of photosynthetic light harvesting. *Nature* 403: 391–395
- Lima-Melo Y, Collan PJ, Tikkanen M, Silveira JAG, Aro EM (2019) Consequences of photosystem-I damage and repair on photosynthesis and carbon use in Arabidopsis thaliana. *Plant J* 97: 1061–1072
- Lion Y, Delmelle M, van de Vorst A (1976) New method of detecting singlet oxygen production. *Nature* 263: 442–443
- Liu Y, Edge R, Henbest K, Timmel CR, Hore PJ, Gast P (2005) Magnetic field effect on singlet oxygen production in a biochemical system. *Chem Commun* 174–176
- Longoni P, Samol I, Goldschmidt-Clermont M (2019) The kinase STATE TRANSITION 8 phosphorylates light harvesting complex ii and contributes to light acclimation in Arabidopsis thaliana. *Front Plant Sci* 10: 1156
- Ma F, Chen XB, Sang M, Wang P, Zhang JP, Li LB, Kuang TY (2009) Singlet oxygen formation and chlorophyll a triplet excited state deactivation in the cytochrome b6f complex from Bryopsis corticulans. *Photosynth Res* 100: 19–28
- Majer P, Neugart S, Krumbein A, Schreiner M, Hideg É (2014) Singlet oxygen scavenging by leaf flavonoids contributes to sunlight acclimation in *Tilia platyphyllos*. *Environ Exper Bot* 100: 1–9

- Malnoë A, Schultink A, Shahrasbi S, Rumeau D, Havaux M, Niyogi KK (2018) The plastid lipocalin LCNP is required for sustained photoprotective energy dissipation in *Arabidopsis*. *Plant Cell* 30: 196–208
- Malone LA, Qian P, Mayneord GE, Hitchcock A, Farmer DA, Thompson RF, Swainsbury DJK, Ranson NA, Hunter CN, Johnson MP (2019) Cryo-EM structure of the spinach cytochrome b6 f complex at 3.6 Å resolution. *Nature* 575: 535–539
- Mao HB, Li GF, Ruan X, Wu QY, Gong YD, Zhang XF, Zhao NM (2002) The redox state of plastoquinone pool regulates state transitions via cytochrome b6f complex in *Synechocystis* sp. PCC 6803. *FEBS Lett* 519: 82–86
- Marin B, Nowack ECM, Melkonian M (2005) Plastid in the making: Evidence for a second primary endosymbiosis. *Protist* 156: 425–432
- Martin WF, Bryant DA, Beatty JT (2018) A physiological perspective on the origin and evolution of photosynthesis. *FEMS Microbiol Rev* 42: 205–231
- Martínez-Junza V, Szczepaniak M, Braslavsky SE, Sander J, Nowaczyk M, Rögner M, Holzwarth A (2008) A photoprotection mechanism involving the D2 branch in photosystem II cores with closed reaction centers. *Photochem Photobiol Sci* 7: 1337–1343
- Mattila H, Khorobrykh S, Havurinne V, Tyystjärvi E (2015) Reactive oxygen species: Reactions and detection from photosynthetic tissues. *J Photochem Photobiol B* 152: 176–214
- Maxwell K, Johnson GN (2000) Chlorophyll fluorescence – a practical guide. *J Exp Bot* 51: 659–668
- Mazor Y, Borovikova A, Caspy I, Nelson N (2017) Structure of the plant photosystem I supercomplex at 2.6 Å resolution. *Nat Plants* 3: 17014
- Mccollom TM, Shock EL (1997) Geochemical constraints on chemolithoautotrophic metabolism by microorganisms in seafloor hydrothermal systems. *Geochim Cosmochim Acta* 61: 4375–4391
- Mekala NR, Suorsa M, Rantala M, Aro EM, Tikkanen M (2015) Plants actively avoid state transitions upon changes in light intensity: role of light-harvesting complex II protein dephosphorylation in high light. *Plant Physiol* 168: 721–734
- Mereschkowsky C (1905) Über natur und ursprung der chromatophoren im pflanzenreiche. *Biol Cent* 25: 593–604
- Merkel PB, Kearns DR (1972) Radiationless decay of singlet molecular oxygen in solution. An experimental and theoretical study of electronic-to-vibrational energy transfer. *J Am Chem Soc* 94: 7244–7253
- Meyer B, Schlodder E, Dekker JP, Witt HT (1989) O₂ evolution and Chl a+II (P-680+) nanosecond reduction kinetics in single flashes as a function of pH. *Biochim Biophys Acta Bioenerg* 974: 36–43
- Mitchell P (1979) Possible molecular mechanisms of the protonmotive function of cytochrome systems. *J Theor Biol* 62: 327–367
- Miyake C, Okamura M (2003) Cyclic electron flow within PSII protects PSII from its photoinhibition in thylakoid membranes from spinach chloroplasts. *Plant Cell Physiol* 44: 457–462
- Miyamoto S, Ronsein GE, Prado FM, Uemi M, Corrêa TC, Toma IN, Bertolucci A, Oliveira MCB, Motta FD, Medeiros MHG, Di Mascio P (2007) Biological hydroperoxides and singlet molecular oxygen generation. *IUBMB Life* 59: 322–331
- Moan J (1990) On the diffusion length of singlet oxygen in cells and tissues. *J Photochem Photobiol B* 6: 343–347
- Mohamed A, Jansson C (1989) Influence of light on accumulation of photosynthesis-specific transcripts in the cyanobacterium *Synechocystis* 6803. *Plant Mol Biol* 13: 693–700
- Moore KR, Magnabosco C, Momper L, Gold DA, Bosak T, Fournier GP (2019) An expanded ribosomal phylogeny of Cyanobacteria supports a deep placement of plastids. *Front Microbiol* 10: 1612
- Mor A, Koh E, Weiner L, Rosenwasser S, Sibony-Benjamini H, Fluhr R (2014) Singlet oxygen signatures are detected independent of light or chloroplasts in response to multiple stresses. *Plant Physiol* 165: 249–261

- Mozzo M, Dall'Osto L, Hienerwadel R, Bassi R, Croce R (2008) Photoprotection in the Antenna Complexes of Photosystem II. Role of individual xanthophylls in chlorophyll triplet quenching. *J Biol Chem* 283: 6184–6192
- Mulkidjanian AY (2010) Activated Q-cycle as a common mechanism for cytochrome bcl and cytochrome b6f complexes. *Biochim Biophys Acta* 1797: 1858–1868
- Müller MG, Lambrev P, Reus M, Wientjes E, Croce R, Holzwarth AR (2010) Singlet energy dissipation in the photosystem ii light-harvesting complex does not involve energy transfer to carotenoids. *ChemPhysChem* 11: 1289–1296
- Müller MG, Niklas J, Lubitz W, Holzwarth AR (2003) Ultrafast transient absorption studies on Photosystem I reaction centers from *Chlamydomonas reinhardtii*. 1. A new interpretation of the energy trapping and early electron transfer steps in Photosystem I. *Biophys J* 85: 3899–3922
- Mullineaux CW, Allen JF (1990) State 1-state 2 transitions in the cyanobacterium *Synechococcus* 6301 are controlled by the redox state of electron carriers between Photosystems I and II. *Photosynth Res* 23: 297–311
- Mulo P, Medina M (2017) Interaction and electron transfer between ferredoxin–NADP⁺ oxidoreductase and its partners: structural, functional, and physiological implications. *Photosynth Res* 134: 265–280
- Mulo P, Pursiheimo S, Hou CX, Tyystjärvi T, Aro EM (2003) Multiple effects of antibiotics on chloroplast and nuclear gene expression. *Funct Plant Biol* 30: 1097–1103
- Munekage Y, Hashimoto M, Miyake C, Tomizawa K, Endo T, Tasaka M, Shikanai T (2004) Cyclic electron flow around photosystem I is essential for photosynthesis. *Nature* 429: 579–582
- Munekage YN, Hojo M, Meurer J, Endo T, Tasaka M, Shikanai T (2002) PGR5 is involved in cyclic electron flow around photosystem I and is essential for photoprotection in *Arabidopsis*. *Cell* 110: 361–371
- Mur LA, Aubry S, Mondhe M, Kingston-Smith A, Gallagher J, Timms-Taravella E, James C, Papp I, Hörtensteiner S, Thomas H, Ougham H (2010) Accumulation of chlorophyll catabolites photosensitizes the hypersensitive response elicited by *Pseudomonas syringae* in *Arabidopsis*. *New Phytol* 188: 161–174
- Murata N, Nishiyama Y (2018) ATP is a driving force in the repair of photosystem II during photoinhibition. *Plant Cell Environ* 41: 285–299
- Murray JW, Barber J (2007) Structural characteristics of channels and pathways in photosystem II including the identification of an oxygen channel. *J Struct Biol* 159: 228–237
- Nagao R, Yamaguchi M, Nakamura S, Ueoka-Nakanishi H, Noguchi T (2017) Genetically introduced hydrogen bond interactions reveal an asymmetric charge distribution on the radical cation of the special-pair chlorophyll P680. *J Biol Chem* 292: 7474–7486
- Nagarajan A, Burnap RL (2014) Parallel expression of alternate forms of psbA2 gene provides evidence for the existence of a targeted D1 repair mechanism in *Synechocystis* sp. PCC 6803. *Biochim Biophys Acta* 1837: 1417–1426
- Nagy G, Unnep R, Zsiros O, Tokutsu R, Takizawa K, Porcar L, Moyet L, Petroutsos D, Garab G, Finazzi G, Minagawa J (2014) Chloroplast remodeling during state transitions in *Chlamydomonas reinhardtii* as revealed by noninvasive techniques in vivo. *Proc Natl Acad Sci USA* 111: 5042–5047
- Nakamura S, Hidema J, Sakamoto W, Ishida H, Izumi M (2018) Selective elimination of membrane-damaged chloroplasts via microautophagy. *Plant Physiol* 177: 1007–1026
- Nawrocki W, Tourasse NJ, Taly A, Rappaport F, Wollman FA (2015) The plastid terminal oxidase: its elusive function points to multiple contributions to plastid physiology. *Annu Rev Plant Biol* 66: 49–74
- Nishiyama Y, Allakhverdiev SI, Yamamoto H, Hayashi H, Murata N (2004) Singlet oxygen inhibits the repair of photosystem II by suppressing the translation elongation of the D1 protein in *Synechocystis* sp. PCC 6803. *Biochemistry* 43: 11321–11330

- Nishiyama Y, Murata N (2014) Revised scheme for the mechanism of photoinhibition and its application to enhance the abiotic stress tolerance of the photosynthetic machinery. *Appl Microbiol Biotechnol* 98: 8777–8796
- Nishiyama Y, Yamamoto H, Allakhverdiev SI, Inaba M, Yokota A, Murata N (2001) Oxidative stress inhibits the repair of photodamage to the photosynthetic machinery. *EMBO J* 20: 5587–5594
- Noguchi T, Tomo T, Kato C (2001) Triplet formation on a monomeric chlorophyll in the Photosystem II reaction center as studied by time-resolved infrared spectroscopy. *Biochemistry* 40: 2176–2185
- Novoderezhkin VI, Dekker JP, van Grondelle R (2007) Mixing of exciton and charge-transfer states in photosystem II reaction centers: modeling of Stark spectra with modified Redfield theory. *Biophys J* 93: 1293–1311
- Ohad I, Adir N, Koike H, Kyle DJ, Inoue Y (1990) Mechanism of photoinhibition in vivo. A reversible light-induced conformational change of reaction center II is related to an irreversible modification of the D1 protein. *J Biol Chem* 265: 1972–1979
- Ohad I, Kyle DJ, Arntzen CJ (1984) Membrane protein damage and repair: Removal and replacement of inactivated 32-kilodalton polypeptides in chloroplast membranes. *J Cell Biol* 270: 14919–14927
- Ohnishi N, Allakhverdiev SI, Takahashi S, Higashi S, Watanabe M, Nishiyama Y, Murata N (2005) Two-step mechanism of photodamage to Photosystem II: Step 1 occurs at the oxygen-evolving complex and step 2 occurs at the photochemical reaction center. *Biochemistry* 44: 8494–8499
- Okada K, Satoh K, Katoh S (1991) Chloramphenicol is an inhibitor of photosynthesis. *FEBS Lett* 295: 155–158
- Orf GS, Gisriel C, Redding KE (2018) Evolution of photosynthetic reaction centers: insights from the structure of the heliobacterial reaction center. *Photosynth Res* 138: 11–37
- Pathak V, Prasad A, Pospíšil P (2017) Formation of singlet oxygen by decomposition of protein hydroperoxide in photosystem II. *PLoS One* 12: e0181732
- Pavlou A, Jacques J, Ahmadova N, Mamedov F, Styryng S (2018) The wavelength of the incident light determines the primary charge separation pathway in Photosystem II. *Sci Rep* 8: 2837
- Peers G, Truong TB, Ostendorf E, Busch A, Elrad D, Grossman AR, Hippler M, Niyogi KK (2009) An ancient light-harvesting protein is critical for the regulation of algal photosynthesis. *Nature* 462: 518–521
- Pesaresi P, Hertle A, Pribil M, Kleine T, Wagner R, Strissel H, Ihnatowicz A, Bonardi V, Scharfenberg M, Schneider A, Pfannschmidt T, Leister D (2009) Arabidopsis STN7 kinase provides a link between short- and long-term photosynthetic acclimation. *Plant Cell* 21: 2402–2423
- Pesaresi P, Lunde C, Jahns P, Tarantino D, Meurer J, Varotto C, Hirtz RD, Soave C, Scheller H, Salamini F, Leister D (2002) A stable LHCII-PSI aggregate and suppression of photosynthetic state transitions in the psae1-1 mutant of *Arabidopsis thaliana*. *Planta* 215: 940–948
- Pescheck F, Lohbeck KT, Roleda MY, Bilger W (2014) UVB-induced DNA and photosystem II damage in two intertidal green macroalgae: Distinct survival strategies in UV-screening and non-screening Chlorophyta. *J Photochem Photobiol B* 132: 85–93
- Peterman EJ, Dukker FM, van Grondelle R, van Amerongen H (1995) Chlorophyll a and carotenoid triplet states in light-harvesting complex II of higher plants. *Biophys J* 69: 2670–2678
- Petersen J, Förster K, Turina P, Gräber P (2012) Comparison of the H⁺/ATP ratios of the H⁺-ATP synthases from yeast and from chloroplast. *Proc Natl Acad Sci USA* 109: 11150–11155
- Petrillo E, Godoy Herz MA, Fuchs A, Reifer, Fuller J, Yanovsky MJ, Simpson C, Brown JWS, Barta A, Kalyna M, Kornblihtt AR (2014) A chloroplast retrograde signal regulates nuclear alternative splicing. *Science* 344: 427–430
- Pham LV, Messinger J (2016) Probing S-state advancements and recombination pathways in photosystem II with a global fit program for flash-induced oxygen evolution pattern. *Biochim Biophys Acta* 1857: 848–859
- Pietrzykowska M, Suorsa M, Semchonok DA, Tikkanen M, Boekema EJ, Aro EM, Jansson S (2014) The light-harvesting chlorophyll a/b binding proteins Lhcb1 and Lhcb2 play complementary roles during state transitions in *Arabidopsis*. *Plant Cell* 26: 3646–3660

- Piippo M, Allahverdiyeva Y, Paakkari V, Suoranta UM, Battchikova N, Aro EM (2006) Chloroplast-mediated regulation of nuclear genes in *Arabidopsis thaliana* in the absence of light stress. *Physiol Genomics* 25: 142–152
- Pinnola A, Bassi R (2018) Molecular mechanisms involved in plant photoprotection. *Biochem Soc Trans* 46: 467–482
- Pons TL, de Jong van Berkel YEM (2004) Species-specific variation in the importance of the spectral quality gradient in canopies as a signal for photosynthetic resource partitioning. *Ann Bot* 94: 725–732
- Pospíšil P, Prasad A, Rác M (2019) Mechanism of the formation of electronically excited species by oxidative metabolic processes: Role of reactive oxygen species. *Biomolecules* 9: 258
- Pospíšil P, Šnyrychová I, Nauš J (2007) Dark production of reactive oxygen species in photosystem II membrane particles at elevated temperature: EPR spin-trapping study. *Biochim Biophys Acta Bioenerg* 1767: 854–859
- Poulson ME, Thai T (2015) Effect of high light intensity on photoinhibition, oxyradicals and artemisinin content in *Artemisia annua* L. *Photosynthetica* 53: 403–409
- Pralon T, Shanmugabalaji V, Longoni P, Glauser G, Ksas B, Collombat J, Desmeules S, Havaux M, Finazzi G, Kessler F (2019) Plastoquinone homeostasis by *Arabidopsis* proton gradient regulation 6 is essential for photosynthetic efficiency. *Commun Biol* 2: 220
- Prasad A, Sedlářová M, Kale RS, Pospíšil P (2017) Lipxygenase in singlet oxygen generation as a response to wounding: in vivo imaging in *Arabidopsis thaliana*. *Sci Rep* 7: 9831
- Prasad A, Sedlářová M, Pospíšil P (2018) Singlet oxygen imaging using fluorescent probe Singlet Oxygen Sensor Green in photosynthetic organisms. *Sci Rep* 8 :13685
- Przybyla D, Gobel C, Imboden A, Feussner I, Hamberg M, Apel K (2008) Enzymatic, but not non-enzymatic 1O₂-mediated peroxidation of polyunsaturated fatty acids forms part of the EXECUTER1-dependent stress response program in the flu mutant of *Arabidopsis thaliana*. *Plant J* 54: 236–248
- Pshybytko NL, Kruk J, Kabashnikova LF, Stralka K (2008) Function of plastoquinone in heat stress reactions of plants. *Biochim Biophys Acta* 1777: 1393–1399
- Ptushenko VV, Cherepanov DA, Krishtalik LI, Semenov AY (2008) Semi-continuum electrostatic calculations of redox potentials in photosystem I. *Photosynth Res* 97: 55–74
- Puthiyaveetil S, Kavanagh TA, Cain P, Sullivan JA, Newell CA, Gray JC, Robinson C, van der Giezen M, Rogers MB, Allen JF (2008) The ancestral symbiont sensor kinase CSK links photosynthesis with gene expression in chloroplasts. *Proc Natl Acad Sci USA* 105: 10061–10066
- Ragás X, Jiménez-Banzo A, Sánchez-García D, Batllori X, Nonell S (2009) Singlet oxygen photosensitisation by the fluorescent probe Singlet Oxygen Sensor Green®. *Chem Commun* 20: 2920–2922
- Rajagopal S, Joly D, Gauthier A, Beauregard M, Carpentier R (2005) Protective effect of active oxygen scavengers on protein degradation and photochemical function in photosystem I submembrane fractions during light stress. *FEBS J* 272: 892–902
- Ramel F, Birtic S, Cuié S, Triantaphylidès C, Ravanat JC, Havaux M (2012a) Chemical quenching of singlet oxygen by carotenoids in plants. *Plant Physiol* 158: 1267–1278
- Ramel F, Birtic S, Ginies C, Soubigou-Taconnat L, Triantaphylides C, Havaux M (2012b) Carotenoid oxidation products are stress signals that mediate gene responses to singlet oxygen in plants. *Proc Natl Acad Sci USA* 109: 5535–5540
- Ramel F, Ksas B, Akkari E, Mialoundama AS, Monnet F, Krieger-Liszakay A, Ravanat JC, Mueller MJ, Bouvier F, Havaux M (2013) Light-induced acclimation of the *Arabidopsis thaliana* mutant to singlet oxygen. *Plant Cell* 25: 1445–1462
- Ranjbar Choubeh R, Wientjes E, Struik PC, Kirilovsky D, van Amerongen H (2018) State transitions in the cyanobacterium *Synechococcus elongatus* 7942 involve reversible quenching of the photosystem II core. *Biochim Biophys Acta Bioenerg* 1859: 1059–1066

- Rantala M, Rantala S, Aro EM (2020a) Composition, phosphorylation and dynamic organization of photosynthetic protein complexes in plant thylakoid membrane. *Photochem Photobiol Sci* DOI: 10.1039/d0pp00025f
- Rantala S, Lempiäinen T, Gerotto C, Tiwari A, Aro EM, Tikkanen M (2020b) PGR5 and NDH-1 systems do not function as protective electron acceptors but mitigate the consequences of PSI inhibition. *Biochim Biophys Acta Bioenerg* 1861: 148154
- Rappaport F, Lavergne J (2009) Thermoluminescence: theory. *Photosynth Res* 101: 205–216
- Rastogi A, Yadav DK, Szymańska R, Kruk J, Sedlářová M, Pospíšil P (2014) Singlet oxygen scavenging activity of tocopherol and plastoquinone in *Arabidopsis thaliana*: relevance to photooxidative stress. *Plant Cell Environ* 37: 392–401
- Rastogi VK, Girvin ME (1999) Structural changes linked to proton translocation by subunit c of the ATP synthase. *Nature* 402: 263–268
- Raven JA (2009) Contributions of anoxygenic and oxygenic phototrophy and chemolithotrophy to carbon and oxygen fluxes in aquatic environments. *Aquat Microb Ecol* 56: 177–192
- Redmond RW, Gamlin JN (1999) A compilation of singlet oxygen yields from biologically relevant molecules. *Photochem Photobiol* 70: 391–475
- Rehman AU, Cser K, Sass L, Vass I (2013) Characterization of singlet oxygen production and its involvement in photodamage of Photosystem II in the cyanobacterium *Synechocystis* PCC 6803 by histidine-mediated chemical trapping. *Biochim Biophys Acta* 1827: 689–698
- Rehman AU, Kodru S, Vass I (2016b) Chloramphenicol mediates superoxide production in Photosystem II and enhances its photodamage in isolated membrane particles. *Front Plant Sci* 7: 479
- Rehman AU, Szabó M, Deák Z, Sass L, Larkum A, Ralph P, Vass I (2016a) *Symbiodinium* sp. cells produce light-induced intra- and extracellular singlet oxygen, which mediates photodamage of the photosynthetic apparatus and has the potential to interact with the animal host in coral symbiosis. *New Phytol* 212: 472–484
- Reinman S, Mathis P (1981) Influence of temperature on photosystem II electron transfer reactions. *Biochim Biophys Acta* 635: 249–258
- Renger G, Völker M, Eckert HJ, Fromme R, Hohm-Weit S, Graber P (1989) On the mechanism of Photosystem II deterioration by UV-B irradiation. *Photochem Photobiol* 49: 97–105
- Renger T, Schlodder E (2011) Optical properties, excitation energy and primary charge transfer in photosystem II: Theory meets experiment. *J Photochem Photobiol B* 104: 126–141
- Rintamäki E, Martinsuo P, Pursiheimo S, Aro EM (2000) Cooperative regulation of light-harvesting complex II phosphorylation via the plastoquinol and ferredoxin-thioredoxin system in chloroplasts. *Proc Natl Acad Sci USA* 97: 11644–11649
- Rippka R, Deruelles J, Waterbury JB, Herdman M, Stanier RY (1979) Generic assignments, strain histories and properties of pure cultures of cyanobacteria. *J Gen Microbiol* 111: 1–61
- Roach T, Krieger-Liszkay A (2012) The role of the PsbS protein in the protection of photosystems I and II against high light in *Arabidopsis thaliana*. *Biochim Biophys Acta* 1817: 2158–2165
- Roach T, Sedoud A, Krieger-Liszkay A (2013) Acetate in mixotrophic growth medium affects photosystem II in *Chlamydomonas reinhardtii* and protects against photoinhibition. *Biochim Biophys Acta* 1827: 1183–1190
- Roach T, Sun Na C, Stögg W, Krieger-Liszkay A (2020) The non-photochemical quenching protein LHCSR3 prevents oxygen-dependent photoinhibition in *Chlamydomonas reinhardtii*. *J Exp Bot* 71: 2650–2660
- Rokka A, Suorsa M, Saleem A, Battchikova N, Aro EM (2005) Synthesis and assembly of thylakoid protein complexes: multiple assembly steps of photosystem II. *Biochem J* 388: 159–168
- Romero E, Novoderezhkin VI, van Grondelle R (2017) Quantum design of photosynthesis for bio-inspired solar-energy conversion. *Nature* 543: 355–365
- Romero E, van Stokkum IHM, Novoderezhkin VI, Dekker JP, van Grondelle R (2010) Two different charge separation pathways in photosystem II. *Biochemistry* 49: 4300–4307

- Rova M, Mamedov F, Magnuson A, Fredriksson PO, Styring S (1998) Coupled activation of the donor and the acceptor side of Photosystem II during photoactivation of the oxygen evolving cluster. *Biochemistry* 37: 11039–11045
- Rozema J, Bjorn LO, Bornman JF, Gaberšček A, Hader DP, Trost T, Germ M, Klisch M, Gröniger A, Sinha RP, Lebert M, He YY, Buffoni-Hall R, de Bakker NVJ, van de Staaij J, Meijkamp BB (2002) The role of UV-B radiation in aquatic and terrestrial ecosystems—an experimental and functional analysis of the evolution of UV-absorbing compounds. *J Photochem Photobiol B* 66: 2–12
- Ruettinger WF, Yagi M, Wolf K, Bernasek S, Dismukes GC (2000) O₂ evolution from the manganese-oxo cubane core Mn₄O₄6⁺: a molecular mimic of the photosynthetic water oxidation enzyme? *J Am Chem Soc* 122: 10353–10357
- Rutherford AW, Mullet JE (1981) Reaction center triplet states in Photosystem I and Photosystem II. *Biochim Biophys Acta Bioenerg* 635: 225–235
- Rutherford AW, Osyczka A, Rappaport F (2012) Back-reactions, short-circuits, leaks and other energy wasteful reactions in biological electron transfer: Redox tuning to survive life in O₂. *FEBS Lett* 586: 603–616
- Sacharz J, Bryan SJ, Yu J, Burroughs NJ, Spence EM, Nixon PJ, Mullineaux CW (2015) Sub-cellular location of FtsH proteases in the cyanobacterium *Synechocystis* sp. PCC 6803 suggests localised PSII repair zones in the thylakoid membranes. *Mol Microbiol* 96: 448–462
- Saito K, Rutherford AW, Ishikita H (2013) Mechanism of proton-coupled quinone reduction in Photosystem II. *Proc Natl Acad Sci USA* 110: 954–959
- Sang M, Ma F, Xie J, Chen XB, Wang KB, Qin XB, Wang WD, Zhao JQ, Li LB, Zhang JP, Kuang TY (2010) High-light induced singlet oxygen formation in cytochrome b6f complex from *Bryopsis corticulans* detected by EPR spectroscopy. *Biophys Chem* 146: 7–12
- Santabarbara S, Agostini G, Casazza AP, Syme CD, Heathcote P, Böhles F, Evans MCW, Jennings RC, Carbonera D (2007) Chlorophyll triplet states associated with Photosystem I and Photosystem II in thylakoids of the green alga *Chlamydomonas reinhardtii*. *Biochim Biophys Acta* 1767: 88–105
- Santabarbara S, Bullock B, Rappaport F, Redding KE (2015) Controlling electron transfer between the two cofactor chains of photosystem I by the redox state of one of their components. *Biophys J* 108: 1537–1547
- Santabarbara S, Cazzalini I, Rivadossi A, Garlaschi FM, Zucchelli G, Jennings RC (2002) Photoinhibition in vivo and in vitro involves weakly coupled chlorophyll-protein complexes. *Photochem Photobiol* 75: 613–618
- Santabarbara S, Galuppi L, Casazza AP (2010) Bidirectional electron transfer in the reaction centre of photosystem I. *J Integ Plant Biol* 52: 735–749
- Sarvikas P, Hakala M, Pätsikkä E, Tyystjärvi T, Tyystjärvi E (2006) Action spectrum of photoinhibition in leaves of wild type and npq1-2 and npq4-1 mutants of *Arabidopsis thaliana*. *Plant Cell Physiol* 47: 391–400
- Sarvikas P, Tyystjärvi T, Tyystjärvi E (2010) Kinetics of prolonged photoinhibition revisited. Photoinhibited Photosystem II centres do not protect the active ones against loss of oxygen evolution. *Photosyn Res* 103: 7–17
- Satoh K (1970) Mechanism of photoinactivation in photosynthetic systems II. The occurrence and properties of two different types of photoinactivation. *Plant Cell Physiol* 11: 29–38
- Schansker G, Tóth SZ, Holzwarth AR, Garab G (2014) Chlorophyll a fluorescence: beyond the limits of the QA model. *Photosynth Res* 120: 43–58
- Schatz GH, Brock H, Holzwarth AR (1987) Picosecond kinetics of fluorescence and absorbance changes in photosystem II particles excited at low photon density. *Proc Natl Acad Sci USA* 84: 8414–8418
- Schlodder E, Brettel K (1988) Primary charge separation in closed Photosystem II with a lifetime of 11 ns: Flash-absorption spectroscopy with O₂-evolving Photosystem II complexes from *Synechococcus*. *Biochim Biophys Acta* 933: 22–34

- Schreiber U, Klughammer C (2016) Analysis of Photosystem I donor and acceptor sides with a new type of online-deconvoluting kinetic LED-array spectrophotometer. *Plant Cell Physiol* 57: 1454–1467
- Schuermans RM, Schuurmans JM, Bekker M, Kromkamp JC, Matthijs HCP, Hellingwerf KJ (2014) The redox potential of the plastoquinone pool of the cyanobacterium *Synechocystis* species strain PCC6803 is under strict homeostatic control. *Plant Physiol* 165: 463–475
- Schwarz EM, Tietz S, Froehlich JE (2018) Photosystem I-LHCII megacomplexes respond to high light and aging in plants. *Photosynth Res* 136: 107–124
- Seemann JR, Sharkey TD, Wang J, Osmond CB (1987) Environmental effects on photosynthesis, nitrogen-use efficiency, and metabolite pools in leaves of sun and shade plants. *Plant Physiol* 84: 796–802
- Seelert H, Poetsch A, Dencher NA, Engel A, Stahlberg H, Müller DJ (2000) Proton-powered turbine of a plant motor. *Nature* 405: 418–419
- Serôdio J, Schmidt W, Frankenbach S (2017) A chlorophyll fluorescence-based method for the integrated characterization of the photophysiological response to light stress. *J Exp Bot* 68: 1123–1135
- Sétif P, Hervo G, Mathis P (1981) Flash-induced absorption changes in Photosystem-I - radical pair or triplet-state formation. *Biochim Biophys Acta* 638: 257–267
- Shikanai T, Endo T, Hashimoto T, Yamada Y, Asada K, Yokota A (1998) Directed disruption of the tobacco *ndhB* gene impairs cyclic electron flow around photosystem I. *Proc Natl Acad Sci USA* 95: 9705–9709
- Shinopoulos KE, Brudvig GW (2012) Cytochrome b559 and cyclic electron transfer within photosystem II. *Biochim Biophys Acta Bioenerg* 1817: 66–75
- Shinopoulos KE, Yu J, Nixon PJ, Brudvig GW (2014) Using site-directed mutagenesis to probe the role of the D2 carotenoid in the secondary electron-transfer pathway of photosystem II. *Photosynth Res* 120: 141–152
- Shipman LL (1979) A theoretical study of excitons in chlorophyll a photosystems on a picosecond timescale. *Photochem Photobiol* 31: 157–167
- Siegbahn PEM (2013) Water oxidation mechanism in photosystem II, including oxidations, proton release pathways, O–O bond formation and O₂ release. *Biochim Biophys Acta* 1827: 1003–1019
- Silva P, Thompson E, Bailey S, Kruse O, Mullineaux CW, Robinson C, Mann NH, Nixon PJ (2003) FtsH is involved in the early stages of repair of photosystem II in *Synechocystis* sp PCC 6803. *Plant Cell* 15: 2152–2164
- Sinvany-Villalobo G, Davydov O, Ben-Ari G, Zaltsman A, Raskind A, Adam Z (2004) Expression in multigene families. Analysis of chloroplast and mitochondrial proteases. *Plant Physiol* 135: 1336–1345
- Smith HL, McAusland L, Murchie EH (2017) Don't ignore the green light: exploring diverse roles in plant processes. *J Exp Bot* 68: 2099–2110
- Song YG, Liu B, Wang LF, Li MH, Liu Y (2006) Damage to the oxygen-evolving complex by superoxide anion, hydrogen peroxide, and hydroxyl radical in photoinhibition of photosystem II. *Photosynth Res* 90: 67–78
- Sonoike K (1996) Degradation of *psaB* gene product, the reaction center subunit of photosystem I, is caused during photoinhibition of photosystem I: Possible involvement of active oxygen species. *Plant Sci* 115: 157–164
- Sonoike K (2011) Photoinhibition of photosystem I. *Physiol Plant* 142: 56–64
- Spetea C, Herdean A, Alloreant G, Carraretto L, Finazzi G, Szabo I (2017) An update on the regulation of photosynthesis by thylakoid ion channels and transporters in *Arabidopsis*. *Physiol Plant* 161: 16–27
- Spetea C, Hideg É, Vass I (1997) Low pH accelerates light-induced damage of photosystem II by enhancing the probability of the donor-side mechanism of photoinhibition. *Biochim Biophys Acta Bioenerg* 1318: 275–283

- Stamatakis K, Papageorgiou GC, Govindjee (2016) Effects of exogenous beta-carotene, a chemical scavenger of singlet oxygen, on the millisecond rise of chlorophyll a fluorescence of cyanobacterium *Synechococcus* sp PCC 7942. *Photosynth Res* 130: 317–324
- Stemler A, Jursinic P (1983) The effects of carbonic anhydrase inhibitors formate, bicarbonate, acetazolamide, and imidazole on Photosystem II in maize chloroplasts. *Arch Biochem Biophys* 221: 227–237
- Stiehl HH, Witt HT (1969) Quantitative treatment of the function of plastoquinone in photosynthesis. *Z Naturforsch B* 24: 1588–1598
- Suga M et al. (2019) An oxyl/oxo mechanism for oxygen-oxygen coupling in PSII revealed by an x-ray free-electron laser. *Science* 366: 334–338
- Sumi H (1999) Theory on rates of excitation-energy transfer between molecular aggregates through distributed transition dipoles with application to the antenna system in bacterial photosynthesis. *J Phys Chem B* 103: 252–260
- Sun H, Zhang SB, Liu T, Huan W (2020) Decreased photosystem II activity facilitates acclimation to fluctuating light in the understory plant *Paris polyphylla*. *Biochim Biophys Acta Bioenerg* 1861: 148135
- Suorsa M, Järvi S, Grieco M, Nurmi M, Pietrzykowska M, Rantala M, Kangasjärvi S, Paakkanen V, Tikkanen M, Jansson S, Aro EM (2012) PROTON GRADIENT REGULATION5 is essential for proper acclimation of *Arabidopsis* photosystem I to naturally and artificially fluctuating light conditions. *Plant Cell* 24: 2934–2948
- Suter GW, Mazzola P, Wendler J, Holzwarth AR (1984) Fluorescence decay kinetics in phycobilisomes isolated from the bluegreen alga *Synechococcus* 6301. *Biochim Biophys Acta Bioenerg* 766: 269–276
- Suzuki H, Sugiura M, Noguchi T (2012) Determination of the miss probabilities of individual S state transitions during photosynthetic water oxidation by monitoring electron flow in Photosystem II using FTIR spectroscopy. *Biochemistry* 51: 6776–6785
- Szechyńska-Hebda M, Kruk J, Górecka M, Karpińska B and Karpiński S (2010) Evidence for light wavelength-specific photoelectrophysiological signaling and memory of excess light episodes in *Arabidopsis*. *Plant Cell* 22: 2201–2218
- Szilárd A, Sass L, Deák Z, Vass I (2007) The sensitivity of Photosystem II to damage by UV-B radiation depends on the oxidation state of the water-splitting complex. *Biochim Biophys Acta* 1767: 876–882
- Takagi D, Takumi S, Hashiguchi M, Sejima T, Miyake C (2016) Superoxide and singlet oxygen produced within the thylakoid membranes both cause Photosystem I photoinhibition. *Plant Physiol* 171: 1626–1634
- Takahashi M, Asada K (1988) Superoxide production in aprotic interior of chloroplast thylakoids. *Arch Biochem Biophys* 267: 714–722
- Takahashi Y, Hansson Ö, Mathis P, Satoh K (1987) Primary radical pair in the Photosystem II reaction centre. *Biochim Biophys Acta Bioenerg* 893: 49–59
- Takegawa Y, Nakamura M, Nakamura S, Noguchi T, Sellés J, Rutherford AW, Boussac A, Sugiura M (2019) New insights on ChlD1 function in Photosystem II from site-directed mutants of D1/T179 in *Thermosynechococcus elongatus*. *Biochim Biophys Acta Bioenerg* 1860: 297–309
- Takizawa K, Cruz JA, Kanazawa A, Kramer DM (2007) The thylakoid proton motive force in vivo. Quantitative, non-invasive probes, energetics, and regulatory consequences of light-induced pmf. *Biochim Biophys Acta* 1767: 1233–1244
- Tarahi Tabrizi S, Sawicki A, Zhou S, Luo M, Willows RD (2016) GUN4-Protochlorophyllin IX is a singlet oxygen generator with consequences for plastid retrograde signaling. *J Biol Chem* 291: 8978–8984
- Taylor RM, Sallans L, Frankel LK, Bricker TM (2018) Natively oxidized amino acid residues in the spinach cytochrome b6/f complex. *Photosynth Res* 137: 141–151
- Telfer A, Barber J (1995) Role of carotenoid bound to the photosystem II reaction centre. In: Mathis P (ed) *Photosynthesis: from Light to Biosphere*, Kluwer, Dordrecht, pp 15–20

- Telfer A, Bishop SM, Phillips D, Barber J (1994) Isolated photosynthetic reaction center of Photosystem II as a sensitizer for the formation of singlet oxygen. Detection and quantum yield determination using a chemical trapping technique. *J Biol Chem* 269: 13244–13253
- Telfer A, De Las Rivas J, Barber J (1991) β -Carotene within the isolated Photosystem II reaction centre: photooxidation and irreversible bleaching of this chromophore by oxidised P680. *Biochim Biophys Acta* 1060: 106–114
- Terashima I, Fujita T, Inoue T, Chow WS, Oguchi R (2009) Green light drives leaf photosynthesis more efficiently than red light in strong white light: revisiting the enigmatic question of why leaves are green. *Plant Cell Physiol* 50: 684–697
- Terashima I, Funayama S, Sonoike K (1994) The site of photoinhibition in leaves of *Cucumis sativus* L. At low temperatures is photosystem I, not photosystem II. *Planta* 193: 300–306
- Theis J, Schroda M (2016) Revisiting the photosystem II repair cycle. *Plant Signal Behav* 11: e1218587
- Thomas JB, Van Der Wal UP (1963) Absorption responses of chlorophyll in vivo to treatment with acetone. *Biochim Biophys Acta* 79: 490–499
- Thompson LK, Brudvig GW (1988) Cytochrome b-559 may function to protect Photosystem II from photoinhibition. *Biochemistry* 27: 6653–6658
- Tian L, Xu P, Chukhutsina VU, Holzwarth AR, Croce R (2017) Zeaxanthin-dependent nonphotochemical quenching does not occur in photosystem I in the higher plant *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 114: 4828–4832
- Tichý M, Lupínková L, Sicora C, Vass I, Kuvíková S, Prásl O, Komenda J (2003) Synechocystis 6803 mutants expressing distinct forms of the Photosystem II D1 protein from *Synechococcus* 7942: relationship between the psbA coding region and sensitivity to visible and UV-B radiation. *Biochim Biophys Acta* 1605: 55–66
- Tikhonov AN (2018) The cytochrome b6f complex: Biophysical aspects of its functioning in chloroplasts. In: Harris JR, Boekema EJ (Eds) *Membrane Protein Complexes: Structure and Function*, Springer Nature, Singapore, pp. 287–328
- Tikhonov AN, Khomutov GB, Ruuge EK (1984) Electron transport control in chloroplasts. Effects of magnesium ions on the electron flow between two photosystems. *Photobiochem Photobiophys* 8: 261–269
- Tikkanen M, Grieco M, Kangasjärvi S, Aro EM (2010) Thylakoid protein phosphorylation in higher plant chloroplasts optimizes electron transfer under fluctuating light. *Plant Physiol* 152: 723–735
- Tikkanen M, Mekala NR, Aro EM (2014) Photosystem II photoinhibition-repair cycle protects Photosystem I from irreversible damage. *Biochim Biophys Acta* 1837: 210–215
- Tilbrook K, Dubois M, Crocco CD, Yin R, Chappuis R, Allore G, Schmid-Siegert E, Goldschmidt-Clermont M, Ulm R (2016) UV-B perception and acclimation in *Chlamydomonas reinhardtii*. *Plant Cell* 28: 966–983
- Tiwari A, Mamedov F, Grieco M, Suorsa M, Jajoo A, Styring S, Tikkanen M, Aro EM (2016) Photodamage of iron-sulphur clusters in photosystem I induces non-photochemical energy dissipation. *Nat Plants* 2: 16035
- Tjus SE, Lindberg Møller B, Scheller HV (1998) Photosystem I is an early target of photoinhibition in barley illuminated at chilling temperatures. *Plant Physiol* 116: 755–764
- Tomo T, Kusakabe H, Nagao R, Ito H, Tanaka A, Akimoto S, Mimuro M, Okazaki S (2012) Luminescence of singlet oxygen in photosystem II complexes isolated from cyanobacterium *Synechocystis* sp. PCC6803 containing monovinyl or divinyl chlorophyll a. *Biochim Biophys Acta Bioenerg* 1817: 1299–1305
- Tóth SZ, Shansker G, Strasser RJ (2007) A non-invasive assay of the plastoquinone pool redox state based on the OJIP-transient. *Photosynth Res* 93: 193–203
- Townsend AJ, Retkute R, Chinnathambi K, Randall JWP, Foulkes J, Carmo-Silva E, Murchie EH (2018) Suboptimal acclimation of photosynthesis to light in wheat canopies. *Plant Physiol* 176: 1233–1246

- Treves H, Raanan H, Kedem I, Murik O, Keren N, Zer H, Berkowicz SM, Giordano M, Norici A, Shotland Y, Ohad I, Kaplan A (2016) The mechanisms whereby the green alga *Chlorella ohadii*, isolated from desert soil crust, exhibits unparalleled photodamage resistance. *New Phytol* 210: 1229–1243
- Triantaphylidès C, Kruschke M, Hoerberichts FA, Ksas B, Gresser G, Havaux M, Van Breusegem F, Mueller MJ (2008) Singlet oxygen is the major reactive oxygen species involved in photooxidative damage to plants. *Plant Physiol* 148: 960–968
- Trotta A, Suorsa M, Rantala M, Lundin B, Aro EM (2016) Serine and threonine residues of plant STN7 kinase are differentially phosphorylated upon changing light conditions and specifically influence the activity and stability of the kinase. *Plant J* 87: 484–494
- Tschiersch H, Ohmann E (1993) Photoinhibition in *Euglena gracilis*: involvement of reactive oxygen species. *Planta* 191: 316–323
- Tsonev TD, Hikosaka K (2003) Contribution of photosynthetic electron transport, heat dissipation, and recovery of photoinactivated Photosystem II to photoprotection at different temperatures in *Chenopodium album* leaves. *Plant Cell Physiol* 44: 828–835
- Tsuyama M, Kobayashi Y (2008) Reduction of the primary donor P700 of photosystem I during steady-state photosynthesis under low light in *Arabidopsis*. *Photosynth Res* 99: 37
- Tullberg A, Alexciev K, Pfannschmidt T, Allen JF (2000) Photosynthetic electron flow regulates transcription of the *psaB* gene in pea (*Pisum sativum* L.) chloroplasts through the redox state of the plastoquinone pool. *Plant Cell Physiol* 41: 1045–1054
- Tyystjärvi E (2004) Phototoxicity. In: Noodén LD (ed) *Plant Cell Death Processes*, Elsevier Academic Press, Amsterdam, pp 271–283
- Tyystjärvi E (2013) Photoinhibition of Photosystem II. *Int Rev Cell Mol Biol* 300: 343–303
- Tyystjärvi E, Ali-Yrkkö K, Kettunen R, Aro EM (1992) Slow degradation of the D1 protein is related to the susceptibility of low-light-grown pumpkin plants to photoinhibition. *Plant Physiol* 100: 1310–1317
- Tyystjärvi E, Aro EM (1996) The rate constant of photoinhibition, measured in lincomycin-treated leaves, is directly proportional to light intensity. *Proc Natl Acad Sci USA* 93: 2213–2218
- Tyystjärvi E, Hakala M, Sarvikas P (2005) Mathematical modelling of the light response curve of photoinhibition of Photosystem II. *Photosyn Res* 84: 21–27
- Tyystjärvi E, Kettunen R, Aro EM (1994) The rate constant of photoinhibition in vitro is independent of the antenna size of Photosystem II but depends on temperature. *Biochim Biophys Acta* 1186: 177–185
- Tyystjärvi T, Tuominen I, Herranen M, Aro EM, Tyystjärvi E (2002) Action spectrum of *psbA* gene transcription is similar to that of photoinhibition in *Synechocystis* sp. PCC 6803. *FEBS Lett* 516: 167–171
- Ueno M, Sae-Tang P, Kusama Y, Hihara Y, Matsuda M, Hasunuma T, Nishiyama Y (2016) Moderate heat stress stimulates repair of photosystem ii during photoinhibition in *Synechocystis* sp. PCC 6803. *Plant Cell Physiol* 57: 2417–2426
- Umena Y, Kawakami K, Shen JR, Kamiya N (2011) Crystal structure of oxygen-evolving photosystem II at a resolution of 1.9 Å. *Nature* 473: 55–60
- Ünlü C, Drop B, Croce R, van Amerongen H (2014) State transitions in *Chlamydomonas reinhardtii* strongly modulate the functional size of photosystem II but not of photosystem I. *Proc Natl Acad Sci USA* 111: 3460–3465
- van Mieghem F, Brettel K, Hillmann B, Kamlowski A, Rutherford AW, Schlodder E (1995) Charge recombination reactions in Photosystem II. 1. Yields, recombination pathways, and kinetics of the primary pair. *Biochemistry* 34: 4798–4813
- Van Walraven HS, Strotmann H, Schwarz O, Rumberg B (1996) The H⁺/ATP coupling ratio of the ATP synthase from thiol-modulated chloroplasts and two cyanobacterial strains is four. *FEBS Lett* 379: 309–313

- Vasil'ev S, Bergmann A, Redlin H, Eichler HJ, Renger G (1996) On the role of exchangeable hydrogen bonds for the kinetics of P680+QA⁻ formation and P680+Pheo⁻ recombination in photosystem II. *Biochim Biophys Acta* 1276: 35–44
- Vass I (2012) Molecular mechanisms of photodamage in the Photosystem II complex. *Biochim Biophys Acta Bioenerg* 1817: 209–217
- Vass I, Cser K (2009) Janus-faced charge recombinations in photosystem II photoinhibition. *Trends Plant Sci* 14: 4
- Vass I, Defik Z, Hideg É (1990) Charge equilibrium between the water-oxidizing complex and the electron donor tyrosine-D in Photosystem II. *Biochim Biophys Acta* 1017: 63–69
- Vass I, Sass L, Spetea C, Bakou A, Ghanotakis D, Petrouleas V (1996) UV-B induced inhibition of Photosystem II electron transport studied by EPR and chlorophyll fluorescence. Impairment of donor and acceptor side components. *Biochemistry* 35: 8964–8973
- Vassiliev S, Zارايسкая T, Bruce D (2013) Molecular dynamics simulations reveal highly permeable oxygen exit channels shared with water uptake channels in photosystem II. *Biochim Biophys Acta* 1827: 1148–1155
- Vener AV, Van Kan PJM, Gal A, Andersson B, Ohad I (1995) Activation/deactivation cycle of redox-controlled thylakoid protein phosphorylation. Role of plastoquinol bound to the reduced cytochrome b_f complex. *J Biol Chem* 270: 25225–25232
- Verhoeven A (2014) Sustained energy dissipation in winter evergreens. *New Phytol* 201: 57–65
- Vermaas WFJ, Renger G, Dohnt G (1985) The reduction of the oxygen evolving system in chloroplasts by thylakoid components. *Biochim Biophys Acta* 764: 194–202
- Vinyard DJ, Gimpel J, Ananyev GM, Cornejo MA, Golden SS, Mayfield SP, Dismukes GC (2013) Natural variants of Photosystem II subunit D1 tune photochemical fitness to solar intensity. *J Biol Chem* 288: 5451–5462
- Vladkova R (2016) Chlorophyll a is the crucial redox sensor and transmembrane signal transmitter in the cytochrome b₆f complex. Components and mechanisms of state transitions from the hydrophobic mismatch viewpoint. *J Biomol Struct Dyn* 34: 824–854
- Vonshak A, Novoplansky N (2008) Acclimation to low temperature of two *Arthrospira platensis* (cyanobacteria) strains involves down-regulation of PSII and improved resistance to photoinhibition. *J Phycol* 44: 1071–1079
- von Sydow L, Schwenkert S, Meurer J, Funk C, Mamedov F, Schröder WP (2016) The PsbY protein of *Arabidopsis* Photosystem II is important for the redox control of cytochrome b₅₅₉. *Biochim Biophys Acta Bioenerg* 1857: 1524–1533
- Wagner R, Dietzel L, Bräutigam K, Fischer W, Pfannschmidt T (2008) The long-term response to fluctuating light quality is an important and distinct light acclimation mechanism that supports survival of *Arabidopsis thaliana* under low light conditions. *Planta* 228: 573–587
- Wang F, Qi Y, Malnoë A, Choquet Y, Wollman FA, de Vitry C (2017) The high light response and redox control of thylakoid FtsH protease in *Chlamydomonas reinhardtii*. *Mol Plant* 10: 99–114
- Wang L, Kim C, Xu X, Piskurewicz U, Dogra V, Singh S, Mahler H, Apel K (2016a) Singlet oxygen- and EXECUTER1-mediated signaling is initiated in grana margins and depends on the protease FtsH2. *Proc Natl Acad Sci USA* 113: E3792–E3800
- Wang WH, He EM, Chen J, Guo Y, Chen J, Liu X, Zheng HL (2016b) The reduced state of the plastoquinone pool is required for chloroplast-mediated stomatal closure in response to calcium stimulation. *Plant J* 86: 132–144
- Watanabe M, Ikeuchi M (2013) Phycobilisome: architecture of a light-harvesting supercomplex. *Photosynth Res* 116: 265–276
- Weisz DA, Johnson VM, Niedzwiedzki DM, Shinn MK, Liu H, Klitzke CF, Gross ML, Blankenship RE, Lohman TM, Pakrasi HB (2019) A novel chlorophyll protein complex in the repair cycle of photosystem II. *Proc Natl Acad Sci USA* 116: 21907–21913

- Wiciarz M, Niewiadomska E, Kruk J (2018) Effects of salt stress on low molecular antioxidants and redox state of plastoquinone and P700 in *Arabidopsis thaliana* (glycophyte) and *Eutrema salsgineum* (halophyte). *Photosynthetica* 56: 811–819
- Wientjes E, Drop B, Kouril R, Boekema EJ, Croce R (2013a) During state 1 to state 2 transition in *Arabidopsis thaliana*, the photosystem II supercomplex gets phosphorylated but does not disassemble. *J Biol Chem* 288: 32821–32826
- Wientjes E, van Amerongen H, Croce R (2013b) LHCII is an antenna of both photosystems after long-term acclimation. *Biochim Biophys Acta* 1827: 420–426
- Wientjes E, van Stokkum IH, van Amerongen H, Croce R (2011) The role of the individual Lhcas in photosystem I excitation energy trapping. *Biophys J* 101: 745–754
- Willig A, Shapiguzov A, Goldschmidt-Clermont M, Rochaix JD (2011) The phosphorylation status of the chloroplast protein kinase STN7 of *Arabidopsis* affects its turnover. *Plant Physiol* 157: 2102–2107
- Wilson A, Ajlani G, Verbavatz JM, Vass I, Kerfeld CA, Kirilovsky D (2006) A soluble carotenoid protein involved in phycobilisome-related energy dissipation in cyanobacteria. *Plant Cell* 18: 992–1007
- Wilson S, Ruban AV (2020) Rethinking the influence of chloroplast movements on non-photochemical quenching and photoprotection. *Plant Physiol* 183: 1213–1223
- Wood WHJ, MacGregor-Chatwin C, Barnett SFH, Mayneord GE, Huang X, Hobbs JK, Hunter CN, Johnson MP (2018) Dynamic thylakoid stacking regulates the balance between linear and cyclic photosynthetic electron transfer. *Nat Plants* 4: 116–127
- Yadav DK, Pospíšil P (2012) Evidence on the formation of singlet oxygen in the donor side photoinhibition of Photosystem II: EPR spin-trapping study. *PLoS One* 7: e45883
- Yamamoto T, Burke J, Autz G, Jagendorf AT (1981) Bound ribosomes of pea chloroplast thylakoid membranes: location and release in vitro by high salt, puromycin, and RNase. *Plant Physiol* 67: 940–949
- Yamashita A, Nijo N, Pospíšil P, Morita N, Takenaka D, Aminaka R, Yamamoto Y, Yamamoto Y (2008) Quality control of Photosystem II. Reactive oxygen species are responsible for the damage to Photosystem II under moderate heat stress. *J Biol Chem* 283: 28380–28391
- Yamori W, Shikanai T (2016) Physiological functions of cyclic electron transport around Photosystem I in sustaining photosynthesis and plant growth. *Annu Rev Plant Biol* 67: 81–106
- Yang DH, Andersson B, Aro EM, Ohad I (2001) The redox state of the plastoquinone pool controls the level of the light-harvesting chlorophyll a/b binding protein complex II (LHC II) during photoacclimation. *Photosynth Res* 68: 163–174
- Yao M, Liu Y, Fei L, Zhou Y, Wang F, Chen J (2018) Self-adaptable quinone–quinol exchange mechanism of Photosystem II. *J Phys Chem B* 122: 10478–10489
- Yoshida K, Shibata M, Terashima I, Noguchi K (2010) Simultaneous determination of in vivo plastoquinone and ubiquinone redox states by HPLC-based analysis. *Plant Cell Physiol* 51: 836–841
- Yutthanasirikul R, Nagano T, Jimbo H, Hihara Y, Kanamori T, Ueda T, Haruyama T, Konno H, Yoshida K, Hisabori T, Nishiyama Y (2016) Oxidation of a cysteine residue in elongation factor EF-Tu reversibly inhibits translation in the cyanobacterium *Synechocystis* sp. PCC 6803. *J Biol Chem* 291: 5860–5870
- Zaltsman A, Feder A, Adam Z (2005) Developmental and light effects on the accumulation of FtsH protease in *Arabidopsis* chloroplasts: Implications for thylakoid formation and photosystem II maintenance. *Plant J* 42: 609–617
- Zang LY, van Kуйk FJGM, Misra BR, Misra HP (1995) The specificity and product of quenching singlet oxygen by 2, 2, 6, 6-tetramethylpiperidine. *Biochem Mol Biol Int* 37: 283–293
- Zavafer A, Cheah MH, Hillier W, Chow WS, Takahashi S (2015b) Photodamage to the oxygen evolving complex of photosystem II by visible light. *Sci Rep* 5: 16363

- Zavafer A, Chow WS, Cheah MH (2015a) The action spectrum of Photosystem II photoinactivation in visible light. *J Photochem Photobiol B* 152: 247–260
- Zavafer A, Koinuma W, Chow WS, Cheah MH, Mino H (2017) Mechanism of photodamage of the oxygen evolving mn cluster of Photosystem II by excessive light energy. *Sci Rep* 7: 7604
- Zhang C, Chen C, Dong H, Shen JR, Dau H, Zhao J (2015) A synthetic Mn4Ca-cluster mimicking the oxygen-evolving center of photosynthesis. *Science* 348: 690–693
- Zhang T, Maruhnich SA, Folta KM (2011) Green light induces shade avoidance symptoms. *Plant Physiol* 157: 1528–1536
- Zhang L, Paakkarinen V, Suorsa M, Aro EM (2001) A SecY homologue is involved in chloroplast-encoded D1 protein biogenesis. *J Biol Chem* 276: 37809–37814
- Zhang L, Paakkarinen V, van Wijk KJ, Aro EM (1999) Co-translational assembly of the D1 protein into photosystem II. *J Biol Chem* 274: 16062–16067
- Zhang S, Scheller HV (2004) Photoinhibition of photosystem I at chilling temperature and subsequent recovery in *Arabidopsis thaliana*. *Plant Cell Physiol* 45: 1595–1602
- Zhao LS, Huokko T, Wilson S, Simpson S, Wang Q, Ruban AV, Mullineaux CW, Zhang YZ, Liu LN (2020) Structural variability, coordination and adaptation of a native photosynthetic machinery. *Nat Plants* 6: 869, DOI: 10.1038/s41477-020-0694-3
- Zito F, Finazzi G, Delosme R, Nitschke W, Picot D, Wollman FA (1999) The Qo site of cytochrome b6f complexes controls the activation of the LHCII kinase. *EMBO J* 18: 2961–2969
- Zobnina V, Lambrevia MD, Rea G, Campi G, Antonacci A, Scognamiglio V, Giardi MT, Polticelli F (2017) The plastoquinol-plastoquinone exchange mechanism in photosystem II: insight from molecular dynamics simulations. *Photosynth Res* 131: 15–30
- Zoeller M, Stingl N, Krischke M, Fekete A, Waller F, Berger S, Mueller MJ (2012) Lipid profiling of the *Arabidopsis* hypersensitive response reveals specific lipid peroxidation and fragmentation processes: Biogenesis of pimelic and azelaic acid. *Plant Physiol* 160: 365–378



**UNIVERSITY
OF TURKU**

ISBN 978-951-29-8269-1 (PRINT)
ISBN 978-951-29-8270-7 (PDF)
ISSN 0082-7002 (Print)
ISSN 2343-3175 (Online)