

Potential Biomarkers, Risk Factors, and Their Associations with IgE-Mediated Food Allergy in Early Life: A Narrative Review

Caroline E Childs,^{1,2} Daniel Munblit,^{3,4,5} Laurien Ulfman,⁶ Carlos Gómez-Gallego,⁷ Liisa Lehtoranta,⁸ Tobias Recker,⁹ Seppo Salminen,¹⁰ Machteld Tiemessen,¹¹ and Maria Carmen Collado¹²

¹School of Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, United Kingdom; ²Institute for Life Sciences, University of Southampton, Southampton, United Kingdom; ³Imperial College London, London, United Kingdom; ⁴Department of Paediatrics and Paediatric Infectious Diseases, Institute of Child's Health, Sechenov First Moscow State Medical University (Sechenov University), Moscow, Russia; ⁵Inflammation, Repair and Development Section, National Heart and Lung Institute, Faculty of Medicine, Imperial College London, London, United Kingdom; ⁶FrieslandCampina, Amersfoort, The Netherlands; ⁷Institute of Public Health and Clinical Nutrition, University of Eastern Finland, Kuopio, Finland; ⁸IFF Health, International Flavors & Fragrances, Finland; ⁹ILSI Europe, Belgium; ¹⁰University of Turku, Finland; ¹¹Danone Nutricia Research, The Netherlands; and ¹²Institute of Agrochemistry and Food Technology-National Research Council (IATA-CSIC), Valencia, Spain

ABSTRACT

Food allergy (FA) affects the quality of life of millions of people worldwide and presents a significant psychological and financial burden for both national and international public health. In the past few decades, the prevalence of allergic disease has been on the rise worldwide. Identified risk factors for FA include family history, mode of delivery, variations in infant feeding practices, prior diagnosis of other atopic diseases such as eczema, and social economic status. Identifying reliable biomarkers that predict the risk of developing FA in early life would be valuable in both preventing morbidity and mortality and by making current interventions available at the earliest opportunity. There is also the potential to identify new therapeutic targets. This narrative review provides details on the genetic, epigenetic, dietary, and microbiome influences upon the development of FA and synthesizes the currently available data indicating potential biomarkers. Whereas there is a large body of research evidence available within each field of potential risk factors, there is a very limited number of studies that span multiple methodological fields, for example, including immunology, microbiome, genetic/epigenetic factors, and dietary assessment. We recommend that further collaborative research with detailed cohort phenotyping is required to identify biomarkers, and whether these vary between at-risk populations and the wider population. The low incidence of oral food challenge–confirmed FA in the general population, and the complexities of designing nutritional intervention studies will provide challenges for researchers to address in generating high-quality, reliable, and reproducible research findings. *Adv Nutr* 2021;00:1–19.

Statement of Significance: Food allergy affects the quality of life of millions of people worldwide and presents a significant psychological and financial burden for both national and international public health. Identifying reliable biomarkers that predict the risk of developing food allergy would be valuable in both preventing morbidity and mortality and by making current interventions available at the earliest opportunity. This review provides details on the genetic, epigenetic, dietary, and microbiome influences upon the development of food allergy. This helps in identifying reliable biomarkers to predict the risk of developing food allergy, which could be valuable in both preventing morbidity and mortality and by making interventions available at the earliest opportunity.

Keywords: IgE-mediated food allergy, biomarkers, pathways, risk factors, microbiota, nutrition, infant diet

Introduction

Food allergy (FA) is defined as an adverse immunological response to a food protein (1). It affects the quality of life of millions of people worldwide and presents a significant psychological (2) and financial (3) burden for both national and international public health. The European Academy

of Allergy and Clinical Immunology (EAACI) systematic review estimates FA prevalence in Europe at between 0.1% and 6.0% (4). Risk factors for developing FA are multiple and contextual, ranging from genetic predisposition to environmental factors (such as mode of birth delivery, type and timing of solid food introduction, changes in hygiene

TABLE 1 Summary of the most common and specific determinants impacting microbiota and risk of developing food allergy

	Factors associated with higher risk of food allergy	Factors associated with lower risk of food allergy	Factors with no association with higher/lower risk of food allergy
Factors increasing microbial dysbiosis	Antibiotic use during pregnancy and first year of life Cesarean delivery Exposure to bacterial enterotoxins Vitamin D deficiency	—	Formula feeding Low-fiber/high-fat diet
Factors improving microbial equilibrium	—	Farm/rural lifestyle Pet exposure in early life Having older siblings Exposure to an increased diversity of foods in early life Ingestion of aryl hydrocarbon receptor ligands (cruciferous vegetables) n-3 PUFAs Early oral exposure to foods	Outdoor activities Breastfeeding Probiotics/fermented products Less processed food
Factors with no proven impact on microbial dysbiosis/equilibrium	Early cutaneous exposure to food allergens in the environment Family history of allergic disease Prior diagnosis of atopic disease like eczema Higher socioeconomic status Living in developed societies	—	—

practices, and socioeconomic status) and the interaction between these factors (Table 1).

Identifying biomarkers that reflect either the risk of developing FA, the severity of FA, or induction of tolerance (i.e., reaching nonreactivity toward a substance that would previously cause a reaction) would be valuable in both preventing morbidity and mortality arising from FA, by allowing earlier interventions and by potentially highlighting new targets for intervention. The Health Biomarkers Definitions Working Group defined a biomarker as “a characteristic that is objectively measured and evaluated as an indicator

of normal biological processes, pathogenic processes, or pharmacologic responses to a therapeutic intervention” (5).

Biomarkers can also provide value in the regulatory context. The European Food Safety Authority health claim substantiation requires that “a food or one of its constituents significantly reduces a risk factor in the development of a human disease” (6). The regulation additionally requires that the risk factor is “generally accepted.” A classic example is cholesterol, a biomarker found to be associated with heart disease development. In labeling or advertising, health claims that constitute a “reduction of disease risk” shall also bear a statement indicating that the disease to which the claim is referring has multiple risk factors and that altering one of these risk factors might or might not have a beneficial effect. Thus, the optimal risk biomarker to be altered would be a combination of risk factors or a chain of events reflecting changes in the RR.

This article reviews available evidence in human studies in early life about well-described pathways with well-defined biomarkers and risk factors that are associated with IgE-mediated FA.

Current Status of Knowledge

Recent efforts have focused on the identification of biomarkers for prediction and diagnosis of IgE-mediated FA. IgE-mediated reactions induce a variety of symptoms that range from erythema, urticaria and angioedema, nausea, abdominal pain or vomiting, to severe respiratory distress, or cardiovascular collapse among others (7). Differences in the outcomes and manifestations might be related to genetic components but also to environmental factors, dietary factors, and the intestinal microbiota (8). The exact

Funding for this work was provided by the European branch of the International Life Sciences Institute (ILSI, Europe), as this work was developed by an expert group of ILSI Europe.

Author disclosures: This work was conducted by an expert group of the European branch of the International Life Sciences Institute, ILSI Europe. The research question addressed in this publication and potential contributing experts in the field were identified by the Nutrition, Immunity & Inflammation Task Force. Members of this task force are listed on the ILSI Europe website at www.ils.eu/task_forces/nutrition/nutrition-immunity-and-inflammation/.

According to ILSI Europe policies, the expert group is composed of $\geq 50\%$ of external nonindustry members. The research project was handed over to the experts to independently refine the research question and carry out the work, that is, collecting/analyzing data/information and writing the scientific paper independently of other activities of the Task Force. The research reported is the result of a scientific evaluation in line with ILSI Europe's framework to provide a precompetitive setting for public-private partnership. Experts are not paid for the time spent on this work. However, the nonindustry members within the expert group were offered support for travel and accommodation costs from the Task Force to attend meetings to discuss the manuscript and a small compensatory sum (honorarium) with the option to decline. For further information about ILSI Europe, please email info@ilsieurope.be. CEC, DM, LU, and CG-G* have shared authorship and all of them contributed equally.

Address correspondence to MCC (e-mail: mcolam@iata.csic.es; publications@ilsieurope.be). Abbreviations used: CHILd, Canadian Healthy Infant Longitudinal Development; CMA, cow milk allergy; C11orf30, chromosome 11 open reading frame 30 (EMSY transcriptional repressor, BRCA2 interacting); EAACI, European Academy of Allergy and Clinical Immunology; EAT, Enquiring About Tolerance; FA, food allergy; FOXP3, forkhead box P3; HLA, human leukocyte antigen; LEAP, The Learning Early about Peanut Allergy; OIT, oral immunotherapy; RCT, randomized controlled trial; STAT, signal transducer and activator of transcription; Treg, regulatory T cell.

diagnosis and prevalence of FA is difficult to ascertain due to the imprecision of laboratory tests and the lack of specific biomarkers, relying on the combination of the clinical history of characteristic symptoms together with test results (7), the use of IgE as a biomarker in FA, and the potential associations with genetic and epigenetic origins that would be targets of potential interventions (breast milk compared with others, weaning, diet, etc.).

Genetic and epigenetic biomarkers of FA

The link between the risk of FA in children and allergic diseases and/or allergic sensitization in their family has been extensively reported (9–14), with estimates that FA/sensitization risk doubles if 1 parent has an allergic disease, and is 3-fold higher if both parents have an allergic disease. A meta-analysis of genome-wide association studies identified 10 loci in or near *TLR6* (toll like receptor 6), *C11orf30* (EMSY transcriptional repressor, BRCA2 interacting), *STAT6* (signal transducer and activator of transcription 6), *SLC25A46* (solute carrier family 25 member 46), *HLA-DQB1* (human leukocyte antigen DQ isotype B1), *IL1RL1* (interleukin 1 receptor like 1), *LPP* (LIM domain containing preferred translocation partner in lipoma), *MYC* (MYC proto-oncogene, bHLH transcription factor), *IL2* (interleukin 2), and *HLA-B* (major histocompatibility complex, class I, B), that are associated with allergic sensitization (15). Allergen-specific genetic modifications in the HLA DR and DQ isotype gene region have also been associated with peanut allergy (16). Conflicting results were reported with regard to gender association with FA and no conclusive studies are available (10, 11, 17, 18). Some data suggest that 5 loci at genome-wide significance (clade B serpin, or SERPINB) gene cluster at 18q21.3, the cytokine gene cluster at 5q31.1, the filaggrin gene, the *C11orf30/LRRC32* (leucine rich repeat containing 32) locus, and the HLA region increase the risk of FA (19).

Eczema and FA often coexist, and evidence suggests that an impaired skin barrier is a significant risk factor for FA development later in life (20, 21) with loss-of-function variants in the filaggrin gene suggested as a causative factor; moreover, filaggrin mutation is associated with eczema and asthma later in life (22, 23). Identified genetic loci associated with FA, their potential mode of action, and evidence supporting their use as biomarkers are presented in **Table 2**.

Extrinsic environmental factors including diet, pollutants, and infections, and intrinsic factors such as the intestinal microbiota and inflammatory state are likely to play a crucial role in inducing epigenetic changes (24, 25). Postnatal factors and environmental influence are risk factors for FA development and this exposure accumulates while the infant develops (9, 10, 18). The route of exposure (e.g., placental, skin, breast milk, airway, gut), timing, dose of allergen exposure, and host immune system status are likely to impact upon the potential for epigenetic change (26). Investigations of targeted and untargeted methylation profiles of immune cells are methodologies that can help to find biomarkers

that reflect the different stages of FA: those at risk, those who are tolerant, and those with active disease (27, 28). An overview of studies on epigenetic changes associated with FA is presented in **Table 3**.

The role of breastfeeding, and time of food introduction in FA

Breastfeeding.

Human milk is the first food available to a newborn baby, and exclusive breastfeeding for the duration of 6 mo is recommended by the WHO. Available evidence suggests that breastfeeding protects against infections as well as offering long-term benefits, reducing the risk of hypertension and diabetes, and improving cognitive development (29). The protective effect of breast milk on allergy development has not been fully demonstrated (29–33). However, there are conflicting data concerning the relation between breastfeeding and FA, with some cohort studies reporting a reduced risk of FA development in the general population (20, 21) and in high-risk children (34) and others reporting an increased risk (35, 36). One meta-analysis investigating this relation reported no evidence of breastfeeding's protective effect in preventing FA development (OR: 1.02; 95% CI: 0.88, 1.18), although the authors suggested that the risk of bias and major differences in the outcome definitions in the current studies might be responsible for the inconclusive results (31). Because human milk contains food proteins, their concentrations in the milk and maternal diet might also contribute to tolerance development (37), particularly in the presence of the biologically active molecules (38). Both aspects are not normally considered in the studies assessing associations between breastfeeding and noncommunicable diseases development.

A recent systematic review on FA prevention suggests that although breastfeeding has many benefits for infants and mothers, it might not reduce the risk of FA (39). Human breast milk constituents vary (over time postpartum, within and between women, and even within the same feed), which could, in part, explain some of the conflicting results of general observational studies regarding the provision of breastfeeding (40, 41). It has been described that immunological compounds in breast milk (including cytokines and Igs) are modulated by multiple factors, including maternal allergic status, parity, and geographical location among others (42–45), but overall evidence on the topic is conflicting with most of the studies not identifying clear associations between the immunological composition of breast milk and allergic disease development in infants (38). Dietary peptides from proteins in food are excreted in breast milk, but these have relatively short sequences and are in small amounts; therefore, their sensitization or tolerogenic potential remains to be explored (46). The presence of specific peptides has also been shown in infant formula (47). However, so far, systematic reviews (48, 49) have not found sufficient evidence that hydrolyzed formula prevents eczema or milk allergy (50).

TABLE 2 Genetic loci associated with food allergy, their potential link with food allergy, and evidence supporting their use as biomarkers¹

Name	Genetic risk factor	Role	Potential link with FA	Reported utility as biomarker?	References
Toll-like receptor 6	<i>TLR6</i>	Pathogen recognition and activation of innate immunity	TLR function can be altered by early environmental and microbial exposures	Generally associated with allergic sensitization	(15, 145)
EMSY transcriptional repressor	<i>Ct1orf30</i>	Repressor of BRCA2 protein	Involved in epigenetic regulation of gene expression	Identified as genetic risk factor for peanut allergy and food allergy	(15, 146)
Signal transducer and activator of transcription 6	<i>STAT6</i>	Central role in IL4-mediated responses	Polymorphisms have been associated with age of tolerance induction	Age of tolerance development for cow milk was significantly higher in children with the GG genotype at rs324015 of the <i>STAT6</i> gene compared with those with the AA + AG genotype [2 y (range = 1.5–3.9 y) vs. 1.2 y (range = 1.0–2.2 y); <i>P</i> = 0.02]	(15, 147)
Solute carrier family 25 member 46	<i>SLC25A46</i>	Promotes mitochondrial fission and prevents the formation of hyperfilamentous mitochondria	Involved in the association between food allergy and atopic dermatitis	Polymorphism <i>SLC25A46</i> was associated with higher risk of food allergy	(15, 148)
Major histocompatibility complex, class II, DQ beta 1	<i>HLA-DQB1</i>	Plays a central role in the immune system by presenting peptides derived from extracellular proteins	Peanut allergic-specific loci in the human leukocyte antigen (HLA)-DQ and -DR regions were found in a large cohort study	Several polymorphisms associated with peanut, milk, and egg allergy	(15, 16, 149)
Interleukin 1 receptor-like 1	<i>IL1RL1</i>	Involved in the function of helper T cells	ST2, β -chain of IL33 receptor	Generally associated with allergic sensitization	(15)
LIM domain containing preferred translocation partner in lipoma	<i>LPP</i>	Involved in cell-cell adhesion and cell motility. This protein also shuttles through the nucleus and may function as a transcriptional coactivator	Allergic sensitization	Generally associated with allergic sensitization	(15)
MYC proto-oncogene, bHLH transcription factor	<i>MYC</i>	Plays a role in cell cycle progression, apoptosis, and cellular transformation	Downregulated in children with food allergy	Generally associated with allergic sensitization and food allergy	(15, 150)
Interleukin 2	<i>IL2</i>	Proliferation of T and B lymphocytes	Allergic sensitization	Generally associated with allergic sensitization	(15)
Major histocompatibility complex, class I, B	<i>HLA-B</i>	Central role in the immune system by presenting peptides derived from the endoplasmic reticulum lumen	Allergic sensitization	Generally associated with allergic sensitization	(15)

(Continued)

TABLE 2 (Continued)

Name	Genetic risk factor	Role	Potential link with FA	Reported utility as biomarker?	References
Filaggrin	<i>FLG</i>	Role in skin barrier function	Indirect association with food allergy	Filaggrin loss-of-function mutations are associated with food allergy in older children via eczema and food allergen sensitization in their early childhood	(20, 21, 23, 151)
Interleukin 13	<i>IL13</i>	Involved in several stages of B-cell maturation and differentiation	IL13 polymorphism rs1295686 (in complete linkage disequilibrium with functional variant rs20541) is associated with challenge-proven food allergy	IL13 gene polymorphisms have also been identified as biomarkers of IgE-mediated food allergy and are a predictor of cord blood IgE concentrations	(152)
Catenin alpha 3	<i>CTNNA3</i>	Cell-cell adhesion	Knockdown of <i>CTNNA3</i> resulted in upregulation of <i>CD63</i> and <i>CD203c</i> in mononuclear cells upon PMA stimulation	Copy number variation impacting <i>CTNNA3</i> has been associated with pediatric food allergy	(153)
RNA binding fox-1 homolog 1	<i>RBFOX1</i>	Regulates alternative splicing events	Association with food allergy at a genome-wide scale	Generally associated with pediatric food allergy	(153)
GC vitamin D binding protein	<i>GC/DBP</i>	Binds to vitamin D and its plasma metabolites and transports them to target tissues	GG genotype produces less vitamin D binding protein (DBP)	Vitamin D deficiency linked with GG genotype producing less vitamin D binding protein (DBP) was associated with a higher prevalence of egg and peanut allergy in 1- and 2-y-old infants	(154)
Indoleamine 2,3-dioxygenase 1	<i>IDO1</i>	Modulates T-cell behavior	High IDO activity is associated with nonresponsiveness to food allergens despite allergen sensitization	Associated with tolerance to food allergens	(155)
Sirtuin 1	<i>SIRT1</i>	Functions of human sirtuins have not yet been determined	Negatively regulates FcεR1-stimulated mast cell activation and anaphylaxis	Generally associated with antiallergic response	(156, 157)

¹BRCA2, Breast Cancer Type 2 susceptibility protein; FcεR1, high-affinity IgE receptor; PMA, phorbol myristyl acetate.

TABLE 3 Epigenetic changes associated with food allergy¹

Study	Where identified	Main findings	Potential mechanism of action	Reported utility as biomarker?	References
DNA methylation profiles (~450,000 CpGs) of peripheral immune cells (CD4+ T cells)	Children with IgE-mediated food allergy	179 differentially methylated sites of loci associated with the disease phenotype, and 96 CpG sites DNA methylation profile discriminated food-allergic vs. healthy infants	MAP kinase pathway → dysregulation of DNA methylation at MAPK signalling-associated genes during early CD4+ T-cell development may contribute to suboptimal T-lymphocyte responses in early childhood associated with the development of food allergy Methylation of metabolic (<i>RPTOR</i> , <i>PIK3CD</i> , <i>MAPK1</i> , <i>FOXO1</i>) and inflammatory genes (<i>IL1R</i> , <i>IL18RAP</i> , <i>CD82</i>) affected	Predicted clinical outcomes with an accuracy of almost 80% MAP kinase pathway was most prominently associated with CpGs that were predictive of food challenge	(158, 159)
DNA methylation profiles	Egg allergy	DNA methylation profiles of T cells discriminate infants with persistent egg allergy compared with those who had outgrown egg allergy Cow milk allergic infants showed hypermethylation in whole blood compared with controls and tolerant group		Data about predictive potential not available	(150)
DNA methylation profiles	Cow milk allergy		Differential methylation patterns on <i>DHX58</i> (innate immune response), <i>ZNF287</i> (transcriptional regulation), <i>EIF42A</i> (interferon pathway), and <i>HTRA2</i> (smooth muscle contraction) between groups	Data about predictive potential not available	(160)
DNA methylation profiles and single-nucleotide polymorphisms	Peanut allergy	DNA methylation of the <i>HLA-DQB1</i> and <i>HLA-DRB1</i> , <i>IL4</i> , <i>IL12B</i> , <i>IL2</i> , <i>BDNF</i> , <i>IL17F</i> , <i>CXCL12</i> , <i>CCR7</i> , runt-related transcription factor 1 (<i>RUNX1</i>), <i>CD3e</i> , and <i>SERPINE1</i> <i>IL1B</i> and <i>IL6</i> has been associated with peanut allergy	Increased protein secretion in response to allergen-specific stimulation Additional functional studies are needed	DNA methylation signature combinations may have superior diagnostic potential than serum peanut-specific IgE	(16)
Th1-Th2	Cow milk allergy	DNA methylation profiles differ with cow milk allergy	DNA methylation profiles of <i>IL4</i> , <i>IL5</i> , <i>IL10</i> , and <i>IFNγ</i> genes between infants with active cow milk allergy and those who outgrew their cow milk allergy	<i>GATA3</i> in Th2 cells Ex vivo PBMC cytokine profile in predicting cow milk allergy: TNF, IL10, IL12 higher in cow milk allergy patients compared with controls	(161–163)

(Continued)

TABLE 3 (Continued)

Study	Where identified	Main findings	Potential mechanism of action	Reported utility as biomarker?	References
Th1-Th2	Cow milk allergy	DNA methylation of FOXP3, Th1/Th2 cytokine genes in IgE-mediated allergy, in children with cow milk allergy treated with an extensively hydrolyzed formula including a probiotic (test formula) vs. a control formula	FOXP3, IL10, and IFN γ demethylation rate was higher, and IL4 and IL5 demethylation rate was lower in the test formula group	Intervention promotes regulatory and immune suppressive immune factors and at the same time decreases activity of Th2 type genes	(164)
FOXP3	Peanut-allergic infants and cow milk-allergic infants	Immune-tolerant participants had \uparrow ai-Treg with greater suppressive function, and with \uparrow FOXP3 hypomethylation	Oral immunotherapy in peanut allergic infants increased antigen-induced regulatory T-cell function and hypomethylation of FOXP3 in infants that became tolerant	Data about predictive potential not available	(165)
	Cow milk allergy	\downarrow FOXP3 gene demethylation in children with active IgE-mediated cow milk allergy	Formula selection influenced the FOXP3 T-cell-specific demethylation region	Data about predictive potential not available	(166)
Methylation levels taken from mononuclear blood cells at 405,658 CpG islands across the genome (machine learning approach)	40 samples for training, 10 samples for cross-validation, and 8 completely hidden samples for testing	Novel 13-gene signature to diagnose clinical reactivity: chr1p13 (SARS), chr7p22 (MAFK), chr11q14 (PANX1), chr9p22 (SLC24A2), chr8p21 (KIF13B), chr10q26 (CTBP2), chr10q11 (ARID5B), and chr10q23 (FAM190B)	These genes are mapped to several canonical Wnt pathways, GO, and positional gene sets with functional association with the immune system	The 18-CpG signature mapped to 13 genes is a strong biomarker of FA with a 94–96% accuracy	(167)

^aai, antigen-induced; ARID5B, AT-rich interaction domain 5B; BDNF, brain-derived neurotrophic factor; CCR7, C-C motif chemokine receptor 7; chr, chromosome; CTBP2, C-terminal binding protein 2; CXCL12, C-X-C motif chemokine ligand 12; DHX58, DEXH-box helicase 58; EIF4A2, eukaryotic translation initiation factor 4A2; FA, food allergy; FOXO1, forkhead box O1; FOXP3, forkhead box P3; GATA3, GATA binding protein 3; GO, The Generic Gene Ontology; HLA, human leukocyte antigen; HTRA2, Htra serine peptidase 2; KIF13B, kinesin family member 13B; MAFK, MAF bZIP transcription factor K; MAPK1, mitogen-activated protein kinase 1; PANX1, pannexin 1; PBMCs, peripheral blood mononuclear cells; PK3CCD, phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit delta; RPTOR, regulatory associated protein of MTOR complex 1; SARS, seryl-tRNA synthetase; SLC24A2, solute carrier family 24-member 2; Th, helper T cells; Treg, regulatory T cells; ZNF281, Zinc Finger Protein 281.

Thus, claims currently appearing on infant formula products need better substantiation and many reputable organizations, including the American Academy of Pediatrics; American Academy of Allergy, Asthma, and Immunology; American College of Allergy, Asthma, and Immunology; and Canadian Society for Allergy and Clinical Immunology, concluded that “there is no protective benefit from the use of hydrolyzed formula in the first year of life against food allergy or food sensitization” (51, 52). A recent study suggested that avoiding temporary supplementation with conventional cow milk formula in the first 3 d of life can result in a large decrease in the risk of FA in early childhood (53), but this requires further confirmation.

Weaning and food introduction.

Delaying the introduction of solid food until 6 mo remains the current WHO recommendation. Yet recent expert opinion has investigated the hypothesis that oral tolerance can be induced by modifying the timing and diversity of early food exposure (54). Supportive data for this hypothesis are coming predominantly from 2 large high-quality randomized controlled trials (RCTs), The Learning Early about Peanut Allergy (LEAP) and Enquiring About Tolerance (EAT). The LEAP study demonstrated a significant reduction in peanut allergy prevalence in children at high risk of allergy development, who were consuming peanuts between 4 and 11 mo old on a regular basis (55). There was an earlier and greater increase in peanut-specific IgG and IgG4 in the early consumption group compared with the avoidance group. In both groups the mean peanut-specific IgE concentrations were highly comparable and increased over time, albeit there were more participants in the avoidance group with very high IgE concentrations (55). The EAT trial looked at early food introduction (from 3 mo old) and concluded that it might decrease the risk of FA development (56). The authors reported significantly lower RRs of peanut and egg allergy in the early introduction group, with no difference in the prevalence of milk, sesame, fish, or wheat allergy. Risk reduction was shown in per protocol analysis only, whereas no statistically significant difference was found in intention-to-treat analysis. Studies reporting contradictory results to EAT exist (57), but they are often considered of lower robustness.

With an apparent shift in expert opinion toward early introduction of certain highly allergenic foods, the American National Institute of Allergy and Infectious Diseases updated its guidelines on peanut allergy prevention in 2017 (58), recommending that peanut-containing food introduction should occur between 4 and 6 mo of age in egg-allergic infants and/or babies with severe eczema, and at 6 mo of age for infants with mild-to-moderate eczema. Recent guidelines from the American Academy of Pediatrics support these recommendations (52).

In their systematic review on FA prevention, the authors concluded that available evidence suggests that “introduction of small amounts of cooked egg into the infant diet as part of complementary feeding probably reduces

the risk of egg allergy in infancy and in countries with a high prevalence of peanut allergy, introducing regular peanut consumption from 4–11 months of life in infants at increased risk probably results in a large reduction in peanut allergy in early childhood compared to completely avoiding peanut for the first five years” (39). In contrast, no reduction in FA incidence was found when multiple potential food allergens were simultaneously introduced into the infant diet from age 3 mo (56). Diet diversity during the first year of life might also have a positive role in determining the risk of FA. An increased diversity of complementary foods introduced in the first 12 mo of life was inversely associated with FA development up to 6 y old (59).

Is there a need for biomarkers to monitor dietary interventions to induce tolerance?

Food avoidance remains the main therapeutic approach in FA management, but researchers and clinicians are continuously seeking for intervention options. Controlled exposure to the allergens was suggested as a potential option for tolerance induction. Indeed, in recent years, oral immunotherapy (OIT) has been applied for several allergens to investigate whether desensitization and/or sustained unresponsiveness development is possible. A meta-analysis on the effect of OIT in reducing prevalence of cow milk allergy (CMA) concluded it is an effective therapy (60); however, frequency of adverse events is high and validity of outcome selection used to measure the efficacy of OIT is still unclear. Looking at an individual study level, there was no association of OIT in children (aged 6–17 y) and IgE concentrations between the treated and the control group, whereas IgG4 was significantly increased in the posttreatment group after OIT but there was only a slight increase in the control group (61). Recently, a cohort of 137 peanut-allergic child and adult patients (aged 6–26 y) were compared with non-peanut-allergic controls and differences between IgE, IgG4, and the ratio of IgG4/IgE were examined (62).

These observations would imply that more data are needed on specific immunoglobulin E (sIgE) and IgG4 in monitoring tolerance induction over time before it can be concluded that these are reliable biomarkers for tolerance induction. There could be more potential for the increase in IgG4 in oral tolerance induction than the decrease in IgE. It is very important to note that there are no agreed core outcome measures in FA trials, which do not allow for appropriate effectiveness/efficacy evaluation (63). Different immunological parameters are currently used as end points in OIT trials, but available evidence of their importance is very limited (64).

What Is the Role of the Microbiota in FA?

A link between IgE-mediated FA and the gut microbiota composition and metabolic activity has been suggested. A recent study including 233 infants (>4 y old) with FA (milk, sesame, peanut, and tree nuts), and nonallergic controls showed a distinct microbial profile for FA to different foods




FIGURE 1 Level of evidence of different biomarkers and interactions between genetic background, lifestyle, and epigenetics factors on the interplay between microbiota and the immune system on food allergy. CD103, integrin α E β 7; CX3CR1, C-X3-C motif chemokine receptor 1; FOXP3, forkhead box P3; LAP, latency associated peptide; mMCP-1, mouse mast cell protease-1; PPAR, peroxisome proliferator-activated receptors; TGF- β , transforming growth factor β ; Th, helper T cell; TLR, Toll-like receptor; Treg, regulatory T cell.

characterized with an underrepresentation of *Prevotella copri* (65). In agreement, maternal carriage of *Prevotella copri* during pregnancy was also linked to a decreased risk of FA during infancy (66). Growing evidence supports a role for the gut microbiome in the pathogenesis and course of FA, with microbial dysbiosis preceding the development of FA (67). It has been reported that an elevated *Enterobacteriaceae/Bacteroidaceae* ratio in early infancy as well as lower microbial species richness in the infant ($n = 166$, ages 3 and 12 mo) might be a predictor of egg, milk, and peanut sensitization (determined by skin prick test) at age 12 mo, adjusting for birth delivery mode, antibiotic use, or breastfeeding (68). This raises the question of whether FA can be predicted using gut microbiome biomarkers (69). A study with 319 subjects enrolled in the Canadian Healthy Infant Longitudinal Development (CHILD) study showed that infants at risk of asthma exhibited transient gut

microbial dysbiosis during the first 100 d of life characterized by lower relative abundance of *Lachnospira*, *Veillonella*, *Faecalibacterium*, and *Rothia* species (70). Another study reported lower relative abundance of *Citrobacter*, *Oscillospira*, *Lactococcus*, and *Dorea* in stool samples collected at age 3–6 mo in children who had FA (milk, egg, peanut, wheat, soy, or other nut allergy) by the age of 3 y (71). In addition, *Firmicutes*, including clostridia, were enriched in the gut microbiota of infants at age 3–6 mo whose milk allergy resolved by 8 y of age (72), suggesting a potential predictive role of gut microbiota composition for FA.

Interestingly, the specific microbiota signature can distinguish infants with IgE-mediated from non-IgE-mediated FA. Infants with IgE-mediated FA had increased concentrations of (*cluster I*) and *Anaerobacter* and decreased concentrations of *Bacteroides* and *Clostridium cluster XVIII*, with a positive correlation between *Clostridium sensu stricto* and serum

sIgE (73). However, as with observational studies, it is not possible to assess causation between changes in microbial composition and FA (74). A study in adults with FA showed the opposite results with reduced *Clostridiales*, and increased *Bacteroidales* (75), suggesting that the changes observed in microbiota associated with allergy can be different depending on other factors such as age, ethnicity, geographical location, and lifestyle.

It is widely known that the early infant microbiota is influenced by several factors, including mode of birth, antibiotic use, and environmental exposures, that can contribute to the dysbiosis linked to allergy development (Figure 1) and would provide opportunities to develop strategies aimed at microbial modulation and decreasing the risk of FA (76).

C-section delivery and antibiotic exposition

Available evidence indicates that C-section is a possible risk factor for FA because the newborn infant bypasses the microbial exposure happening naturally during vaginal delivery, whereby a distinct gut microbiota is obtained (77). In general, infants born by C-section have lower concentrations of *Bacteroides* and lower diversity, which is a pattern also observed to precede the development of allergic symptoms in several studies (78). However, there is no clear evidence on C-section association with a higher risk of FA development, with studies producing contradictory results (79, 80). However, a 7-fold increased risk of parental-reported fish or nut allergy and a 4-fold increased risk of confirmed egg allergy were reported (81) in high-risk children born via C-section. C-section was found to be associated with other allergic diseases, such as allergic rhinitis (OR: 1.23; 95% CI: 1.12, 1.35), asthma (OR: 1.18; 95% CI: 1.05, 1.32), and allergic sensitization to foods (OR: 1.32; 95% CI: 1.12, 1.55) (82). Most of the C-sections are associated with antibiotic intrapartum. Antibiotic use (particularly cephalosporins and sulfonamides), including its frequency during pregnancy and first year of life, was linked with an increased risk of FA development (83), and is likely to reflect an indirect effect via infant gut microbiota dysbiosis (84, 85).

Breastfeeding practices

It has been shown that infants with CMA had an increased gut microbiota diversity and a higher prevalence of members belonging to the *Lachnospiraceae* family (*Firmicutes* phylum) compared with nonallergic infants (86). However, another study showed an inverse association between the early gut microbial diversity and the risk of allergic sensitization (87). A low gut microbiota richness, overrepresentation of *Enterobacteriaceae*, and underrepresentation of *Bacteroidaceae* (*Bacteroidetes* phylum) at 3 mo of age were associated with food sensitization in a subset of the CHILD study (68). Those associations were found in infants who were vaginally delivered, exclusively breastfed, and unexposed to antibiotics.

Breastfeeding practices were associated with lower diversity and higher concentrations of *Bifidobacterium breve*

and *B. bifidum* (*Actinobacteria* phylum), and the cessation of breastfeeding resulted in faster maturation of the gut microbiota, as marked by an increase in the members belonging to the *Firmicutes* phylum (88). However, formula-fed infants had a more diverse microbiota with higher proportions of *Clostridium* spp. (*Firmicutes* phylum), and *Enterobacteriaceae* members (*Proteobacteria* phylum), but with lower bacterial count (89). Recent studies have shown that breast milk with a reduced microbial richness in the first month of life could play an important role in allergy development during childhood (90). Thus, the protection against allergy development provided by human milk might be attributable to the effect on the infant gut microbiota or direct effects on immune system; however, further studies are needed to evaluate the effect of breastfeeding and milk-specific compounds on FA (91).

Environmental exposures

Associations between living in affluent countries and allergic disease development are well known, and FA is no exception to the rule. A higher socioeconomic status (92) or living in developed societies were associated with an increased risk of FA development, although it is possible that variations in frequencies of studies and methodological variation also contribute to these geographic variations (4). Researchers suggest that farming lifestyle exposes pregnant women and their offspring to a wide variety of microorganisms, which urban inhabitants lack. Data from 2 large, prospective cohorts showed that exposure to a greater variety of environmental microorganisms was associated with a reduced risk of asthma development in "Prevention of Allergy—Risk Factors for Sensitization Related to Farming and Anthroposophic Lifestyle" (PARSIFAL study) (OR: 0.62; 95% CI: 0.44, 0.89) and in Multidisciplinary Study to Identify the Genetic and Environmental Causes of Asthma in the European Community (GABRIEL) Advanced Study (OR: 0.86; 95% CI: 0.75, 0.99) (93).

Dietary Interventions

Macronutrient and micronutrient associations with FA

A recent systematic review suggested that supplementation with fish oil [a source of long-chain omega-3 (n-3) fatty acids] during pregnancy and lactation can reduce risk of allergic sensitization to egg (RR: 0.69; 95% CI: 0.53, 0.90; I² = 15%; absolute risk reduction: 31 cases per 1000; 95% CI: 10, 47) (94). The Grading of Recommendations Assessment, Development and Evaluation certainty of these findings was moderate. In addition, in vitro and in vivo studies have demonstrated that n-3 PUFAs can modulate the activity of dendritic cells, T cells, and IgE production by B cells, reducing allergic sensitization (95).

Although vitamin D deficiency was linked with the development of allergic diseases (96), data relevant for FA are limited. Vitamin D deficiency linked with GG genotype producing less vitamin D binding protein was associated with

a higher prevalence of egg and peanut allergy in 1- and 2-year-old infants (97). Use of vitamin D supplements during pregnancy to prevent FA was, however, unsuccessful, both in an RCT (RR: 1.92; 95% CI: 0.57, 6.50) (98) and a case-control study (OR: 1.50; 95% CI: 0.78, 2.88) (99). Supplementation during the first year of life resulted in a reduced risk of FA development during the first 12 mo of life (RR: 0.49, 95% CI: 0.27, 0.88) (99). However, the confidence in this estimate is also very low owing to indirectness of the evidence and risk of bias, as reported in a recent systematic review on the subject (100). Overall, there is currently not enough evidence to suggest that vitamin D supplements for pregnant and/or breastfeeding women or infants have an effect on FA development (39).

Dietary interventions targeting microbiota modulation: prebiotics and probiotics

Targeted and personalized nutrition is an emerging strategy to approach FA in early infancy including microbiome-modifying interventions with probiotics (*Lactobacillus acidophilus* LAVRI A1, *Lactocaseibacillus rhamnosus* GG), prebiotics (long-chain fructo-oligosaccharides, short-chain galacto-oligosaccharides), and human milk oligosaccharides (2'-fucosyllactose, lacto-*N*-neotetraose) (101). The pathogenesis of FA in early infancy and other associated events such as dermatitis or asthma are still largely unknown, but increasing evidence suggests that they are associated with a perturbation of the gut microbiome, or microbial dysbiosis, leading to alterations in the immune system that could influence the occurrence of FA (102). In addition, FA derives from a defect in immune tolerance mechanisms. Immune tolerance is modulated by gut microbiota composition and function. Therefore, the potential use of probiotics has been highlighted to counteract microbial dysbiosis linked to FA and boost microbially modulated tolerance because probiotics could interact with the host microbiota and the host immune system at the same time (103). In infants, supplementation with specific probiotic strains might reduce the risk of sensitization to cow milk (RR: 0.60; 95% CI: 0.37, 0.96) (104) although the quality of evidence is considered low. In general, those studies combined maternal and infant supplementation, and it is unclear if the effect was due to the combination or the specific intervention (104–106). A systematic review and meta-analysis was published recently, suggesting that probiotic intake during late pregnancy and lactation might reduce the risk of eczema (RR: 0.78; 95% CI: 0.68, 0.90; I^2 : 61%; absolute risk reduction: 44 cases per 1000; 95% CI: 20, 64) (94). There are some studies associating the consumption of oligosaccharides in early life with reduced incidence of atopic dermatitis and other allergy manifestations (107, 108) with a lack of evidence in FA and human studies. However, the evidence on the use of prebiotics, probiotics, and synbiotics in breastfeeding mothers and infants to reduce the risk of FA is inconclusive (39). In an RCT, specialized infant formula enriched with fructo-oligosaccharides and *Bifidobacterium breve* M-16V could restore altered microbiota in non-IgE-mediated cow

milk-allergic infants bringing it close to the healthy breastfed microbial profile when compared with the same formula without the synbiotic (109). Increasing evidence suggests that shifts in the neonatal gut microbiota composition, activity, and diversity are implicated in the pathogenesis of FA (Table 4).

Evidence for the Role of Microbial Metabolites in FA

Increasing data are showing the key role of metabolites in the host–microbe interaction as messengers and signals between the microbiota and the immune system with an impact on human health. A comprehensive understanding of how microbiota-derived metabolites influence the human immune system and health is critical for the rational design of therapies for microbiota-driven diseases (110). Different dietary patterns change the proportions and type of microbial groups, influencing host exposure to microbial metabolites (111), which in turn produce epigenetic changes. Although no data are available for infants in their first year of life, in older children and adults, a balanced low-fat and high-fiber diet could be important in preventing perturbation of the gut microbiome and preserving a functional immune system (112). Little is known about the role of microbial metabolites in FA but evidence is showing the impact of diet including prebiotics on the production of microbial metabolites such as SCFAs, polyamines, and even other compounds as toxins (LPS, staphylococcal enterotoxin B, etc.).

SCFAs

Metabolites produced by intestinal microbiota, and in particular SCFAs, play a critical role in mediating the effect of the gut microbiota on regulatory T-cell (Treg) proliferation and differentiation both in vitro and in vivo (113). The molecular mechanisms for this are not clearly elucidated but butyrate can suppress NF- κ B and STAT1 activation and induce differentiation of colonic Treg cells by enhanced histone acetylation (113–116). Moreover, these effects are not confined to the gastrointestinal tract, and both butyrate and propionate have been reported to influence peripheral Treg development (117). The mechanisms involved in SCFA regulation of T-cell differentiation would include the control of cellular metabolism and the G-protein-coupled receptor signaling pathways (118), and involve strong epigenetic regulation through inhibition of histone deacetylases (102). In particular, the effect of butyrate on Treg differentiation could be through the increase of histone H3 acetylation in the *FOXP3* locus (117), and propionate seems to increase the expression of *FOXP3* and *IL10* (119). These results could explain the benefits of dietary fiber and bacteria, such as *Akkermansia muciphila*, *Faecalibacterium prausnitzii*, *Eubacterium*, *Bifidobacterium*, *Clostridium*, and *Ruminococcus*, typical SCFA producers, that can increase colonic luminal SCFA concentrations and modulate the immune system response (120, 121).

Some specific SCFAs have been reported to influence FA. In detail, butyrate has a well-known inhibitory effect on

TABLE 4 Perinatal probiotics, prebiotics, and synbiotics for cow milk allergy management and allergy prevention: summary of clinical studies and meta-analyses¹

Strain(s)	No. subjects	Intervention time	Target	Outcome(s)	Study type	Reference
Clinical studies <i>Lactobacillus</i> LGG 1×10^9 CFU	100 infants diagnosed with CMA	4 wk	Management of CMA	Significant improvement in symptoms of infants diagnosed with CMA No impact on abdominal pain, constipation, and dermatitis	Randomized, double-blind, placebo-controlled study	(168)
Synbiotic formula with a combination of <i>Bifidobacterium breve</i> M-16V and chicory-derived neutral oligofructose, long-chain inulin	122 infants [Synbiotic $n = 35$; Control $n = 36$; Reference $n = 51$]	8 wk	Management of severe or complex non-IgE-mediated CMA	↑% of <i>Bifidobacterium</i> and ↓% of <i>Eubacterium rectale/Clostridium coccoides</i> group in the test group No significant results for the fecal secretory IgA and SCFAs	Double-blind, randomized clinical trial with nonrandomized breastfed reference group	(169)
<i>Lactobacillus rhamnosus</i> and <i>Bifidobacterium animalis</i> ssp. <i>lactis</i>	290 infants aged ~10 mo [Probiotic $n = 144$; Placebo $n = 146$]	6 mo	Allergic diseases and sensitization	↓Incidence of eczema No effect on the incidence of asthma and conjunctivitis or sensitization	Randomized, double-blind, placebo-controlled intervention	(170)
Amino acid-based formula (AAF) with fructo-oligosaccharides and <i>Bifidobacterium breve</i> M-16V <i>Lactobacillus rhamnosus</i> GG; <i>L. rhamnosus</i> LC705 (DSM 7061), <i>Bifidobacterium breve</i> Bb99 (DSM 13,692), and <i>Propionibacterium freudenreichii</i> ssp. <i>shermanii</i> JS (DSM 7076)	51 infants aged <13 mo [Test $n = 35$; Control $n = 36$] 891 mothers with infants at high risk of allergy [Probiotic $n = 445$; Placebo $n = 446$]	Infant intervention for 8 wk Maternal-infant intervention Follow-up until 5 y	Management of infants with suspected/proven CMA Allergy prevention	↑ <i>Bifidobacterium</i> in the AAF with prebiotic and probiotic ↓IgE-associated allergic disease occurred in cesarean-delivered children No allergy-preventive effect that extended to age 5 y	Randomized, double-blind, placebo-controlled intervention Randomized, double-blind, placebo-controlled study	(109) (171)
Meta-analyses Different strains	10 RT; $n = 845$ infants [Probiotics $n = 422$; control $n = 423$]	Different intervention times	Management of infants with suspected/proven CMA	No impact on hematochezia In confirmed CMA, probiotics ↑acquisition of tolerance to CMA at the end of 3 y	Meta-analysis	(172)

(Continued)

TABLE 4 (Continued)

Strain(s)	No. subjects	Intervention time	Target	Outcome(s)	Study type	Reference
Single or multiple organisms, given as capsules, powder, or part of a drink or infant formula milk	28 trials RT ; n=6705 participants	Maternal-infant intervention	Allergy prevention	↓Risk of eczema and/or atopic eczema at age ≤4 y ↓Allergic sensitization to cow milk at age 1–2 y ↓Risk of atopic eczema ↓Risk of food hypersensitivity When probiotics were administered either only prenatally or only postnatally, no effects on atopy and food hypersensitivity	Systematic review and meta-analysis	(94)
Combinations of lactobacilli and bifidobacteria	17 trials ; n=2947 infants	Maternal-infant intervention	Allergy prevention	↓Risk of atopic eczema ↓Risk of food hypersensitivity When probiotics were administered either only prenatally or only postnatally, no effects on atopy and food hypersensitivity	Systematic review and meta-analysis	(173)
Combinations of lactobacilli and bifidobacteria	17 RT ; n=4755 children [Probiotic n = 2381 ; Control n = 2374]	Maternal intervention during pregnancy	Allergy prevention	↓Risk ratio for eczema No impact on asthma, wheezing, or rhinoconjunctivitis	Meta-analysis	(174)

¹AAF, amino acid-based formula; CMA, cow milk allergy; DSM, German Collection of Microorganisms and Cell Cultures; RT, randomized trial.

histone deacetylases (114) and can induce the expression of noncoding RNAs (113, 116). Furthermore, a lower butyrate production and shifted gut microbiota composition toward an enrichment of *Bacteroides* and *Alistipes* genera have been reported in infants with non-IgE-mediated CMA (122). Low concentrations of SCFAs at 1 y of age have been associated with questionnaire-reported symptoms of FA at 4 y (123). In addition, propionate has been associated with increased expression of FOXP3 and IL10 in colonic Treg cells (119). There are signals of an association between SCFA and Treg cell development and function by epigenetic mechanisms, but the influence of this association in the risk of FA is still not clear.

Other microbial metabolites

It has been suggested that some other microbial metabolites such as staphylococcal enterotoxin B could act as adjuvants of food allergens during simultaneous exposure via skin (74). *Staphylococcus aureus* colonization of the skin has been associated with FA to peanut, egg white, and cow milk in patients with atopic dermatitis, and would be associated with skin barrier dysfunction and immune system dysregulation (124). Bacterial LPSs are strong immunostimulants that can induce tolerance at certain doses (125). Their role in allergy seems to be conditioned by the timing of exposure, the presence of pre-existing disease, and polymorphisms in the genes that encode endotoxin receptors (126). Evidence in humans is unclear but results from animal studies indicate LPS might prevent adverse IgE-mediated reactions by regulation of type 2 helper T-cell responses (127) and suppression of mast cell responses (128).

There is substantial evidence that intestinal bacteria can produce significant amounts of folate as well as other B vitamins complementing the dietary intake (129), including generally recognized beneficial microorganisms such as bifidobacteria and lactic acid bacteria (115). These B vitamins, and particularly folate, play a crucial role in epigenetic regulation as donors of methyl groups for DNA, RNA, and protein methylation (130, 131). Folate-induced changes in DNA methylation can modify gene expression in helper T cells (132), which has been proposed as a plausible mechanism underlying associations between folate and several diseases such as asthma (129), child wheeze (133), and allergy (134). For FA, it is still largely underexplored with contradictory results depending on the studies (132). Most of the few studies conducted to date suggest that maternal folate exposure is not associated with the development of FA (132). However, a retrospective study suggested that maternal folic acid supplementation in dosages higher than recommended might be a risk factor for allergy development (135).

Emerging evidence on the role of biotinylation upon immune function (136–138) and microbial metabolites such as polyamines (139–143) indicate potential further links between the gut microbiome and allergy by epigenetic regulation of genes modulating the activity of T and B lymphocytes, and proinflammatory cytokine expression (111, 136–144).

Recommendation/Guidance for Future Research

FA research is now experiencing an exciting new era thanks to advances on immunological, microbiological, and epigenetic factors and their integration, increasing knowledge of risk factors and potential biomarkers. However, limited data are available to identify potential biomarker or biomarker combinations determining a risk reduction in FA. The EAACI has recently published a systematic review as a source of evidence to support the development of FA prevention guidelines (39). This systematic review included 46 intervention studies to reduce the risk of FA in infancy (≤ 1 y) or early childhood. Different interventions during pregnancy, lactation, and infancy, including dietary avoidance of food allergens, vitamin supplements, fish oil, probiotics, prebiotics, synbiotics, and emollients, were included. Results showed that interventions have little or no effect in preventing FA, but the evidence is very uncertain. The systematic review concluded that most of the evidence has been published in the last 10 y, and still no clear data are available on preventing FA. There is a need to validate the potential benefits of early introduction of food allergens in a wider range of populations. Furthermore, there is a lack of studies analyzing serial and longitudinal biomarkers from birth up to adulthood, and clear biomarkers have not been identified until now. Promising potential biomarkers associated with FA, such as the depletion of key microbial components (e.g., *Bifidobacterium* and *Bacteroides* genus) or methylation profiles in the *FOXP3* and *IL10* genes, should be deeply evaluated in future studies.

To bridge the gap, more data are required on the maternal impact during gestation on fetal immune regulation as well as the immunometabolic profile of breast milk composition (immune cells, cytokines, hormones). There are also a limited number of studies focusing on immunology, microbiome, and diet, but few assess across the board. More cohort and intervention studies are needed to confirm which methylation profiles are suitable as biomarkers to monitor risk reduction of FA. Thus, designing nutritional intervention trials aimed at risk reduction of FA, or induction of tolerance, could need stratification based on specific risk factors to determine a design that is still feasible to execute. Indeed, the low incidence of oral food challenge–confirmed FA in the general population requires high numbers of infants to be able to detect a significant effect of an intervention. This review of currently available and emerging biomarkers linked to allergy can inform the design of future intervention studies. The available literature suggests that a highly collaborative approach spanning nutritional, genetic, and microbial biomarkers will be valuable in identifying panels of biomarkers that best predict FA, its severity, or its remission.

Acknowledgments

We thank Professor Philip Calder (University of Southampton, United Kingdom), Dr Jalil Benyacoub (Nestlé Health Science, Switzerland), Dr Stein-Erik Birkeland (Tine SA

R&D, Norway), Dr Bruno Pot (Yakult Europe BV, The Netherlands), Dr Patrizia Bohnhorst (Procter & Gamble, Germany), Dr Elizabeth Forbes-Blom (Nestlé Research Center, Switzerland), and Professor Ascensión Marcos (ICTAN-CSIC, Spain) for their contributions to the discussions that form the basis of this article and their help in the reviewing phase. Dr Siméon Bourdoux and Mr Adam Coventry (ILSI Europe, Belgium) coordinated the work of authors and facilitated meetings and discussions.

The authors' responsibilities were as follows—CEC, DM, LU, CG-G, MCC: wrote and revised the manuscript; LL, TR, SS, MT: assisted the previously cited authors in designing the structure of the manuscript and in reviewing the final content; and all authors: read and approved the final manuscript.

References

1. Wasserman S, Bégin P, Watson W. IgE-mediated food allergy. *Allergy Asthma Clin Immunol* 2018;14:55.
2. DunnGalvin A, Blumchen K, Timmermans F, Regent L, Schnadt S, Podestà M, Sánchez Á, Couratier P, Feeney M, Hjørth B, et al. APPEAL-1: a multiple country European survey assessing the psychosocial impact of peanut allergy. *Allergy* 2020;75(11):2899–908.
3. Bilaver LA, Chadha AS, Doshi P, O'Dwyer L, Gupta RS. Economic burden of food allergy: a systematic review. *Ann Allergy Asthma Immunol* 2019;122(4):373–380.e1.
4. Nwaru BI, Hickstein L, Panesar SS, Muraro A, Werfel T, Cardona V, Dubois AE, Halken S, Hoffmann-Sommergruber K, Poulsen LK, et al. The epidemiology of food allergy in Europe: a systematic review and meta-analysis. *Allergy* 2014;69(1):62–75.
5. Biomarkers Definitions Working Group. Biomarkers and surrogate endpoints: preferred definitions and conceptual framework. *Clin Pharmacol Ther* 2001;69(3):89–95.
6. European Commission. Regulation (EC) no 1924/2006 of the European Parliament and of the Council of 20 December 2006 on nutrition and health claims made on foods. *Official Journal of the European Union* 2006;404:9–25.
7. Anvari S, Miller J, Yeh CY, Davis CM. IgE-mediated food allergy. *Clin Rev Allergy Immunol* 2019;57(2):244–60.
8. Berin MC, Sampson HA. Food allergy: an enigmatic epidemic. *Trends Immunol* 2013;34(8):390–7.
9. Gelincik A, Büyükköztürk S, Gül H, Işık E, Işsever H, Özşeker F, Çolakoglu B, Dal M, Ayvaz Ö, Güngör G, et al. Confirmed prevalence of food allergy and non-allergic food hypersensitivity in a Mediterranean population. *Clin Exp Allergy* 2008;38(8):1333–41.
10. Pyrhönen K, Hiltunen L, Kaila M, Nayha S, Laara E. Heredity of food allergies in an unselected child population: an epidemiological survey from Finland. *Pediatr Allergy Immunol* 2011;22(1pt2):e124–32.
11. Sasaki M, Peters RL, Koplin JJ, Field MJ, McWilliam V, Sawyer SM, Vuillermin PJ, Pezic A, Gurrin LC, Douglass JA, et al. Risk factors for food allergy in early adolescence: the schoolnuts study. *J Allergy Clin Immunol Pract* 2018;6(2):496–505.
12. Venter C, Hasan Arshad S, Grundy J, Pereira B, Bernie Clayton C, Voigt K, Higgins B, Dean T. Time trends in the prevalence of peanut allergy: three cohorts of children from the same geographical location in the UK. *Allergy* 2010;65(1):103–8.
13. Venter C, Patil V, Grundy J, Glasbey G, Twiselton R, Arshad SH, Dean T. Prevalence and cumulative incidence of food hyper-sensitivity in the first 10 years of life. *Pediatr Allergy Immunol* 2016;27(5):452–8.
14. Venter C, Pereira B, Voigt K, Grundy J, Clayton CB, Higgins B, Arshad SH, Dean T. Prevalence and cumulative incidence of food hypersensitivity in the first 3 years of life. *Allergy* 2008;63(3):354–9.
15. Bonnelykke K, Matheson MC, Pers TH, Granell R, Strachan DP, Alves AC, Linneberg A, Curtin JA, Warrington NM, Standl M, et al.

- Meta-analysis of genome-wide association studies identifies ten loci influencing allergic sensitization. *Nat Genet* 2013;45(8):902–6.
16. Hong X, Hao K, Ladd-Acosta C, Hansen KD, Tsai H-J, Liu X, Xu X, Thornton TA, Caruso D, Keet CA, et al. Genome-wide association study identifies peanut allergy-specific loci and evidence of epigenetic mediation in US children. *Nat Commun* 2015;6(1):1–12.
 17. Nicolaou N, Poorafshar M, Murray C, Simpson A, Winell H, Kerry G, Harlin A, Woodcock A, Ahlstedt S, Custovic A. Allergy or tolerance in children sensitized to peanut: prevalence and differentiation using component-resolved diagnostics. *J Allergy Clin Immunol* 2010;125(1):191–7.e13.
 18. Pyrhönen K, Nayha S, Kaila M, Hiltunen L, Laara E. Occurrence of parent-reported food hypersensitivities and food allergies among children aged 1–4 yr. *Pediatr Allergy Immunol* 2009;20(4):328–38.
 19. Marenholz I, Grosche S, Kalb B, Ruschendorf F, Blumchen K, Schlags R, Harandi N, Price M, Hansen G, Seidenberg J, et al. Genome-wide association study identifies the SERPINB gene cluster as a susceptibility locus for food allergy. *Nat Commun* 2017;8(1):1056.
 20. Kull I, Melen E, Alm J, Hallberg J, Svartengren M, van Hage M, Pershagen G, Wickman M, Bergstrom A. Breast-feeding in relation to asthma, lung function, and sensitization in young schoolchildren. *J Allergy Clin Immunol* 2010;125(5):1013–19.
 21. Saarinen UM, Kajosaari M. Breastfeeding as prophylaxis against atopic disease: prospective follow-up study until 17 years old. *Lancet* 1995;346(8982):1065–9.
 22. Chalmers JR, Haines RH, Bradshaw LE, Montgomery AA, Thomas KS, Brown SJ, Ridd MJ, Lawton S, Simpson EL, Cork MJ, et al. Daily emollient during infancy for prevention of eczema: the BEEP randomised controlled trial. *Lancet* 2020;395(10228):962–72.
 23. Rice NE, Patel BD, Lang IA, Kumari M, Frayling TM, Murray A, Melzer D. Filaggrin gene mutations are associated with asthma and eczema in later life. *J Allergy Clin Immunol* 2008;122(4):834–6.
 24. Alam R, Abdolmaleky HM, Zhou JR. Microbiome, inflammation, epigenetic alterations, and mental diseases. *Am J Med Genet B Neuropsychiatr Genet* 2017;174(6):651–60.
 25. Bowers EC, McCullough SD. Linking the epigenome with exposure effects and susceptibility: the epigenetic seed and soil model. *Toxicol Sci* 2017;155(2):302–14.
 26. Freeland DMH, Manohar M, Andorf S, Hobson BD, Zhang W, Nadeau KC. Oral immunotherapy for food allergy. *Semin Immunol* 2017;30:36–44.
 27. Lin PI, Shu H, Mersha TB. Comparing DNA methylation profiles across different tissues associated with the diagnosis of pediatric asthma. *Sci Rep* 2020;10(1):151.
 28. Thibeault AAH, Laprise C. Cell-specific DNA methylation signatures in asthma. *Genes (Basel)* 2019;10(11):932.
 29. Victora CG, Bahl R, Barros AJD, França GVA, Horton S, Krasevec J, Murch S, Sankar MJ, Walker N, Rollins NC, et al. Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect. *Lancet* 2016;387(10017):475–90.
 30. Ip S, Chung M, Raman G, Chew P, Magula N, DeVine D, Trikalinos T, Lau J. Breastfeeding and maternal and infant health outcomes in developed countries. *Evid Rep Technol Assess (Full Rep)* 2007;153(153):1–186.
 31. Lodge C, Tan D, Lau M, Dai X, Tham R, Lowe A, Bowatte G, Allen K, Dharmage S. Breastfeeding and asthma and allergies: a systematic review and meta-analysis. *Acta Paediatr* 2015;104:38–53.
 32. van den Elsen LWJ, Garssen J, Burcelin R, Verhasselt V. Shaping the gut microbiota by breastfeeding: the gateway to allergy prevention? *Front Pediatr* 2019;7:47.
 33. Van Odijk J, Kull I, Borres MP, Brandtzaeg P, Edberg U, Hanson LÅ, Høst A, Kuitunen M, Olsen SF, Skerfving S, et al. Breastfeeding and allergic disease: a multidisciplinary review of the literature (1966–2001) on the mode of early feeding in infancy and its impact on later atopic manifestations. *Allergy* 2003;58(9):833–43.
 34. Lucas A, Brooke OG, Morley R, Cole TJ, Bamford MF. Early diet of preterm infants and development of allergic or atopic disease: randomised prospective study. *BMJ* 1990;300(6728):837–40.
 35. Mihrshahi S, Ampon R, Webb K, Almqvist C, Kemp AS, Hector D, Marks GB. The association between infant feeding practices and subsequent atopy among children with a family history of asthma. *Clin Exp Allergy* 2007;37(5):671–9.
 36. Pesonen M, Kallio MJT, Ranki A, Siimes MA. Prolonged exclusive breastfeeding is associated with increased atopic dermatitis: a prospective follow-up study of unselected healthy newborns from birth to age 20 years. *Clin Exp Allergy* 2006;36(8):1011–18.
 37. Munblit D, Perkin MR, Palmer DJ, Allen KJ, Boyle RJ. Assessment of evidence about common infant symptoms and cow's milk allergy. *JAMA Pediatr* 2020;174(6):599–608.
 38. Boix-Amorós A, Collado MC, Van't Land B, Calvert A, Le Doare K, Garssen J, Hanna H, Khaleva E, Peroni DG, Geddes DT, et al. Reviewing the evidence on breast milk composition and immunological outcomes. *Nutr Rev* 2019;77(8):541–56.
 39. de Silva D, Halken S, Singh C, Muraro A, Angier E, Arasi S, Arshad H, Beyer K, Boyle R, du Toit G, et al. Preventing food allergy in infancy and childhood: systematic review of randomised controlled trials. *Pediatr Allergy Immunol* 2020;31(7):813–26.
 40. Munblit D, Boyle RJ. Modulating breast milk composition—the key to allergy prevention? *Int Arch Allergy Immunol* 2012;159(2):107–8.
 41. Munblit D, Peroni DG, Boix-Amorós A, Hsu PS, Van't Land B, Gay MCL, Kolotilina A, Skevaki C, Boyle RJ, Collado MC, et al. Human milk and allergic diseases: an unsolved puzzle. *Nutrients* 2017;9(8):894.
 42. Munblit D, Boyle RJ, Warner JO. Factors affecting breast milk composition and potential consequences for development of the allergic phenotype. *Clin Exp Allergy* 2015;45(3):583–601.
 43. Munblit D, Treneva M, Peroni DG, Colicino S, Chow LY, Dissanayake S, Abrol P, Sheth S, Pampura A, Boner AL, et al. Colostrum and mature human milk of women from London, Moscow, and Verona: determinants of immune composition. *Nutrients* 2016;8(11):695.
 44. Ruiz L, Espinosa-Martos I, García-Carral C, Manzano S, McGuire MK, Meehan CL, McGuire MA, Williams JE, Foster J, Sellen DW, et al. What's normal? Immune profiling of human milk from healthy women living in different geographical and socioeconomic settings. *Front Immunol* 2017;8:696.
 45. Tomićić S, Johansson G, Voor T, Björkstén B, Böttcher MF, Jenmalm MC. Breast milk cytokine and IgA composition differ in Estonian and Swedish mothers—relationship to microbial pressure and infant allergy. *Pediatr Res* 2010;68(4):330–4.
 46. Picariello G, De Cicco M, Nocerino R, Paparo L, Mamone G, Addeo F, Berni Canani R. Excretion of dietary cow's milk derived peptides into breast milk. *Front Nutr* 2019;6:25.
 47. Gouw JW, Jo J, Meulenbroek L, Heijer TS, Kremer E, Sandalova E, Knulst AC, Jeurink PV, Garssen J, Rijnierse A, et al. Identification of peptides with tolerogenic potential in a hydrolysed whey-based infant formula. *Clin Exp Allergy* 2018;48(10):1345–53.
 48. Boyle RJ, Ierodiakonou D, Khan T, Chivinge J, Robinson Z, Geoghegan N, Jarrold K, Afxentiou T, Reeves T, Cunha S, et al. Hydrolysed formula and risk of allergic or autoimmune disease: systematic review and meta-analysis. *BMJ* 2016;352:i974.
 49. Osborn DA, Sinn JKH, Jones LJ. Infant formulas containing hydrolysed protein for prevention of allergic disease. *Cochrane Database Syst Rev* 2018;10(10):CD003664.
 50. Knol EF, de Jong NW, Ulfman LH, Tiemessen MM. Management of cow's milk allergy from an immunological perspective: what are the options? *Nutrients* 2019;11(11):2734.
 51. Fleischer DM, Chan ES, Venter C, Spergel JM, Abrams EM, Stukus D, Groetch M, Shaker M, Greenhawt M. A consensus approach to the primary prevention of food allergy through nutrition: guidance from the American Academy of Allergy, Asthma, and Immunology; American College of Allergy, Asthma, and Immunology; and the

- Canadian Society for Allergy and Clinical Immunology. *J Allergy Clin Immunol Pract* 2021;9(1):22–43.e4.
52. Greer FR, Sicherer SH, Burks AW. The effects of early nutritional interventions on the development of atopic disease in infants and children: the role of maternal dietary restriction, breastfeeding, hydrolyzed formulas, and timing of introduction of allergenic complementary foods. *Pediatrics* 2019;143(4):e20190281.
 53. Urashima M, Mezawa H, Okuyama M, Urashima T, Hirano D, Gocho N, Tachimoto H. Primary prevention of cow's milk sensitization and food allergy by avoiding supplementation with cow's milk formula at birth: a randomized clinical trial. *JAMA Pediatr* 2019;173(12):1137–45.
 54. Du Toit G, Sampson HA, Plaut M, Burks AW, Akdis CA, Lack G. Food allergy: update on prevention and tolerance. *J Allergy Clin Immunol* 2018;141(1):30–40.
 55. Du Toit G, Roberts G, Sayre PH, Bahnson HT, Radulovic S, Santos AF, Brough HA, Phippard D, Basting M, Feeney M, et al. Randomized trial of peanut consumption in infants at risk for peanut allergy. *N Engl J Med* 2015;372(9):803–13.
 56. Perkin MR, Logan K, Tseng A, Raji B, Ayis S, Peacock J, Brough H, Marrs T, Radulovic S, Craven J, et al. Randomized trial of introduction of allergenic foods in breast-fed infants. *N Engl J Med* 2016;374(18):1733–43.
 57. Elbert NJ, Kiefte-de Jong JC, Voortman T, Nijsten TEC, De Jong NW, Jaddoe VVW, De Jongste JC, Van Wijk RG, Duijts L, Pasmans S. Allergenic food introduction and risk of childhood atopic diseases. *PLoS One* 2017;12(11):e0187999.
 58. Togias A, Cooper SF, Acebal ML, Assa'ad A, Baker JR, Beck LA, Block J, Byrd-Bredbenner C, Chan ES, Eichenfield LF, et al. Addendum guidelines for the prevention of peanut allergy in the United States: report of the National Institute of Allergy and Infectious Diseases–Sponsored Expert Panel. *J Allergy Clin Immunol* 2017;139(1):29–44.
 59. Roduit C, Frei R, Depner M, Schaub B, Loss G, Genuneit J, Pfeffler P, Hyvärinen A, Karvonen AM, Riedler J, et al. Increased food diversity in the first year of life is inversely associated with allergic diseases. *J Allergy Clin Immunol* 2014;133(4):1056–64.e7.
 60. Martorell Calatayud C, Muriel García A, Martorell Aragonés A, De La Hoz Caballer B. Safety and efficacy profile and immunological changes associated with oral immunotherapy for IgE-mediated cow's milk allergy in children: systematic review and meta-analysis. *J Investig Allergol Clin Immunol* 2014;24(5):298–307.
 61. Skripak JM, Nash SD, Rowley H, Brereton NH, Oh S, Hamilton RG, Matsui EC, Burks AW, Wood RA. A randomized, double-blind, placebo-controlled study of milk oral immunotherapy for cow's milk allergy. *J Allergy Clin Immunol* 2008;122(6):1154–60.
 62. Datema MR, Eller E, Zwinderman AH, Poulsen LK, Versteeg SA, van Ree R, Bindslev-Jensen C. Ratios of specific IgG4 over IgE antibodies do not improve prediction of peanut allergy nor of its severity compared to specific IgE alone. *Clin Exp Allergy* 2019;49(2):216–26.
 63. Sim K, Mijakoski D, Stoleski S, del Rio PR, Sammut P, Le TM, Munblit D, Boyle RJ. Outcomes for clinical trials of food allergy treatments. *Ann Allergy Asthma Immunol* 2020;125(5):535–42.
 64. Schoos AMM, Bullens D, Chawes BL, Costa J, De Vlioger L, DunnGalvin A, Epstein MM, Garssen J, Hilger C, Knipping K, et al. Immunological outcomes of allergen-specific immunotherapy in food allergy. *Front Immunol* 2020;11:568598.
 65. Goldberg MR, Mor H, Magid Neriya D, Magzal F, Muller E, Appel MY, Nachshon L, Borenstein E, Tamir S, Louzoun Y, et al. Microbial signature in IgE-mediated food allergies. *Genome Med* 2020;12(1):92.
 66. Vuillermin PJ, O'Hely M, Collier F, Allen KJ, Tang MLK, Harrison LC, Carlin JB, Saffery R, Ranganathan S, Sly PD, et al. Maternal carriage of *Prevotella* during pregnancy associates with protection against food allergy in the offspring. *Nat Commun* 2020;11(1):1452.
 67. Zhao W, He Ho, Bunyavanich S. The gut microbiome in food allergy. *Ann Allergy Asthma Immunol* 2019;122(3):276–82.
 68. Azad MB, Konya T, Guttman DS, Field CJ, Sears MR, HayGlass KT, Mandhane PJ, Turvey SE, Subbarao P, Becker AB, et al. Infant gut microbiota and food sensitization: associations in the first year of life. *Clin Exp Allergy* 2015;45(3):632–43.
 69. Kozyrskyj AL. Can we predict future allergies from our infant gut microbiota? *Expert Rev Respir Med* 2015;9(6):667–70.
 70. Arrieta M-C, Stiemsma LT, Dimitriu PA, Thorson L, Russell S, Yurist-Doutsch S, Kuzeljevic B, Gold MJ, Britton HM, Lefebvre DL, et al. Early infancy microbial and metabolic alterations affect risk of childhood asthma. *Sci Transl Med* 2015;7(307):307ra152.
 71. Savage JH, Lee-Sarwar KA, Sordillo J, Bunyavanich S, Zhou Y, O'Connor G, Sandel M, Bacharier LB, Zeiger R, Sodergren E, et al. A prospective microbiome-wide association study of food sensitization and food allergy in early childhood. *Allergy* 2018;73(1):145–52.
 72. Bunyavanich S, Shen N, Grishin A, Wood R, Burks W, Dawson P, Jones SM, Leung DYM, Sampson H, Sicherer S, et al. Early-life gut microbiome composition and milk allergy resolution. *J Allergy Clin Immunol* 2016;138(4):1122–30.
 73. Ling Z, Li Z, Liu X, Cheng Y, Luo Y, Tong X, Yuan L, Wang Y, Sun J, Li L, et al. Altered fecal microbiota composition associated with food allergy in infants. *Appl Environ Microbiol* 2014;80(8):2546–54.
 74. Benedé S, Blázquez AB, Chiang D, Tordesillas L, Berin MC. The rise of food allergy: environmental factors and emerging treatments. *EBioMedicine* 2016;7:27–34.
 75. Hua X, Goedert JJ, Pu A, Yu G, Shi J. Allergy associations with the adult fecal microbiota. *EBioMedicine* 2016;3:172–9.
 76. Renz H, Skevaki C. Early life microbial exposures and allergy risks: opportunities for prevention. *Nat Rev Immunol* 2021;21(3):177–91.
 77. Dominguez-Bello MG, De Jesus-Laboy KM, Shen N, Cox LM, Amir A, Gonzalez A, Bokulich NA, Song SJ, Hoashi M, Rivera-Vinas JJ, et al. Partial restoration of the microbiota of cesarean-born infants via vaginal microbial transfer. *Nat Med* 2016;22(3):250–3.
 78. Shu SA, Yuen AWT, Woo E, Chu KH, Kwan HS, Yang GX, Yang Y, Leung PSC. Microbiota and food allergy. *Clin Rev Allergy Immunol* 2019;57(1):83–97.
 79. Papathoma E, Triga M, Fouzas S, Dimitriou G. Cesarean section delivery and development of food allergy and atopic dermatitis in early childhood. *Pediatr Allergy Immunol* 2016;27(4):419–24.
 80. Pyrhönen K, Nayha S, Hiltunen L, Laara E. Caesarean section and allergic manifestations: insufficient evidence of association found in population-based study of children aged 1 to 4 years. *Acta Paediatr* 2013;102(10):982–9.
 81. Eggesbø M, Botten G, Stigum H, Nafstad P, Magnus P. Is delivery by cesarean section a risk factor for food allergy? *J Allergy Clin Immunol* 2003;112(2):420–6.
 82. Bager P, Wohlfahrt J, Westergaard T. Cesarean delivery and risk of atopy and allergic disease: meta-analyses. *Clin Exp Allergy* 2008;38(4):634–42.
 83. Love BL, Mann JR, Hardin JW, Lu ZK, Cox C, Amrol DJ. Antibiotic prescription and food allergy in young children. *Allergy Asthma Clin Immunol* 2016;12(1):41.
 84. Huang YJ, Marsland BJ, Bunyavanich S, O'Mahony L, Leung DYM, Muraro A, Fleisher TA. The microbiome in allergic disease: current understanding and future opportunities—2017 PRACTALL document of the American Academy of Allergy, Asthma & Immunology and the European Academy of Allergy and Clinical Immunology. *J Allergy Clin Immunol* 2017;139(4):1099–110.
 85. Kim S, Covington A, Pamer EG. The intestinal microbiota: antibiotics, colonization resistance, and enteric pathogens. *Immunol Rev* 2017;279(1):90–105.
 86. Canani RB, Sangwan N, Stefa AT, Nocerino R, Paparo L, Aitoro R, Calignano A, Khan AA, Gilbert JA, Nagler CR. Lactobacillus rhamnosus GG-supplemented formula expands butyrate-producing bacterial strains in food allergic infants. *ISME J* 2016;10(3):742–50.
 87. Bisgaard H, Li N, Bonnelykke K, Chawes BLK, Skov T, Paludan-Müller G, Stokholm J, Smith B, Krogfelt KA. Reduced diversity of the intestinal microbiota during infancy is associated with increased risk of allergic disease at school age. *J Allergy Clin Immunol* 2011;128(3):646–52.e5.

88. Stewart CJ, Ajami NJ, O'Brien JL, Hutchinson DS, Smith DP, Wong MC, Ross MC, Lloyd RE, Doddapaneni H, Metcalf GA, et al. Temporal development of the gut microbiome in early childhood from the TEDDY study. *Nature* 2018;562(7728):583–8.
89. Bezirtzoglou E, Tsiotsias A, Welling GW. Microbiota profile in feces of breast-and formula-fed newborns by using fluorescence in situ hybridization (FISH). *Anaerobe* 2011;17(6):478–82.
90. Dzidic M, Mira A, Artacho A, Abrahamsson TR, Jenmalm MC, Collado MC. Allergy development is associated with consumption of breastmilk with a reduced microbial richness in the first month of life. *Pediatr Allergy Immunol* 2020;31(3):250–7.
91. Järvinen KM, Martin H, Oyoshi MK. Immunomodulatory effects of breast milk on food allergy. *Ann Allergy Asthma Immunol* 2019;123(2):133–43.
92. Kotz D, Simpson CR, Sheikh A. Incidence, prevalence, and trends of general practitioner-recorded diagnosis of peanut allergy in England, 2001 to 2005. *J Allergy Clin Immunol* 2011;127(3):623–30.e1.
93. Ege MJ, Mayer M, Normand AC, Genuneit J, Cookson W, Braun-Fahrlander C, Heederik D, Piarroux R, Von Mutius E. Exposure to environmental microorganisms and childhood asthma. *N Engl J Med* 2011;364(8):701–9.
94. Garcia-Larsen V, Ierodiakonou D, Jarrold K, Cunha S, Chivinge J, Robinson Z, Geoghegan N, Ruparelia A, Devani P, Trivella M, et al. Diet during pregnancy and infancy and risk of allergic or autoimmune disease: a systematic review and meta-analysis. *PLoS Med* 2018;15(2):e1002507.
95. Hoppenbrouwers T, Cvejić Hogervorst JH, Garssen J, Wichers HJ, Willemsen LEM. Long chain polyunsaturated fatty acids (LCPUFAs) in the prevention of food allergy. *Front Immunol* 2019;10:1118.
96. Aryan Z, Rezaei N, Camargo CA. Vitamin D status, aeroallergen sensitization, and allergic rhinitis: a systematic review and meta-analysis. *Int Rev Immunol* 2017;36(1):41–53.
97. Koplin JJ, Suaini NHA, Vuillermin P, Ellis JA, Panjari M, Ponsonby AL, Peters RL, Matheson MC, Martino D, Dang T, et al. Polymorphisms affecting vitamin D-binding protein modify the relationship between serum vitamin d (25[OH]D3) and food allergy. *J Allergy Clin Immunol* 2016;137(2):500–6.e4.
98. Goldring ST, Griffiths CJ, Martineau AR, Robinson S, Yu C, Poulton S, Kirkby JC, Stocks J, Hooper R, Shaheen SO, et al. Prenatal vitamin D supplementation and child respiratory health: a randomised controlled trial. *PLoS One* 2013;8(6):e66627.
99. Allen KJ, Koplin JJ, Ponsonby AL, Gurrin LC, Wake M, Vuillermin P, Martin P, Matheson M, Lowe A, Robinson M, et al. Vitamin D insufficiency is associated with challenge-proven food allergy in infants. *J Allergy Clin Immunol* 2013;131(4):1109–16.e6.
100. Yepes-Nuñez JJ, Brożek JL, Fiocchi A, Pawankar R, Cuello-García C, Zhang Y, Morgano GP, Agarwal A, Gandhi S, Terracciano L, et al. Vitamin D supplementation in primary allergy prevention: systematic review of randomized and non-randomized studies. *Allergy* 2018;73(1):37–49.
101. Heine RG. Food allergy prevention and treatment by targeted nutrition. *Ann Nutr Metab* 2018;72(Suppl. 3):33–45.
102. Canani RB, Paparo L, Nocerino R, Scala CD, Gatta GD, Maddalena Y, Buono A, Bruno C, Voto L, Ercolini D. Gut microbiome as target for innovative strategies against food allergy. *Front Immunol* 2019;10:191.
103. Paparo L, Nocerino R, Di Scala C, Della Gatta G, Di Costanzo M, Buono A, Bruno C, Berni Canani R. Targeting food allergy with probiotics. *Adv Exp Med Biol* 2019;1125:57–68.
104. Turner PJ, Campbell DE, Boyle RJ, Levin ME. Primary prevention of food allergy: translating evidence from clinical trials to population-based recommendations. *J Allergy Clin Immunol Pract* 2018;6(2):367–75.
105. de Silva D, Geromi M, Halken S, Host A, Panesar SS, Muraro A, Werfel T, Hoffmann-Sommergruber K, Roberts G, Cardona V, et al. Primary prevention of food allergy in children and adults: systematic review. *Allergy* 2014;69(5):581–9.
106. Fiocchi A, Pawankar R, Cuello-Garcia C, Ahn K, Al-Hammadi S, Agarwal A, Beyer K, Burks W, Canonica GW, Ebisawa M, et al. World Allergy Organization-McMaster University guidelines for allergic disease prevention (GLAD-P): probiotics. *World Allergy Organ J* 2015;8(1):4.
107. Arslanoglu S, Moro GE, Schmitt J, Tandoi L, Rizzardi S, Boehm G. Early dietary intervention with a mixture of prebiotic oligosaccharides reduces the incidence of allergic manifestations and infections during the first two years of life. *J Nutr* 2008;138(6):1091–5.
108. Moro G, Arslanoglu S, Stahl B, Jelinek J, Wahn U, Boehm G. A mixture of prebiotic oligosaccharides reduces the incidence of atopic dermatitis during the first six months of age. *Arch Dis Child* 2006;91(10):814–19.
109. Wopereis H, Van Ampting MTJ, Cetinyurek-Yavuz A, Slump R, Candy DCA, Butt AM, Peroni DG, Vandenplas Y, Fox AT, Shah N, et al. A specific synbiotic-containing amino acid-based formula restores gut microbiota in non-IgE mediated cow's milk allergic infants: a randomized controlled trial. *Clin Transl Allergy* 2019;9(1):27.
110. Levy M, Thaiss CA, Elinav E. Metabolites: messengers between the microbiota and the immune system. *Genes Dev* 2016;30(14):1589–97.
111. Hullar MAJ, Fu BC. Diet, the gut microbiome, and epigenetics. *Cancer J* 2014;20(3):170–5.
112. Brown K, DeCoffe D, Molcan E, Gibson DL. Diet-induced dysbiosis of the intestinal microbiota and the effects on immunity and disease. *Nutrients* 2012;4(8):1095–119.
113. Luo A, Leach ST, Barres R, Hesson LB, Grimm MC, Simar D. The microbiota and epigenetic regulation of T helper 17/regulatory T cells: in search of a balanced immune system. *Front Immunol* 2017;8:417.
114. Bourassa MW, Alim I, Bultman SJ, Ratan RR. Butyrate, neuroepigenetics and the gut microbiome: can a high fiber diet improve brain health? *Neurosci Lett* 2016;625:56–63.
115. Qin Y, Wade PA. Crosstalk between the microbiome and epigenome: messages from bugs. *J Biochem* 2018;163(2):105–12.
116. Tzur G, Levy A, Meiri E, Barad O, Spector Y, Bentwich Z, Mizrahi L, Katzenellenbogen M, Ben-Shushan E, Reubinoff BE, et al. MicroRNA expression patterns and function in endodermal differentiation of human embryonic stem cells. *PLoS One* 2008;3(11):e3726.
117. Arpaia N, Campbell C, Fan X, Dikiy S, Van Der Veeke J, Deroos P, Liu H, Cross JR, Pfeffer K, Coffey PJ, et al. Metabolites produced by commensal bacteria promote peripheral regulatory T-cell generation. *Nature* 2013;504(7480):451–5.
118. Kim CH, Park J, Kim M. Gut microbiota-derived short-chain fatty acids, T cells, and inflammation. *Immune Network* 2014;14(6):277.
119. Smith PM, Howitt MR, Panikov N, Michaud M, Gallini CA, Bohlooly-Y M, Glickman JN, Garrett WS. The microbial metabolites, short-chain fatty acids, regulate colonic T reg cell homeostasis. *Science* 2013;341(6145):569–73.
120. Baxter NT, Schmidt AW, Venkataraman A, Kim KS, Waldron C, Schmidt TM. Dynamics of human gut microbiota and short-chain fatty acids in response to dietary interventions with three fermentable fibers. *mBio* 2019;10(1):02566–18.
121. Morrison DJ, Preston T. Formation of short chain fatty acids by the gut microbiota and their impact on human metabolism. *Gut Microbes* 2016;7(3):189–200.
122. Berni Canani R, De Filippis F, Nocerino R, Paparo L, Di Scala C, Cosenza L, Della Gatta G, Calignano A, De Caro C, Laiola M, et al. Gut microbiota composition and butyrate production in children affected by non-IgE-mediated cow's milk allergy. *Sci Rep* 2018;8(1):12500.
123. Sandin A, Bråbäck L, Norin E, Björkstén B. Faecal short chain fatty acid pattern and allergy in early childhood. *Acta Paediatr* 2009;98(5):823–7.
124. Jones AL, Curran-Everett D, Leung DYM. Food allergy is associated with *Staphylococcus aureus* colonization in children with atopic dermatitis. *J Allergy Clin Immunol* 2016;137(4):1247–8.e3.

125. Wassenaar TM, Zimmermann K. Lipopolysaccharides in food, food supplements, and probiotics: should we be worried? *Eur J Microbiol Immunol* 2018;8(3):63–9.
126. Williams LK, Ownby DR, Maliarik MJ, Johnson CC. The role of endotoxin and its receptors in allergic disease. *Ann Allergy Asthma Immunol* 2005;94(3):323–32.
127. Torii I, Shimizu S, Daimon T, Shinohara Y, Kudo T, Sato A, Tsujimura T. Exposure to high doses of lipopolysaccharide during ovalbumin sensitization prevents the development of allergic Th2 responses to a dietary antigen. *J Toxicol Pathol* 2014;27(3+4):205–15.
128. Wang N, McKell M, Dang A, Yamani A, Waggoner L, Vanoni S, Noah T, Wu D, Kordowski A, Köhl J, et al. Lipopolysaccharide suppresses IgE-mast cell-mediated reactions. *Clin Exp Allergy* 2017;47(12):1574–85.
129. Kok DE, Steegenga WT, McKay JA. Folate and epigenetics: why we should not forget bacterial biosynthesis. *Epigenomics* 2018;10(9):1147–50.
130. Crider KS, Yang TP, Berry RJ, Bailey LB. Folate and DNA methylation: a review of molecular mechanisms and the evidence for folate's role. *Adv Nutr* 2012;3(1):21–38.
131. McKay JA, Mathers JC. Diet induced epigenetic changes and their implications for health. *Acta Physiologica* 2011;202(2):103–18.
132. Brown SB, Reeves KW, Bertone-Johnson ER. Maternal folate exposure in pregnancy and childhood asthma and allergy: a systematic review. *Nutr Rev* 2014;72(1):55–64.
133. Roy A, Kocak M, Hartman TJ, Vereen S, Adgent M, Piyathilake C, Tylavsky FA, Carroll KN. Association of prenatal folate status with early childhood wheeze and atopic dermatitis. *Pediatr Allergy Immunol* 2018;29(2):144–50.
134. Socha-Banasiak A, Kamer B, Gach A, Wysocka U, Jakubowski L, Głowacka E, Czkwianianc E. Folate status, regulatory t cells and MTHFR C677T polymorphism study in allergic children. *Adv Med Sci* 2016;61(2):300–5.
135. Socha-Banasiak A, Kamer B, Pacześ K, Ślusarek B, Pawlikowski B, Czkwianianc E. Trends in folic acid supplementation during pregnancy – the effect on allergy development in children. *Adv Dermatol Allergol* 2018;35(2):139–44.
136. Bhat MI, Kapila R. Dietary metabolites derived from gut microbiota: critical modulators of epigenetic changes in mammals. *Nutr Rev* 2017;75(5):374–89.
137. Elahi A, Sabui S, Narasappa NN, Agrawal S, Lambrecht NW, Agrawal A, Said HM. Biotin deficiency induces Th1- and Th17-mediated proinflammatory responses in human CD4+ T lymphocytes via activation of the mTOR signaling pathway. *J Immunol* 2018;200(8):2563–70.
138. Xue J, Zempleni J. Epigenetic synergies between biotin and folate in the regulation of pro-inflammatory cytokines and repeats. *Scand J Immunol* 2013;78(5):419–25.
139. Hesterberg RS, Cleveland JL, Epling-Burnette PK. Role of polyamines in immune cell functions. *Med Sci* 2018;6(1):22.
140. Kanemura A, Yoshikawa Y, Fukuda W, Tsumoto K, Kenmotsu T, Yoshikawa K. Opposite effect of polyamines on in vitro gene expression: enhancement at low concentrations but inhibition at high concentrations. *PLoS One* 2018;13(3):e0193595.
141. McCabe LR, Parameswaran N. Advances in probiotic regulation of bone and mineral metabolism. *Calcif Tissue Int* 2018;102(4):480–8.
142. Skrypnik K, Suliburska J. Association between the gut microbiota and mineral metabolism. *J Sci Food Agric* 2018;98(7):2449–60.
143. Timmons J, Chang ET, Wang J-Y, Rao JN. Polyamines and gut mucosal homeostasis. *J Gastrointest Dig Syst* 2012;2(Suppl 7):001.
144. Aleksandrova K, Romero-Mosquera B, Hernandez V. Diet, gut microbiome and epigenetics: emerging links with inflammatory bowel diseases and prospects for management and prevention. *Nutrients* 2017;9(9):1–13.
145. Prescott SL, Noakes P, Chow BWY, Breckler L, Thornton CA, Hollams EM, Ali M, van den Biggelaar AHJ, Tulic MK. Presymptomatic differences in Toll-like receptor function in infants who have allergy. *J Allergy Clin Immunol* 2008;122(2):391–9.e5.
146. Asai Y, Eslami A, van Ginkel CD, Akhbari L, Wan M, Ellis G, Ben-Shoshan M, Martino D, Ferreira MA, Allen K, et al. Genome-wide association study and meta-analysis in multiple populations identifies new loci for peanut allergy and establishes C11orf30/EMSY as a genetic risk factor for food allergy. *J Allergy Clin Immunol* 2018;141(3):991–1001.
147. Yavuz ST, Buyuktiryaki B, Sahiner UM, Birben E, Tuncer A, Yakarisik S, Karabulut E, Kalayci O, Sackesen C. Factors that predict the clinical reactivity and tolerance in children with cow's milk allergy. *Ann Allergy Asthma Immunol* 2013;110(4):284–9.
148. Hirota T, Nakayama T, Tamari M, Sato S, Yanagida N, Ebisawa M, Matsui T, Sugiura S, Ito K, Takaoka Y, et al. Association study of childhood food allergy with genome-wide association studies—discovered loci of atopic dermatitis and eosinophilic esophagitis. *J Allergy Clin Immunol* 2017;140(6):1713–16.
149. Dimitrov I, Doytchinova I. Associations between milk and egg allergens and the HLA-DRB1/DQ polymorphism: a bioinformatics approach. *Int Arch Allergy Immunol* 2016;169(1):33–9.
150. Martino D, Neeland M, Dang T, Cobb J, Ellis J, Barnett A, Tang M, Vuillermin P, Allen K, Saffery R. Epigenetic dysregulation of naive CD4+ T-cell activation genes in childhood food allergy. *Nat Commun* 2018;9(1):3308.
151. Venkataraman D, Soto-Ramirez N, Kurukulaaratchy RJ, Holloway JW, Karmaus W, Ewart SL, Arshad SH, Erlewyn-Lajeunesse M. Filaggrin loss-of-function mutations are associated with food allergy in childhood and adolescence. *J Allergy Clin Immunol* 2014;134(4):876–82.e4.
152. Ashley SE, Tan HTT, Peters R, Allen KJ, Vuillermin P, Dharmage SC, Tang MLK, Koplin J, Lowe A, Ponsonby AL, et al. Genetic variation at the Th2 immune gene IL13 is associated with IgE-mediated paediatric food allergy. *Clin Exp Allergy* 2017;47(8):1032–7.
153. Li J, Fung I, Glessner JT, Pandey R, Wei Z, Bakay M, Mentch FD, Pellegrino R, Wang T, Kim C, et al. Copy number variations in CTNNA3 and RFXO1 associate with pediatric food allergy. *J Immunol* 2015;195(4):1599–607.
154. Clark A, Mach N. Role of vitamin D in the hygiene hypothesis: the interplay between vitamin D, vitamin D receptors, gut microbiota, and immune response. *Front Immunol* 2016;7:627.
155. Buyuktiryaki B, Sahiner UM, Girgin G, Birben E, Soyer OU, Cavkaytar O, Cetin C, Arik Yilmaz E, Yavuz ST, Kalayci O, et al. Low indoleamine 2,3-dioxygenase activity in persistent food allergy in children. *Allergy* 2016;71(2):258–66.
156. Li X, Lee YJ, Jin F, Park YN, Deng Y, Kang Y, Yang JH, Chang JH, Kim DY, Kim JA, et al. Sirt1 negatively regulates FcεRI-mediated mast cell activation through AMPK- and PTP1B-dependent processes. *Sci Rep* 2017;7(1):6444.
157. Li X, Lee YJ, Jin F, Park YN, Deng Y, Kang Y, Yang JH, Chang JH, Kim DY, Kim JA, et al. Author correction: Sirt1 negatively regulates FcεRI-mediated mast cell activation through AMPK- and PTP1B-dependent processes. *Sci Rep* 2020;10(1):3552.
158. Martino D, Dang T, Sexton-Oates A, Prescott S, Tang MLK, Dharmage S, Gurrin L, Koplin J, Ponsonby AL, Allen KJ, et al. Blood DNA methylation biomarkers predict clinical reactivity in food-sensitized infants. *J Allergy Clin Immunol* 2015;135(5):1319–28.e12.
159. Martino D, Joo JE, Sexton-Oates A, Dang T, Allen K, Saffery R, Prescott S. Epigenome-wide association study reveals longitudinally stable DNA methylation differences in CD4+ T cells from children with IgE-mediated food allergy. *Epigenetics* 2014;9(7):998–1006.
160. Petrus NCM, Henneman P, Venema A, Mul A, Van Sinderen F, Haagsmans M, Mook O, Hennekam RC, Sprickelman AB, Mannens M. Cow's milk allergy in Dutch children: an epigenetic pilot survey. *Clin Transl Allergy* 2016;6(1):16.
161. Canani RB, Paparo L, Nocerino R, Cosenza L, Pezzella V, Di Costanzo M, Capasso M, Del Monaco V, D'Argenio V, Greco L, et al. Differences in DNA methylation profile of Th1 and Th2 cytokine genes are associated with tolerance acquisition in children with IgE-mediated cow's milk allergy. *Clin Epigenetics* 2015;7(1):38.

162. Hong X, Ladd-Acosta C, Hao K, Sherwood B, Ji H, Keet CA, Kumar R, Caruso D, Liu X, Wang G, et al. Epigenome-wide association study links site-specific DNA methylation changes with cow's milk allergy. *J Allergy Clin Immunol* 2016;138(3):908–11.e9.
163. Van Bilsen JHM, Sienkiewicz-Szłapka E, Lozano-Ojalvo D, Willemsen LEM, Antunes CM, Molina E, Smit JJ, Wróblewska B, Wichers HJ, Knol EF, et al. Application of the adverse outcome pathway (AOP) concept to structure the available in vivo and in vitro mechanistic data for allergic sensitization to food proteins. *Clin Transl Allergy* 2017;7(1):13.
164. Paparo L, Nocerino R, Bruno C, Di Scala C, Cosenza L, Bedogni G, Di Costanzo M, Mennini M, D'Argenio V, Salvatore F, et al. Randomized controlled trial on the influence of dietary intervention on epigenetic mechanisms in children with cow's milk allergy: the EPICMA study. *Sci Rep* 2019;9:2828.
165. Syed A, Garcia MA, Lyu SC, Bucayu R, Kohli A, Ishida S, Berglund JP, Tsai M, Maecker H, O'Riordan G, et al. Peanut oral immunotherapy results in increased antigen-induced regulatory T-cell function and hypomethylation of forkhead box protein 3 (FOXP3). *J Allergy Clin Immunol* 2014;133(2):500–10.e11.
166. Paparo L, Nocerino R, Cosenza L, Aitoro R, D'Argenio V, Del Monaco V, Di Scala C, Amoroso A, Di Costanzo M, Salvatore F, et al. Epigenetic features of FoxP3 in children with cow's milk allergy. *Clin Epigenetics* 2016;8:86.
167. Alag A. Machine learning approach yields epigenetic biomarkers of food allergy: a novel 13-gene signature to diagnose clinical reactivity. *PLoS One* 2019;14(6):e0218253.
168. Basturk A, Isik İ, Atalay A, Yilmaz A. Investigation of the efficacy of *Lactobacillus rhamnosus* GG in infants with cow's milk protein allergy: a randomised double-blind placebo-controlled trial. *Probiotics Antimicrob Proteins* 2020;12(1):138–43.
169. Fox AT, Wopereis H, Van Ampting MTJ, Oude Nijhuis MM, Butt AM, Peroni DG, Vandenplas Y, Candy DCA, Shah N, West CE, et al. A specific synbiotic-containing amino acid-based formula in dietary management of cow's milk allergy: a randomized controlled trial. *Clin Trans Allergy* 2019;9:5.
170. Schmidt RM, Pilmann Laursen R, Bruun S, Larnkjær A, Mølgaard C, Michaelsen KF, Høst A. Probiotics in late infancy reduce the incidence of eczema: a randomized controlled trial. *Pediatr Allergy Immunol* 2019;30(3):335–40.
171. Kuitunen M, Kukkonen K, Juntunen-Backman K, Korpela R, Poussa T, Tuure T, Haahtela T, Savilahti E. Probiotics prevent IgE-associated allergy until age 5 years in cesarean-delivered children but not in the total cohort. *J Allergy Clin Immunol* 2009;123(2):335–41.
172. Qamer S, Deshmukh M, Patole S. Probiotics for cow's milk protein allergy: a systematic review of randomized controlled trials. *Eur J Pediatr* 2019;178(8):1139–49.
173. Zhang GQ, Hu HJ, Liu CY, Zhang Q, Shakya S, Li ZY. Probiotics for prevention of atopy and food hypersensitivity in early childhood: a PRISMA-compliant systematic review and meta-analysis of randomized controlled trials. *Medicine (Baltimore)* 2016;95(8):e2562.
174. Zuccotti G, Meneghin F, Aceti A, Barone G, Callegari ML, Di Mauro A, Fantini MP, Gori D, Indrio F, Maggio L, et al. Probiotics for prevention of atopic diseases in infants: systematic review and meta-analysis. *Allergy* 2015;70(11):1356–71.