

A note on subgroups of automorphism groups of full shifts

Ville Salo
University of Chile

August 1, 2016

Abstract

We discuss the set of subgroups of the automorphism group of a full shift, and submonoids of its endomorphism monoid. We prove closure under direct products in the monoid case, and free products in the group case. We also show that the automorphism group of a full shift embeds in that of an uncountable sofic shift. Some undecidability results are obtained as corollaries.

1 Introduction

Automorphism groups of two-sided full shifts were first studied in [2], and the transitive SFT case was later studied in more detail in [1]. More results on them were shown in [4]. In particular, these papers prove interesting results about the set of subgroups of $\text{Aut}(S^{\mathbb{Z}})$. It is known for example that (non-abelian) free groups, finite groups and finitely generated abelian groups can be embedded in this group [2, 1], and so can all locally finite residually finite countable groups [4]. Other examples are ‘graph groups’

$$\langle (g_i)_{i \in \mathbb{N}} \mid (g_i g_j = g_j g_i)_{(i,j) \in R} \rangle$$

where $R \subset \mathbb{N}^2$ is arbitrary, fundamental groups of 2-manifolds, and the free product of all finite groups [4]. Despite the long list of examples, to our knowledge there is no known characterization of the set of subgroups of $\text{Aut}(S^{\mathbb{Z}})$.

In the absence of a characterization, we turn to closure properties. One closure property for this set of groups is proved explicitly in [4], namely closure under extensions by finite groups. Another one can be found by a direct application of the ideas of [1, 4], namely closure under direct sums. We prove this more generally for the endomorphism monoid.

Theorem 1. *For any alphabet S , the set of submonoids of $\text{End}(S^{\mathbb{Z}})$ is closed under direct products.*

Our main result is that the set of subgroups of the automorphism groups is also closed under free products, which answers a question I asked in [9].

Theorem 2. *For any alphabet S , the set of subgroups of $\text{Aut}(S^{\mathbb{Z}})$ is closed under free products.*

The closest result of this type in the literature seems to be the embedding of the free product of all finite groups into the automorphism group of a full shift. We are not aware of a published proof, but the result is mentioned in [4] and is attributed to R. C. Alperin.

The results work not just with binary products, but arbitrary countable products: we prove that in general a set of submonoids of a monoid is closed under direct (resp. free) products if and only if it is closed under countable direct (resp. countable free) products.

In [4], the group $\text{Aut}(S^{\mathbb{Z}})$ is embedded into the automorphism group of a transitive SFT. With a similar proof, we extend this in two directions, by showing that $\text{End}(S^{\mathbb{Z}})$ is contained in the endomorphism monoid of every uncountable sofic shift.

Lemma 1. *Let X be any uncountable sofic shift, and $A^{\mathbb{Z}}$ any full shift. Then $\text{End}(A^{\mathbb{Z}}) \leq \text{End}(X)$.*

From our results it follows that this is optimal among sofic shifts in the sense that $\text{End}(A^{\mathbb{Z}})$ cannot be computably embedded into $\text{End}(X)$ when X is countable.¹

As corollaries of the results, we obtain undecidability results for cellular automata on uncountable sofic shifts. The most important of these is that the embedding of free products is undecidable.

Theorem 3. *Given two local rules $F : S^{2r+1} \rightarrow S$ and $G : S^{2s+1} \rightarrow S$ that define reversible cellular automata f and g (respectively) on $S^{\mathbb{Z}}$, it is undecidable whether f and g satisfy a nontrivial relation, and whether $\langle f, g \rangle \cong F_2$.*

Given our embedding result, this follows for essentially abstract reasons from the known fact that the torsion problem is undecidable on full shifts [3].

The ‘marker method’ (the use of unbordered words) usually plays an important role in the construction of endomorphisms, and this note makes no exception. However, of even more importance in our constructions is the concept of ‘conveyor belts’. This technique is implicit in many constructions found in the literature, and variants of it are found for example in [1, 4, 6]. We make this idea more explicit.

While our main interest is in automorphism groups, we prove results for the whole endomorphism monoid when possible, since the results about monoids are strictly stronger, and the proofs are typically the same. In some results, however, the fact we consider groups instead of monoids is important. For example, we do not know whether the set of submonoids of the endomorphism monoid of a full shift is closed under free product.

We also leave some open questions. Of course, our main question² is whether there is a characterization of the whole set of subgroups:

Question 1. *Which groups are subgroups of $\text{Aut}(S^{\mathbb{Z}})$?*

This seems (to us) to be hard, and we do not know whether it can be approached with the current methods. A question that seems more approachable,

¹We will also rule out the existence of abstract embeddings in later work, see Remark 1.

²We state this in a question block as it is our main motivating question, but it is not a rigorous one, but rather an open-ended characterization problem. It does not necessarily have a ‘correct’ answer, and should be judged on an ‘I know it when I see it’ basis.

but that we have not succeeded in solving, is the generalization of Lemma 1 to other pairs of sofic shifts: while it is easy to embed the automorphism group of a full shift into that of an uncountable sofic shift, we are rarely able to do the converse.

Question 2. *Let X and Y be two sofic shifts. When do we have $\text{Aut}(X) \leq \text{Aut}(Y)$?*

The question is discussed briefly in [4], where it is noted that the embedding proof for the full shift does not directly generalize.

On the side of computability, we are also interested in the following natural follow-up question to Theorem 3:

Question 3. *Let A be the set of pairs of CA on $S^{\mathbb{Z}}$ generating a finite group, and B the set of pairs generating F_2 . Are A and B recursively inseparable?*

2 Definitions

The letters A, B, C and S stand for finite alphabets. We write A^* for the set of finite words over the alphabet A , and A^+ the set of nonempty words. The concatenation of two words u, v is written as $u \cdot v$ or simply uv , and *points* or *configurations* $x \in A^{\mathbb{Z}}$ can be written as infinite concatenations

$$x = \dots w_{-3} \cdot w_{-2} \cdot w_{-1} \cdot w_0 \cdot w_1 \cdot w_2 \dots$$

where the ‘decimal point’ in $w_{-1} \cdot w_0$ need not be at the origin, but simply denotes some fixed position of the point.

If $u \in A^*$ is any word, write u^R for the reversed word $u_i^R = u_{|u|-1-i}$.

Let S be a finite alphabet. Then $S^{\mathbb{Z}}$ with the product topology is called a *full shift*. We define the *shift* $\sigma : S^{\mathbb{Z}} \rightarrow S^{\mathbb{Z}}$ by $\sigma(x)_i = x_{i+1}$. A *subshift* is a topologically closed set $X \subset S^{\mathbb{Z}}$ satisfying $\sigma(X) = X$.

If $x \in X$ we write $w \sqsubset x$ if $\exists i : w = x_{[i, i+|w|-1]}$. We also write $w \sqsubset X$ and $u \sqsubset v$ for words u, v , with obvious meanings, and

$$\mathcal{L}_n(X) = \{w \in S^n \mid w \sqsubset X\}.$$

A *cellular automaton* on a subshift $X \subset S^{\mathbb{Z}}$ is a continuous function $f : X \rightarrow X$ that commutes with σ . Equivalently, it has a *radius* $r \in \mathbb{N}$ and a *local rule* $F : \mathcal{L}_{2r+1}(X) \rightarrow S$ such that $f(x)_i = F(x_{[i-r, i+r]})$ for all $x \in X, i \in \mathbb{Z}$.

For a standard reference on symbolic dynamics, see [7, 5].

We assume the reader is familiar with groups and monoids, but give the basic definitions to clarify our choice of boundary between properties and structure. A *monoid* $(M, \cdot, 1_M)$ is a triple where M is a countable set, \cdot is an associative multiplication operation $(a, b) \mapsto a \cdot b = ab$ and 1_M is an *identity element* satisfying $a1_M = 1_M a = a$ for all $a \in M$. A *group* is a monoid $(G, \cdot, 1_G)$ where for each element $g \in G$ there is an inverse g^{-1} satisfying $gg^{-1} = g^{-1}g = 1_G$. In other words, we think of the existence of g^{-1} as simply a property of the element g , not as an operation. If N and M are monoids, a *homomorphism* $\phi : N \rightarrow M$ is a map satisfying $\phi(a \cdot b) = \phi(a) \cdot \phi(b)$ and $\phi(1_N) = 1_M$.

A *submonoid* of a monoid M is a subset of N that contains the identity element 1_M of M and is closed under multiplication, so that N obtains a monoid

structure from M with $1_N = 1_M$, and we write $N \leq M$. More generally, we write $N \leq M$ if there is an *embedding*, or an injective homomorphism $\phi : N \rightarrow M$. A bijective homomorphism is called an *isomorphism*, and we write $M \cong N$ if M and N are isomorphic. A *subgroup* is a submonoid that is a group. If M is a monoid, write $\text{sub}_m(M)$ for the class of isomorphism classes of its submonoids. We write $\text{sub}_g(M)$ for the class of isomorphism classes of its subgroups.

The *endomorphism monoid* $\text{End}(X)$ of a subshift X is the (discrete) monoid of cellular automata on it under function composition, with $1_{\text{End}(X)} = \text{id}_X$, the identity map on X . The *automorphism group* $\text{Aut}(X)$ of X is the subgroup of $\text{End}(X)$ containing the invertible elements:

$$\text{Aut}(X) = \{f \in \text{End}(X) \mid \exists g \in \text{End}(X) : f \circ g = g \circ f = \text{id}_X\}.$$

By compactness of X , this is precisely the set of elements of $\text{End}(X)$ that are bijective. When $X = S^{\mathbb{Z}}$, this is further precisely the set of injective CA [2].

Our definitions are set up so that the following holds.³

Lemma 2. *Let G be a group such that $G \leq \text{End}(X)$. Then $G \leq \text{Aut}(X)$. In particular, if $\text{sub}_m(\text{End}(X)) = \text{sub}_m(\text{End}(Y))$ then $\text{sub}_g(\text{Aut}(X)) = \text{sub}_g(\text{Aut}(Y))$.*

The following lemma is useful for defining cellular automata.

Lemma 3. *Let $X = S^{\mathbb{Z}}$ and let $Y \subset X$ be a subset such that $\sigma(Y) = Y$ and $\bar{Y} = X$. If $f : Y \rightarrow Y$ is uniformly continuous and commutes with the shift, then there is a unique continuous map $g : X \rightarrow X$ such that $g|_Y = f$, and it is a cellular automaton.*

Write $S(A)$ for the set of permutations of A , and $S(A, B)$ for the set of all bijections $c : A \rightarrow B$. The free group with m generators is written F_m , and F_∞ is the free group with a countably infinite set of generators.

The (external) *direct product* of two monoids M, N is the monoid $M \times N$ with operation $(a, b) \cdot (c, d) = (ac, bd)$. The *free product* of monoids M and N is defined up to isomorphism as follows: Let $M \cong \langle 1_M, a_1, a_2, a_3, \dots \mid r_1, r_2, \dots \rangle$ be a presentation, where each r_i is a relation of the form $u = v$ where $u, v \in \{a_1, a_2, \dots\}^*$, and similarly let $N \cong \langle 1_N, b_1, b_2, \dots \mid t_1, t_2, \dots \rangle$. Then

$$M * N \cong \langle a_1, b_1, a_2, b_2, a_3, b_3, \dots \mid 1_M = 1_N, r_1, t_1, r_2, t_2, r_3, t_3, \dots \rangle.$$

We also write M^k for the direct product of k copies of M . There are obvious embeddings $M^{k-1} \rightarrow M^k$, and their direct limit is written as M^ω . The elements of the countable monoid M^ω are vectors of finite support with component values in M . Write $M * M$, M^{*k} and $M^{*\omega}$ for the corresponding concepts for the free product.

3 Lemmas about submonoids

Lemma 4. *Let M and N be monoids. Then $\text{sub}_m(M) = \text{sub}_m(N)$ if and only if $M \leq N$ and $N \leq M$.*

³For this, it is (at least a priori) important that the identity element of a subgroup of $\text{End}(X)$ is the identity map – thus, the identity element must be part of the structure of a monoid, rather than a property.

In particular, to show that two groups have the same subgroups, we only need to show they contain each other as subgroups. We note that the set of subgroups of a group is not a complete invariant for group isomorphism: for example, free groups with different amounts of generators are non-isomorphic, but contain each other as subgroups.

In this note, we concentrate on automorphism groups of full shifts $\text{Aut}(S^{\mathbb{Z}})$ where S is a finite alphabet of size at least 2. It is not known when two such groups $\text{Aut}(A^{\mathbb{Z}})$ and $\text{Aut}(B^{\mathbb{Z}})$ are isomorphic – in particular the case $|A| = 2, |B| = 3$ is open. Nevertheless, the set of subgroups, and more generally submonoids, is always the same:

Lemma 5. *Let A, B be alphabets of size at least 2. Then*

$$\text{sub}_m(\text{End}(A^{\mathbb{Z}})) = \text{sub}_m(\text{End}(B^{\mathbb{Z}})).$$

In particular, it follows that $\text{sub}_g(\text{End}(A^{\mathbb{Z}})) = \text{sub}_g(\text{End}(B^{\mathbb{Z}}))$ if $|A|, |B| \geq 2$. Lemma 5 follows directly from Lemma 8, which we prove in Section 7. The corollary $\text{sub}_g(\text{End}(A^{\mathbb{Z}})) = \text{sub}_g(\text{End}(B^{\mathbb{Z}}))$ is well-known, and can be found in [4] as Corollary 1.

There is not much to say about finite submonoids and subgroups of $\text{End}(S^{\mathbb{Z}})$, as shown by the following result, which essentially already appears in [2]. The proof illustrates the usefulness of Lemma 5.

Proposition 1. *Let $|A| \geq 2$. Then every finite monoid embeds in $\text{End}(A^{\mathbb{Z}})$.*

Proof. Every finite monoid embeds in the transformation monoid of a finite set S , and thus in $\text{End}(S^{\mathbb{Z}})$. The result then follows from the previous lemma. \square

We note that in general, if $\text{sub}_m(M)$ is closed under binary direct or free products, then it is closed under the corresponding countable products.

Lemma 6. *Let $\text{sub}_m(M) = \text{sub}_m(N) = \text{sub}_m(N')$. Then the following are equivalent:*

- $N \times N' \in \text{sub}_m(M)$,
- $M \times M \in \text{sub}_m(M)$,
- $\text{sub}_m(M)$ is closed under finite direct products.
- $M^\omega \in \text{sub}_m(M)$,
- $\text{sub}_m(M)$ is closed under countable direct products.

The analogous result is true for free products.

Proof. We give the proof for direct products, the case of free products being similar. The equivalence of the first two conditions and the equivalence of the last two conditions are direct, as is the fact that the last two conditions imply the first two. We show that the third and fourth condition follow from the second.

For this, suppose that $M \times M \leq M$. Let $\phi_0 : M \rightarrow M$ and $\phi_1 : M \rightarrow M$ be the embeddings giving the embedding $M \times M \leq M$, that is, $\phi_0(M) \cap \phi_1(M) = \{1_M\}$ and $\phi_0(a)\phi_1(b) = \phi_1(b)\phi_0(a)$ for all $a, b \in M$. If this holds for two maps, we say the *embedding conditions* hold for them. For $w \in \{0, 1\}^*$ define

inductively $\phi_{0w} = \phi_0 \circ \phi_w$ and $\phi_{1w} = \phi_1 \circ \phi_w$. Then all maps ϕ_w are embeddings of M into itself.

For $i \in \mathbb{N}$, define $\psi_i = \phi_{1^i 0}$. It is easy to show that the embedding conditions hold for ψ_i and ψ_j whenever $i \neq j$. It follows that $(\psi_i)_{i \in [0, k-1]}$ gives an embedding of M^k into M , and $(\psi_i)_{i \in \mathbb{N}}$ of M^ω into M .

The claims for free products are proved analogously, but using a different embedding condition, namely that the images of the embeddings satisfy no nontrivial relations. \square

The importance of this lemma and Lemma 5 is that we do not need to worry about changing the alphabets of our full shifts when proving closure properties or about whether we use finite or countable products.

4 Conveyor belts

In our constructions, we will typically embed one automorphism group into another. In practise, this means that in the configurations of one full shift $S^\mathbb{Z}$, we identify subsequences that code (parts of) configurations from another full shift $A^\mathbb{Z}$. In these subsequences, we apply $f \in \text{End}(A^\mathbb{Z})$. To make this into a homomorphism from $\text{End}(A^\mathbb{Z})$ to $\text{End}(S^\mathbb{Z})$, it is important to have natural behavior at the boundary between an area coding (part of) a configuration in $A^\mathbb{Z}$, and an area containing something else. For this, we use conveyor belts.

Definition 1. *Let A be any alphabet. A conveyor belt over A is a word over the alphabet A^2 , that is, $w \in (A^2)^*$. Write $\text{Conv}_{A,k}$ for the set of conveyor belts over A of length k , that is $\text{Conv}_{A,k} = (A^2)^k$. For a cellular automaton $f : A^\mathbb{Z} \rightarrow A^\mathbb{Z}$ and $k \in \mathbb{N}$, we define a function $f_k : \text{Conv}_{A,k} \rightarrow \text{Conv}_{A,k}$ as follows: if $w \in (A^2)^k$, we decompose w as $w = u \times v$ for some $u, v \in A^k$, and we define*

$$f_k(w) = f((uv^R)^\mathbb{Z})_{[0, k-1]} \times (f((uv^R)^\mathbb{Z})_{[k, 2k-1]})^R.$$

Applying a CA to a conveyor belt can alternatively be described as applying its local rule on the first track, applying its local rule in reverse on the second track, and gluing the tracks at the borders in the obvious way, as if the word were laid down on a conveyor belt: it is clear that this is essentially the same as applying the CA to a periodic point of even period. The following is then clear.

Lemma 7. *The map $f \mapsto f_k$ is a monoid homomorphism from $\text{End}(A^\mathbb{Z})$ to the monoid of functions on $\text{Conv}_{A,k}$, and*

$$f \mapsto (f_k)_{k \in \mathbb{N}} : \text{End}(A^\mathbb{Z}) \rightarrow \prod_k S(\text{Conv}_{A,k})$$

is an embedding. Furthermore, the maps f_k are uniformly continuous in the sense that there exists a radius $r \in \mathbb{N}$ such that for all $k \in \mathbb{N}$ and $w, w' \in \text{Conv}_{A,k}$ such that

$$w_{[\max(0, i-r), \min(k-1, i+r)]} = w'_{[\max(0, i-r), \min(k-1, i+r)]},$$

we have $f_k(w)_i = f_k(w')_i$.

The radius in the lemma can be taken to be just the usual radius of $f \in \text{Aut}(A^\mathbb{Z})$.

5 Direct products

Theorem 4. *If $|S| \geq 2$, $\text{sub}_m(\text{End}(S^{\mathbb{Z}}))$ is closed under direct products.*

Proof. By Lemma 5 and Lemma 6, it is enough to show that for two disjoint alphabets A and B , the monoid $\text{End}(A^{\mathbb{Z}}) \times \text{End}(B^{\mathbb{Z}})$ embeds into $\text{End}(S^{\mathbb{Z}})$ for some alphabet S . We choose $S = A^2 \sqcup B^2$. To prove the claim, it is enough to give embeddings $f \mapsto f_A : \text{End}(A^{\mathbb{Z}}) \rightarrow \text{End}(S^{\mathbb{Z}})$ and $f \mapsto f_B : \text{End}(B^{\mathbb{Z}}) \rightarrow \text{End}(S^{\mathbb{Z}})$ such that $f_A \circ g_B = g_B \circ f_A$ for all $f \in \text{End}(A^{\mathbb{Z}})$ and $g \in \text{End}(B^{\mathbb{Z}})$, and $f_A = g_B \implies f = \text{id}_{A^{\mathbb{Z}}} \wedge g = \text{id}_{B^{\mathbb{Z}}}$.

Let $f \in \text{End}(A^{\mathbb{Z}})$. Write Y for the set of points $x \in (A^2 \sqcup B^2)^{\mathbb{Z}}$ which are not left or right asymptotic to a point over A^2 or B^2 , in other words, points where all continuous runs over one of the subalphabets A^2 or B^2 are finite. If $x \in Y$, then we can write

$$x = \dots w_{-3} \cdot w_{-2} \cdot w_{-1} \cdot w_0 \cdot w_1 \cdot w_2 \dots$$

where $w_{2i} \in (A^2)^+$ and $w_{2i+1} \in (B^2)^+$ for all $i \in \mathbb{Z}$ (and the decimal point need not be at the origin), and we define

$$f_A(x) = \dots w_{-3} \cdot f_{|w_{-2}|}(w_{-2}) \cdot w_{-1} \cdot f_{|w_0|}(w_0) \cdot w_1 \cdot f_{|w_2|}(w_2) \cdot w_3 \dots$$

(where the decimal point is in the same position as in x). It is clear that f_A commutes with the shift on Y . From Lemma 7, it follows that f_A is uniformly continuous on Y , and thus extends uniquely to a cellular automaton on X . This gives a function from $\text{End}(A^{\mathbb{Z}})$ to $\text{End}(S^{\mathbb{Z}})$. From Lemma 7, it easily follows that this is an embedding.

We symmetrically construct a homomorphism from $\text{Aut}(B^{\mathbb{Z}})$ to $\text{Aut}(S^{\mathbb{Z}})$ by rewriting the contiguous segments over B^2 . It is clear that these mappings commute, and since f_A fixes all symbols from B^2 and g_B all symbols from A^2 , only the identity map is in the image of both embeddings. \square

Corollary 1. *If $|S| \geq 2$, $\text{sub}_g(\text{Aut}(S^{\mathbb{Z}}))$ is closed under direct products.*

6 Free products

In this section, we prove our main theorem:

Theorem 5. *If $|S| \geq 2$, $\text{sub}_g(\text{Aut}(S^{\mathbb{Z}}))$ is closed under free products.*

In the direct product case, we had two kinds of conveyor belts, and we applied the f_A and g_B maps completely independently on both of them. To embed a free product, the idea is to have these conveyor belts talk to each other, so that any alternating product of elements from the $(f \mapsto f_A)$ -embedding and the $(g \mapsto g_B)$ -embedding can transmit information arbitrarily far over an alternating sequence of A -conveyor belts and B -conveyor belts, as long as the lengths and contents of these belts are chosen suitably.

For this, we increase the size of our alphabet, and add a C -component that allows us to transmit any kind of modification of the left end of an A -conveyor belt to any kind of modification of the right end of a B -conveyor belt on its left, and vice versa. Let us proceed to the details.

Proof. By Lemma 5, it is enough to show how to embed the free product of $\text{Aut}(A^{\mathbb{Z}})$ and $\text{Aut}(B^{\mathbb{Z}})$ into $\text{Aut}(S^{\mathbb{Z}})$, where A and B are two disjoint alphabets with the same cardinality and S is an alphabet of our choosing. We choose an arbitrary abelian group structure on both A^2 and B^2 . Let $C = S(A^2, B^2)$, and choose the alphabet

$$S = C \sqcup A^2 \sqcup B^2.$$

Similarly as in the proof of Theorem 4, for each $f \in \text{Aut}(A^{\mathbb{Z}})$, we will define a CA $f_A : S^{\mathbb{Z}} \rightarrow S^{\mathbb{Z}}$ so that $f \mapsto f_A$ embeds $\text{Aut}(A^{\mathbb{Z}})$ into $\text{Aut}(S^{\mathbb{Z}})$ as a subgroup $G_A \leq \text{Aut}(S^{\mathbb{Z}})$. An embedding $f \mapsto f_B$ of $\text{Aut}(B^{\mathbb{Z}})$ into $G_B \leq \text{Aut}(S^{\mathbb{Z}})$ is defined symmetrically by swapping the roles of A and B , and $G_A \cap G_B = \{\text{id}\}$ will be clear from the construction. To show they generate the free product of $\text{Aut}(A^{\mathbb{Z}})$ and $\text{Aut}(B^{\mathbb{Z}})$ in $\text{Aut}(S^{\mathbb{Z}})$, we must show there are no non-trivial relations between elements of G_A and G_B . Because the roles of A and B are symmetric, this will follow from showing that for any

$$f = f_{k-1} \circ f_{k-2} \circ \cdots \circ f_1 \circ f_0 \in \text{Aut}(S^{\mathbb{Z}})$$

where $k \geq 1$, and $f_i \in G_A \setminus \{\text{id}\}$ for even i and $f_i \in G_B \setminus \{\text{id}\}$ for odd i , we have $f \neq \text{id}$.

Let $f \in \text{Aut}(A^{\mathbb{Z}})$. Let Y be the set of points where contiguous runs over each of the subalphabets A^2 , B^2 and C are finite. The set Y is shift-invariant, so to define cellular automata on $S^{\mathbb{Z}}$, it is enough to define shift-invariant uniformly continuous maps on Y . Let thus $x \in Y$, so that

$$x = \dots w_{-2} \cdot w_{-1} \cdot w_0 \cdot w_1 \cdot w_2 \dots$$

where for all $i \in \mathbb{Z}$, $w_i \in (A^2)^+$, $w_i \in (B^2)^+$ or $w_i \in C^+$, and w_i and w_{i+1} are over different alphabets. We write

$$f(x) = \dots u_{-2} \cdot u_{-1} \cdot u_0 \cdot u_1 \cdot u_2 \dots,$$

where $|u_i| = |w_i|$ for all i . If $w_i \in C^+$, we let $u_i = w_i$. If $w_i \in (A^2)^+$, we apply f to the conveyor belt w_i as in the proof of Theorem 4, and let $u_i = f_{|w_i|}(w_i)$.

Finally, let us define u_i for $w_i \in (B^2)^+$. Suppose $|u_i| = \ell$. First, $(u_i)_j = (w_i)_j$ for $j \in [0, \ell - 2]$, that is, only the last symbol may change. If $w_{i+1} \notin C$ (that is, either $w_{i+1} \notin C^+$ or $w_{i+1} \in C^+$ but $|w_{i+1}| > 1$) or $w_{i+2} \notin (A^2)^+$, we also let $(u_i)_{\ell-1} = (w_i)_{\ell-1}$. If $w_{i+2} \in (A^2)^+$ and $w_{i+1} = c \in C$, we let

$$(u_i)_{\ell-1} = (w_i)_{\ell-1} - c((w_{i+2})_0) + c((u_{i+2})_0),$$

where addition is performed with respect to the abelian group structure of B^2 . It is easy to see that this defines f_A on Y , and since the map defined is uniformly continuous and shift-commuting, we can extend it in a unique way to a CA $f_A : S^{\mathbb{Z}} \rightarrow S^{\mathbb{Z}}$. If $f \in \text{Aut}(A^{\mathbb{Z}})$, it has an inverse $f^{-1} \in \text{Aut}(A^{\mathbb{Z}})$, and it is easy to see that $f_A|_Y \circ f_A^{-1}|_Y = f_A^{-1}|_Y \circ f_A|_Y = \text{id}_Y$. It follows that $f_A \in \text{Aut}(S^{\mathbb{Z}})$, because the only extension of the identity map on Y to a CA on $S^{\mathbb{Z}}$ is the identity map.

For $g \in \text{Aut}(B^{\mathbb{Z}})$, the map $g_B : S^{\mathbb{Z}} \rightarrow S^{\mathbb{Z}}$ is defined symmetrically, modifying the rightmost symbol of a word over A^2 as a function of the leftmost symbol of a word over B^2 to the right of it when separated by the single symbol $c \in C$, by the same formula, but using the abelian structure of A^2 instead of that of B^2 , and the function $c^{-1} : B^2 \rightarrow A^2$ instead of c .

We now show that G_A and G_B indeed give the free product of $\text{Aut}(A^{\mathbb{Z}})$ and $\text{Aut}(B^{\mathbb{Z}})$. Suppose thus that

$$f = f_{k-1} \circ f_{k-2} \circ \cdots \circ f_1 \circ f_0 \in \text{Aut}(S^{\mathbb{Z}})$$

where $k \geq 1$, and $f_i \in G_A \setminus \{\text{id}\}$ for even i and $f_i \in G_B \setminus \{\text{id}\}$ for odd i . We need to show $f \neq \text{id}$. Suppose the minimal radius of f_i is $r_i \geq 0$. We show that $f(x)_0$ depends on at least the cell x_h where $h = \sum_i r_i + 2k$. Clearly this will imply that $f \neq \text{id}$.

For this, we will define two points

$$x = y \cdot u_{k-1} \cdot c_{k-2} u_{k-2} \cdots c_1 u_1 \cdot c_0 u_0 \cdot z$$

$$x' = y \cdot u_{k-1} \cdot c_{k-2} u_{k-2} \cdots c_1 u_1 \cdot c_0 u'_0 \cdot z$$

where $y \in C^{-\mathbb{N}}$, $z \in C^{\mathbb{N}}$, $u_i \in (A^2)^+$ if i is even, $u_i \in (B^2)^+$ if i is odd, $|u'_0| = |u_0|$, $(u_0)_i = (u'_0)_i$ for $i \neq |u_0| - 1$, and $c_i \in C$ for all i . The choices of y and z are arbitrary, and $|u_i| = r_i$ for all i . We will choose the words u_i and symbols c_i carefully so that $f(x)_0 \neq f(x')_0$, while $x_i = x'_i$ for all $i \neq h$. Write $I_i = [\alpha_i, \beta_i]$ for the interval where u_i occurs in x .

Let $i \in [0, k-1]$ be even, and let v_i and v'_i be words of length $2r_i + 1$ with $(v_i)_j = (v'_i)_j$ for $j \in [1, 2r_i - 1]$ such that $F_i(v_i) \neq F_i(v'_i)$, where $F_i : A^{2r_i+1} \rightarrow A$ is the local rule of f_i . We then have $(v_i)_0 \neq (v'_i)_0$ or $(v_i)_{2r_i} \neq (v'_i)_{2r_i}$, and if $r_i > 0$ we can (and do) suppose $(v_i)_0 = (v'_i)_0$ or $(v_i)_{2r_i} = (v'_i)_{2r_i}$. If $(v_i)_0 \neq (v'_i)_0$, i is called a *left dependence* and otherwise a *right dependence*.

If i is a left dependence, define the words $t_i, t'_i \in (A^2)^{r_i+1}$ by

$$(t_i)_j = \begin{cases} ((v_i)_{r_i+j+1}, (v_i)_{r_i-j}) & \text{if } j < r_i \\ (a, (v_i)_0) & \text{if } j = r_i, \end{cases}$$

where a is arbitrary, and t'_i by the same formula using the word v'_i . In other words, the word t_i is a conveyor belt version of v_i where the center of v_i is at the left end of t_i and the leftmost coordinate is at the right end of t_i .

The important property of t_i and t'_i is that they agree apart from their rightmost coordinates, but after applying f_i to the conveyor belt, the images differ in the leftmost coordinate, that is,

$$(t_i)_{[0, r_i-1]} = (t'_i)_{[0, r_i-1]} \wedge (f_i)_{|t_i|}(t_i)_0 \neq (f_i)_{|t_i|}(t'_i)_0.$$

If i is a right dependence, we produce words t_i and t'_i with this property in a similar fashion. Note that when $r_i > 0$, it was important that we chose the words v_i and v'_i so that they differ in one end but not the other, as otherwise $(t_i)_{r_i-1} \neq (t'_i)_{r_i-1}$.

Choose c_i so that $c_i((f_i)_{|t_i|}(t_i)_0) = 0$ and

$$c_i((f_i)_{|t_i|}(t'_i)_0) = (t'_{i+1})_{r_{i+1}} - (t_{i+1})_{r_{i+1}}.$$

For odd coordinates i , we choose words $t_i, t'_i \in (B^2)^+$ and $c_i \in C$ similarly, switching the roles of A and B .

We let $u_0 = t_0$, $u'_0 = t'_0$. The words u_i will be chosen inductively in such a way that

$$f_{i-1} \circ \cdots \circ f_1 \circ f_0(x)_{I_i} = t_i,$$

$$f_{i-1} \circ \cdots \circ f_1 \circ f_0(x')_{I_i} = t'_i,$$

and

$$f_{i-1} \circ \cdots \circ f_1 \circ f_0(x)_{I_j} = f_{i-1} \circ \cdots \circ f_1 \circ f_0(x')_{I_j}$$

for $j > i$.

To see this is possible, observe that information travels only from right to left over C -symbols when maps from $\langle G_A, G_B \rangle$ are applied, and a map $f \in (G_A \cup G_B)^k$ will move information over at most k symbols in C . Thus, when applying our maps $g_\ell = f_{\ell-1} \circ \cdots \circ f_1 \circ f_0$ to the points x, x' , we will automatically have $g_\ell(x)_{I_j} = g_\ell(x')_{I_j}$ for all $j > \ell$. It follows that it is enough to show that the u_i can be chosen so that $g_\ell(x)_{I_\ell} = t_\ell$ for all ℓ : by the choice of the symbols c_i , we will then automatically have $g_\ell(x')_{I_\ell} = t'_\ell$.

But naturally we can choose such words u_i by induction on i , since each of the maps g_ℓ is reversible and information travels only to the left over symbols in C . \square

In [1], it is shown that $\mathbb{Z}_2 * \mathbb{Z}_2 * \mathbb{Z}_2$ embeds in $\text{Aut}(X)$ for a full shift X , and thus also the two-generator free group does. More generally, it is known that every free product of finitely many finite groups embeds in $\text{Aut}(X)$ for a full shift X . In [4], this is attributed to R. C. Alperin. Combining the previous theorem and Proposition 1 gives a new proof of this result.

Corollary 2. *If $|S| \geq 2$, every free product of finite groups embeds in $\text{Aut}(S^{\mathbb{Z}})$.*

7 Embeddings between endomorphism groups

In [4], it was shown that automorphism groups of full shifts embed in those of transitive SFTs. With essentially the same proof, we show that endomorphism monoids of full shifts embed in those of uncountable sofic shifts, equivalently in ones with positive entropy. The proof relies on a number of basic properties of sofic shifts, which can be found in [7, 5].

Lemma 8. *Let X be any uncountable sofic shift, and $A^{\mathbb{Z}}$ any full shift. Then $\text{End}(A^{\mathbb{Z}}) \leq \text{End}(X)$.*

Proof. It is easy to see that for a sofic shift, uncountability is equivalent to having positive entropy.

The *syntactic monoid* of a subshift X is the monoid whose elements are equivalence classes of words in X under the equivalence $w \sim w' \iff \forall u, v : uwv \sqsubset X \iff uw'v \sqsubset X$. Sofic shifts are characterized as the subshifts with a finite syntactic monoid. If X has positive entropy, then its minimal SFT cover Y also does. Then Y has a transitive component with positive entropy, and its image in the covering map is a positive-entropy transitive sofic subshift Z of X .

Let k be such that if $u, v \sqsubset Z$ then $uav \sqsubset Z$ for some word a with $|a| \leq k$ – such k exists because Z is transitive and because its syntactic monoid is finite. Let m be such that Z contains at least $n = |A^2|$ words of length m representing the same element of the syntactic monoid of X – such m exists because the syntactic monoid of X is finite,⁴ and because Z has positive entropy. Let w' be

⁴It is important that the words indeed represent the same element in the syntactic monoid of X and not just Z – we will need to be able to interchange them in any configuration of X .

a *synchronizing word* in Z of some length ℓ , that is, such that

$$u_1 w', w' u_2 \sqsubset Z \implies u_1 w' u_2 \sqsubset Z.$$

Such a word can be found in any sofic shift [5].

In every aperiodic infinite word, one can find unbordered words of arbitrary length [8]. Take any configuration in Z which is aperiodic and where w' appears syndetically (in every long enough subword), to find an unbordered word $w \sqsubset Z$ of length at least $2k + m + 1$ containing the word w' . Then w is synchronizing, since it contains a synchronizing subword.

By the assumptions on k and m , there is a set of words wUw , where $U \subset S^p$ for some $p \leq 2k + m$ and $|U| = n$, such that all words in wUw occur in Z and represent the same element of the syntactic monoid of X . Note that two words in wUw can only overlap nontrivially by sharing the subword w . Since all words in wUw occur in Z and w is synchronizing, the language $(wU)^*w$ is contained in the language of Z .

Now, let $U = \{u_1, \dots, u_n\}$, and fix a bijection $\phi : [1, n] \rightarrow A^2$. Given a CA $f : A^{\mathbb{Z}} \rightarrow A^{\mathbb{Z}}$, the embedding is now constructed as in the previous sections: if we have a maximal finite subword of the form

$$wu_{i_1}wu_{i_2}wu_{i_3}w \cdots wu_{i_\ell}w$$

(note that U -subwords of two such words cannot overlap by the assumptions), we apply f to the corresponding conveyor belt and let $v = f_\ell(\phi(i_1)\phi(i_2) \cdots \phi(i_\ell))$. We rewrite the word $wu_{i_1}w \cdots wu_{i_\ell}w$ by

$$wu_{\phi^{-1}(v_1)}wu_{\phi^{-1}(v_2)}wu_{\phi^{-1}(v_3)}w \cdots wu_{\phi^{-1}(v_\ell)}w.$$

As in the previous proofs, it is easy to check that this gives an embedding of $\text{End}(A^{\mathbb{Z}})$ into $\text{End}(X)$. \square

The converse $\text{End}(X) \leq \text{End}(A^{\mathbb{Z}})$ is not true in general for positive-entropy sofic shifts X :

Example 1. *The group $\text{Aut}(A^{\mathbb{Z}})$ is residually finite [1]. We show that $\text{End}(X)$ need not be residually finite for a sofic shift X . For this, let*

$$X_k = \{x \in \{0, 1\}^{\mathbb{Z}} \mid |\{i \in \mathbb{Z} \mid x_i = 1\}| \leq k\}.$$

Then $\text{Aut}(X_2)$ contains a copy of the group of all permutations of \mathbb{N} with finite support, by permuting the (orbits of) isolated points. This group is not residually finite, so $X_2 \times Y$ is not residually finite for any subshift Y . In particular, by letting Y be a positive-entropy sofic shift we obtain the result. \triangle

Even assuming transitivity, we are not aware of a general technique of embedding $\text{End}(X)$ into $\text{End}(Y)$ for two sofic shifts:

Question 4. *Let X and Y be two sofic shifts. When do we have $\text{Aut}(X) \leq \text{Aut}(Y)$?*

We are mainly interested in the case where X and Y are mixing SFTs. Even in this case, we find it tricky to find embeddings even between concrete examples. For example, let $X \subset \{x \in \{0, 1\}^{\mathbb{Z}} \mid 11 \not\sqsubset x\}$, the *golden mean SFT*. Do we have $\text{Aut}(X) \leq \text{Aut}(\{0, 1\}^{\mathbb{Z}})$?

8 Decidability

In this section, we briefly discuss some decidability corollaries for cellular automata that follow from the constructions. We fix the local rule of a CA as its computable presentation. This allows us to ask decidability questions about cellular automata. We start with a lemma that shows that the translation between cellular automata and their local rules is completely algorithmic. We omit the standard proof.

Lemma 9. *Let X be a sofic shift. Then given a function $F : S^{2r+1} \rightarrow S$, it is decidable whether $f(x)_i = F(x_{[i-r, i+r]})$ defines a cellular automaton on X , and if it does, we can compute a minimal radius $r' \in \mathbb{N}$ and a local rule $F' : \mathcal{L}_{2r'+1}(X) \rightarrow S$ for f .*

Of course, an algorithm that minimizes the local rule implies that given two local rules $F : S^{2r+1} \rightarrow S$ and $F' : S^{2r'+1} \rightarrow S$, it is decidable whether they define the same CA.

Let G be a countable group, with a fixed computable presentation for the elements. The *torsion problem* is the problem of, given $g \in G$, deciding whether there exists $m > 0$ such that $g^m = 1_G$.

Theorem 6 ([3]). *For some S , the torsion problem of $\text{Aut}(S^{\mathbb{Z}})$ is undecidable.*

Theorem 7 ([10]). *The torsion problem of $\text{Aut}(X)$ is decidable for every zero-entropy sofic shift X .*

Combining the theorems with Lemma 8 gives the following.

Theorem 8. *Let X be a sofic shift. Then the torsion problem of $\text{Aut}(X)$ is decidable if and only if X has zero entropy.*

Remark 1. *It follows that for sofic X , there can exist a computable embedding of $\text{Aut}(S^{\mathbb{Z}})$ into $\text{Aut}(X)$ only if X is uncountable. However, this does not prohibit the existence of an abstract embedding. We will address this problem in a later work, where we show that $\text{Aut}(X)$ is amenable when X is countable.*

Another definition of the torsion problem is to decide, given a CA, whether it generates a copy of \mathbb{Z} . Next, we discuss other problems of this type, omitting the easy proofs.

Proposition 2. *Given a finite set $F \subset \text{Aut}(S^{\mathbb{Z}})$ and a finite group G , it is decidable whether $\langle F \rangle \cong G$.*

Abelianness is also easy to check.

Proposition 3. *Given a finite set $F \subset \text{Aut}(S^{\mathbb{Z}})$, it is decidable whether $\langle F \rangle$ is abelian.*

Nevertheless, combining the result of [3] with Theorem 4 and the fundamental theorem of abelian groups we see that it is impossible to check *which* abelian group is generated by a finite set of CA.

Proposition 4. *Let G be any infinite finitely-generated abelian group. Then given a finite set $F \subset \text{Aut}(S^{\mathbb{Z}})$, it is undecidable whether $\langle F \rangle \cong G$.*

Finally, combining the result of [3] with Theorem 5, we obtain the undecidability of free products.

Theorem 9. *Given $f, g \in \text{Aut}(S^{\mathbb{Z}})$, it is undecidable whether $\langle f, g \rangle \cong F_2$, and whether f and g satisfy a nontrivial relation.*

We note that deciding whether $\langle f, g \rangle \cong F_2$ is not equivalent to (the complement of) deciding whether f and g satisfy a nontrivial relation (as the CA might satisfy a set of relations, yet generate a group isomorphic to F_2).

Proof. Let $\psi_1 : \text{Aut}(S^{\mathbb{Z}}) \rightarrow \text{Aut}(S^{\mathbb{Z}})$ and $\psi_2 : \text{Aut}(S^{\mathbb{Z}}) \rightarrow \text{Aut}(S^{\mathbb{Z}})$ be a computable embedding of $\text{Aut}(S^{\mathbb{Z}}) * \text{Aut}(S^{\mathbb{Z}})$ into $\text{Aut}(S^{\mathbb{Z}})$, given by Theorem 5. That is, there are no nontrivial relations between elements of $\psi_1(S^{\mathbb{Z}})$ and elements of $\psi_2(S^{\mathbb{Z}})$.

Now, let $h \in \text{Aut}(S^{\mathbb{Z}})$, and consider the CA $f = \psi_1(h)$ and $g = \psi_2(h)$. If h is not a torsion element, then f and g satisfy no nontrivial relations, and thus $\langle f, g \rangle \cong F_2$. If $h^n = \text{id}$ for some $n > 0$, then both f and g are torsion elements, and thus certainly satisfy a nontrivial relation: $f^n = g^n = \text{id}$ for some n . Taking minimal such n , we have $\langle f, g \rangle \cong \mathbb{Z}_n * \mathbb{Z}_n$. Unlike F_2 , this is not a torsion-free group, so $\langle f, g \rangle \not\cong F_2$. Since it is undecidable whether a given CA h is a torsion element, the claim follows. \square

Finally, we note that the undecidability of the torsion problem tells us that the set of reversible CA that generate a finite group are recursively inseparable from the set of reversible CA that generate a copy of \mathbb{Z} . The corresponding result for the two-generator free group does not seem to follow from our embedding theorem.

Question 5. *Let A be the set of pairs of CA on $S^{\mathbb{Z}}$ generating a finite group, and B the set of pairs generating F_2 . Are A and B recursively inseparable?*

9 Acknowledgements

The author was supported by FONDECYT Grant 3150552.

References

- [1] Mike Boyle, Douglas Lind, and Daniel Rudolph. The automorphism group of a shift of finite type. *Transactions of the American Mathematical Society*, 306(1):pp. 71–114, 1988.
- [2] Gustav A. Hedlund. Endomorphisms and automorphisms of the shift dynamical system. *Math. Systems Theory*, 3:320–375, 1969.
- [3] Jarkko Kari and Nicolas Ollinger. Periodicity and immortality in reversible computing. In *Proceedings of the 33rd international symposium on Mathematical Foundations of Computer Science*, MFCS '08, pages 419–430, Berlin, Heidelberg, 2008. Springer-Verlag.
- [4] K. H. Kim and F. W. Roush. On the automorphism groups of subshifts. *Pure Mathematics and Applications*, 1(4):203–230, 1990.

- [5] Bruce P. Kitchens. *Symbolic dynamics – One-sided, two-sided and countable state Markov shifts*. Universitext. Springer-Verlag, Berlin, 1998.
- [6] Douglas Lind. Entropies of automorphisms of a topological markov shift. *Proceedings of the American Mathematical Society*, 99(3):589–595, 1987.
- [7] Douglas Lind and Brian Marcus. *An introduction to symbolic dynamics and coding*. Cambridge University Press, Cambridge, 1995.
- [8] M. Lothaire. *Algebraic combinatorics on words*, volume 90 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2002.
- [9] Ville Salo. Groups and monoids of cellular automata. In Jarkko Kari, editor, *Cellular Automata and Discrete Complex Systems*, volume 9099 of *Lecture Notes in Computer Science*, pages 17–45. Springer Berlin Heidelberg, 2015.
- [10] Ville Salo and Ilkka Törmä. Computational aspects of cellular automata on countable sofic shifts. *Mathematical Foundations of Computer Science 2012*, pages 777–788, 2012.