



ELSEVIER

Contents lists available at ScienceDirect

Journal of Science and Medicine in Sport

journal homepage: www.elsevier.com/locate/jams

Journal of Science and Medicine in Sport

Corrigendum

Corrigendum to “Standing is associated with insulin sensitivity in adults with metabolic syndrome. Journal of Science and Medicine in Sport 24 (2021) 1255–1260”

Taru Garthwaite ^{a,*}, Tanja Sjöros ^a, Mikko Koivumäki ^a, Saara Laine ^a, Henri Vähä-Yppä ^b, Maria Saarenhovi ^c, Petri Kallio ^{c,d}, Eliisa Löyttyniemi ^e, Harri Sievänen ^b, Noora Houttu ^f, Kirsi Laitinen ^f, Kari Kalliokoski ^a, Tommi Vasankari ^{b,g}, Juhani Knuuti ^a, Ilkka Heinonen ^{a,h}

^a Turku PET Centre, University of Turku and Turku University Hospital, Finland

^b The UKK Institute for Health Promotion Research, Finland

^c Department of Clinical Physiology and Nuclear Medicine, University of Turku and Turku University Hospital, Finland

^d Paavo Nurmi Centre and Unit for Health and Physical Activity, University of Turku, Finland

^e Department of Biostatistics, University of Turku, Finland

^f Institute of Biomedicine, University of Turku, Finland

^g Faculty of Medicine and Health Technology, Tampere University, Finland

^h Rydberg Laboratory of Applied Sciences, University of Halmstad, Sweden

The authors regret an error in the fourth paragraph of the Methods section. This should read as follows:

Whole-body insulin-stimulated glucose uptake (M-value) was measured with a hyperinsulinaemic–euglycaemic clamp after fasting overnight. Insulin (Actrapid, 100 U/ml, Novo Nordisk, Bagsvaerd, Denmark) was infused at a 160 mU/m²/min body surface area/min rate during the first 4 min, at 80 mU/m²/min during minutes 4–7,

and thereafter at 40 mU/m²/min. Four minutes after starting insulin infusion, 20% glucose infusion was started. The rate was adjusted according to plasma glucose concentration measured every 5–10 min. M-value (μmol/kg/min) was calculated in 20-min intervals from steady-state glucose values. Fasting insulin and HOMA-IR were determined as surrogate markers of insulin sensitivity as described below.

The authors regret any inconvenience caused.

DOI of original article: <https://doi.org/10.1016/j.jsams.2021.08.009>.

* Corresponding author.

E-mail address: taru.garthwaite@utu.fi (T. Garthwaite).

<https://doi.org/10.1016/j.jsams.2022.03.002>

1440-2440/© 2022 The Author(s). Published by Elsevier Ltd on behalf of Sports Medicine Australia. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Please cite this article as: T. Garthwaite, T. Sjöros, M. Koivumäki, et al., , Journal of Science and Medicine in Sport, <https://doi.org/10.1016/j.jsams.2022.03.002>