



Vladimir Emelichev | Vadim Mychkov | Yury Nikulin

Postoptimal analysis
for one vector venturesome investment
problem

TURKU CENTRE *for* COMPUTER SCIENCE

TUCS Technical Report
No 1153, February 2016



Postoptimal analysis for one vector venturesome investment problem

Vladimir Emelichev

Belarusian State University, Mechanics and Mathematics Faculty
Nezavisimosti av. 4, 220030 Minsk, Belarus
vemelichev@gmail.com

Vadim Mychkov

Belarusian State University, Mechanics and Mathematics Faculty
Nezavisimosti av. 4, 220030 Minsk, Belarus
vadim.mychkov@gmail.com

Yury Nikulin

University of Turku, Department of Mathematics and Statistics
Vesilinnantie 5, 20014 Turku, Finland
yurnik@utu.fi

TUCS Technical Report

No 1153, February 2016

Abstract

Lower and upper bounds are obtained for the stability radius of a vector venturesome investment problem with the Hölder norms in the parameter spaces. As corollaries several results are obtained for special cases.

Keywords: vector venturesome investment problem, Pareto optimal portfolio, stability radius, perturbing matrix, Hölder's metric

TUCS Laboratory
Turku Optimization Group

1 Introduction

While solving applied optimization problems, we have to take into account different factors of uncertainty and randomness such as inadequacy of mathematical models to real processes, round-off errors, measurement errors etc. In all these cases, a mathematical problem can not be correctly posed and solved without stability theory results. In multicriteria discrete optimization problems stability research is usually tied with studying a discrete analogue of the Hausdorff continuity (semicontinuity) for dot-valued mappings i.e. mappings that assign each gang of possible data to a given set of optimal solutions [1,2].

Despite the abundance of approaches to the stability analysis in discrete optimization problems (the comprehensive idea of multiple studies on stability is provided in [3, 4]), the two mainstream approaches can be defined: qualitative and quantitative.

As a part of the qualitative approach, research is concentrated on identifying the different types of problem stability [5 – 12] or establishing interconnections between different types of stability [13, 14]. In addition, searching and describing the stability region for optimal solutions is also done (see e.g. [15]).

Big part of quantitative approach is particularised in [16], and it is tied with obtaining valid assessments of changes in the initial data, retaining some predetermined property of optimal solutions [17 – 21], and developing algorithms for calculating these bounds [22 – 26]. The key point here is stability radius, which is defined as the radius of the largest uncertain data neighbourhood preserving some optimal property in the space of perturbed problem parameters. Any perturbed problem with parameter point within the neighbourhood is «close» to the original problem

This article belongs to the second approach, and it is a continuation of publication series [27 – 34]. In that previous research, the author estimate bounds for the stability radii of solutions in vector investment Boolean problems with different criteria of optimism and pessimism (Wald, Savage criterion etc.) under a variety of combinations of linear l_1 , Chebyshev l_∞ and Hölder l_p metrics in three-dimensional parameter space. In this study, we consider the most general case where different Hölder norms are used in the parameter space mentioned above. Here we obtain lower and upper bounds for the stability radius of the vector investment problem with well-known in the theory of decision-making criteria of extreme optimism regarding portfolio returns.

The results presented here were also announced in [35].

2 Problem formulation and the basic definitions

We consider a multicriteria discrete version of optimal investment problem. Let introduce some notation:

$N_n = \{1, 2, \dots, n\}$ be various alternative investment projects (assets);

N_m be a set of possible states for the financial market (scenarios, the situations in the market);

N_s be many types of (indicators) economic efficiency of investment projects;

e_{ijk} be expected economic evaluation (profitability) of the form $k \in N_s$ for investment project $j \in N_n$ in the case, when the market is in the state $i \in N_m$;

$E = [e_{ijk}] \in \mathbf{R}^{m \times n \times s}$;

$x = (x_1, x_2, \dots, x_n)^T \in X \subseteq \mathbf{E}^n$ is the investment portfolio, where $\mathbf{E} = \{0, 1\}$,

$$x_j = \begin{cases} 1, & \text{if you select a portfolio } j \in N_n, \\ 0 & \text{otherwise;} \end{cases}$$

So here $X \subset \mathbf{E}^n$ is the set of all possible investment portfolios, i.e. which realization does not exceed the investor's initial capital;

\mathbf{R}^m is the state space of the financial market;

\mathbf{R}^n is the space of the investment projects;

\mathbf{R}^s is the space criterion of the cost efficiency.

For a given set of investment portfolios (Boolean vectors) X , $|X| \geq 2$, the vector criterion is defined

$$f(x, E) = (f_1(x, E_1), f_2(x, E_2), \dots, f_s(x, E_s)),$$

whose components are optimistic criteria (MAX-MAX) of investment portfolio returns:

$$f_k(x, E_k) = \max_{i \in N_m} e_{ik} x = \max_{i \in N_m} \sum_{j \in N_n} e_{ijk} x_j \rightarrow \max_{x \in X}, k \in N_s,$$

where $E_k \in \mathbf{R}^{m \times n}$ is the k -th cut of the matrix $E = [e_{ijk}] \in \mathbf{R}^{m \times n \times s}$ with rows $e_{ik} = (e_{i1k}, e_{i2k}, \dots, e_{ink}) \in \mathbf{R}^n$, $i \in N_m$.

With a help of such criterion a venturesome investor optimizes the efficiency of the portfolio under the assumption that the market is in the most profitable condition, namely when portfolio is in the marginal revenue position. Obviously, this approach is based on the stereotype of reckless optimism behaviour «or sink or swim», «who do not risk, nothing gained» etc.

We mean by vector s -criterion problem $Z_m^s(E)$, $s \in \mathbf{N}$, problem of finding the Pareto set $P^s(E)$, that is set of the Pareto-optimal portfolios

$$P^s(E) = \{x \in X : X(x, E) = \emptyset\},$$

where

$$X(x, E) = \{x' \in X : f(x, E) \leq f(x', E) \ \& \ f(x, E) \neq f(x', E)\}.$$

It's clear that $P^s(E) \neq \emptyset$ for any matrix $E \in \mathbf{R}^{m \times n \times s}$. Note also that the problem $Z_m^s(E)$ can be interpreted as «the best optimisation case».

Let different Hölder's norms l_p , l_q and l_r be given in the spaces \mathbf{R}^n , \mathbf{R}^m , \mathbf{R}^s respectively, where $p, q, r \in [1, \infty]$. We understand by norm of matrix $E \in \mathbf{R}^{m \times n \times s}$ the number

$$\|E\|_{pqr} = \|(\|E_1\|_{pq}, \|E_2\|_{pq}, \dots, \|E_s\|_{pq})\|_r,$$

where

$$\|E_k\|_{pq} = \|(\|e_{1k}\|_p, \|e_{2k}\|_p, \dots, \|e_{mk}\|_p)\|_q, \quad k \in N_s.$$

Recall that the Hölder norm l_p in space \mathbf{R}^n is defined by the formula

$$\|a\|_p = \begin{cases} (\sum_{j \in N_n} |a_j|^p)^{1/p}, & \text{if } 1 \leq p < \infty, \\ \max\{|a_j| : j \in N_n\}, & \text{if } p = \infty, \end{cases}$$

where $a = (a_1, a_2, \dots, a_n)^T \in \mathbf{R}^n$.

It is easy to see that the following inequalities hold

$$\|e_{ik}\|_p \leq \|E_k\|_{pq} \leq \|E\|_{pqr}, \quad i \in N_m, \quad k \in N_s \quad (1)$$

for any $p, q, r \in [1, \infty]$. Following [21, 28, 30, 32], the number

$$\rho = \rho_m^s(p, q, r) = \begin{cases} \sup \Xi, & \text{if } \Xi \neq \emptyset, \\ 0, & \text{if } \Xi = \emptyset, \end{cases}$$

is called the stability radius of the problem $Z^s(E)$, where

$$\Xi = \{\varepsilon > 0 : \forall E' \in \Omega(\varepsilon) \ (P^s(E + E') \subseteq P^s(E))\},$$

$$\Omega(\varepsilon) = \{E' \in \mathbf{R}^{m \times n \times s} : \|E'\|_{pqr} < \varepsilon\}.$$

Here $\Omega(\varepsilon)$ is a set of perturbing matrices, $P^s(E + E')$ is the Pareto set of the perturbed problem $Z^s(E + E')$, $\|E'\|_{pqr}$ is the norm of matrix $E' = \|e'_{ijk}\|$. Thus, the stability radius $\rho_m^s(p, q, r)$ for a problem $Z_m^s(E)$ is the limiting level of elements perturbation for matrix E in the space $\mathbf{R}^{m \times n \times s}$ that does not lead to the emergence of new Pareto optimal portfolios. Obviously, when $P^s(E) = X$, the stability radius of a problem should be considered infinitely large. If $P^s(E) \neq X$, we call the problem nontrivial.

3 Auxiliary assertions

Let u be one of the numbers p, q, r introduced above. With number u we associate a number u' as follows

$$1/u + 1/u' = 1, \quad 1 < u < \infty.$$

Moreover, let assume that $u' = 1$ if $u = \infty$, and $u = \infty$ if $u' = 1$. Therefore in what follows we assume that the range of the numbers u and u' is interval $[1, \infty]$, and the numbers themselves are connected by the conditions above. Additionally let's assume that $1/u = 0$ if $u = \infty$.

Next, we will use the well-known Hölder's inequality

$$|a^T b| \leq \|a\|_u \|b\|_{u'}, \quad (2)$$

that holds for any vector a and b within the same dimension.

Lemma 1. *For any portfolios $x, x^0 \in X$ and indexes $i, i' \in N_n, k \in N_s$, the following inequality holds*

$$e_{i'k}x^0 - e_{ik}x \geq -\|E_k\|_{pq} \|(\|x^0\|_{p'}, \|x\|_{p'})\|_v, \quad (3)$$

where

$$v = \min\{p', q'\}.$$

Indeed, if $i \neq i'$, then using Hölder's inequality (2) we have

$$\begin{aligned} e_{i'k}x^0 - e_{ik}x &\geq -(\|e_{i'k}\|_p \|x^0\|_{p'} + \|e_{ik}\|_p \|x\|_{p'}) \geq \\ &\geq -\|(\|e_{i'k}\|_p, \|e_{ik}\|_p)\|_q \|(\|x^0\|_{p'}, \|x\|_{p'})\|_{q'} \geq \\ &\geq -\|E_k\|_{pq} \|(\|x^0\|_{p'}, \|x\|_{p'})\|_{q'} \geq -\|E_k\|_{pq} \|(\|x^0\|_{p'}, \|x\|_{p'})\|_v. \end{aligned}$$

If $i = i'$, then using (1) and Hölder's inequality (2) it comes out that

$$\begin{aligned} e_{i'k}x^0 - e_{ik}x &\geq -\|e_{ik}\|_p \|x^0 - x\|_{p'} \geq -\|E_k\|_{pq} \|x^0 - x\|_{p'} \geq \\ &-\|E_k\|_{pq} \|(\|x^0\|_{p'}, \|x\|_{p'})\|_{p'} \geq -\|E_k\|_{pq} \|(\|x^0\|_{p'}, \|x\|_{p'})\|_v. \end{aligned}$$

Also it is easy to see that for the vector $a = (a_1, a_2, \dots, a_n)^T \in \mathbf{R}^n$ with conditions $|a_j| = \alpha, j \in N_n$ the equality

$$\|a\|_p = \alpha n^{1/p} \quad (4)$$

holds for any number $p \in [1, \infty]$.

4 Bounds for the stability radius

For the non-trivial problem $Z_m^s(E)$, we assume

$$\begin{aligned}\varphi &= \min_{x \notin P^s(E)} \max_{x' \in P(x, E)} \frac{\gamma(x', x)}{\|(\|x'\|_{p'}, \|x\|_{p'})\|_v}, \\ \psi &= \min_{x \notin P^s(E)} \max_{x' \in P(x, E)} \frac{\gamma(x', x)}{\|x' - x\|_1}, \\ \gamma(x', x) &= \min\{f_k(x', E_k) - f_k(x, E_k) : k \in N_s\},\end{aligned}$$

$$P(x, E) = X(x, E) \cap P^s(E),$$

$$v = \min\{p', q'\},$$

$$\sigma = \min\{\|E_k\|_{pq} : k \in N_s\}.$$

It is easy to see that $\varphi, \psi \geq 0$.

Theorem 1. *For the stability radius $\rho_m^s(p, q, r)$ of non-trivial problem $Z_m^s(E)$, $s \geq 1$, the following estimates are true*

$$\varphi \leq \rho_m^s(p, q, r) \leq \min\{n^{1/p} m^{1/q} s^{1/r} \psi, \sigma\}, \quad (5)$$

for any $s, m \in \mathbf{N}$ and $p, q, r \in [1, \infty]$

Proof. At first we show that $\rho \geq \varphi$. With $\varphi = 0$ this inequality is obvious. Let $\varphi > 0$. We assume that the perturbing matrix $E' = \|e'_{ijk}\| \in \mathbf{R}^{m \times n \times s}$ with cut E'_k , $k \in N_s$, belongs to $\Omega(\varphi)$. According to the definition φ , for any portfolio $x \notin P^s(E)$ there exists portfolio $x^0 \in P(x, E)$ such that

$$\gamma(x^0, x) \geq \varphi \|(\|x^0\|_{p'}, \|x\|_{p'})\|_v, \quad k \in N_s.$$

Hence, taking into account inequality (1) and (3), for any index $k \in N_s$ we deduce

$$\begin{aligned}f_k(x^0, E_k + E'_k) - f_k(x, E_k + E'_k) &= \max_{i \in N_m} (e_{ik} + e'_{ik})x^0 - \max_{i \in N_m} (e_{ik} + e'_{ik})x = \\ &= \min_{i \in N_m} \max_{i' \in N_m} (e_{i'k}x^0 + e'_{i'k}x^0 - e_{ik}x - e'_{ik}x) \geq \\ &\geq f_k(x^0, E_k) - f_k(x, E_k) - \|E'\|_{pqr} \|(\|x^0\|_{p'}, \|x\|_{p'})\|_v \geq \\ &\geq (\varphi - \|E'\|_{pqr}) \|(\|x^0\|_{p'}, \|x\|_{p'})\|_v > 0,\end{aligned}$$

where e'_{ik} is the i -th row of the cut E'_k of matrix E' . Thus, portfolio x doesn't belong to the Pareto set $P^s(E + E')$. Therefore, we conclude that the following inclusion is correct

$$P^s(E + E') \subseteq P^s(E)$$

for any perturbing matrix $E' \in \Omega(\varphi)$.

Therefore, $\rho \geq \varphi$.

Next we prove the inequality $\rho \leq m^{1/p}n^{1/q}s^{1/r}\psi$.

According to the definition of the number ψ , there is a portfolio $x^0 \notin P^s(E)$, such that for any portfolio $x \in P(x^0, E)$ there exists index $l \in N_s$, whereby

$$f_l(x, E_l) - f_l(x^0, E_l) \leq \psi \|x - x^0\|_1. \quad (6)$$

Assuming

$$\varepsilon > n^{1/p}m^{1/q}s^{1/r}\psi,$$

we define elements e_{ijk}^0 for any k -th cut E_0^k , $k \in N_s$, of the perturbed matrix $E^0 = \|e_{ijk}^0\| \in \mathbf{R}^{m \times n \times s}$ according to the rule

$$e_{ijk}^0 = \begin{cases} \delta, & \text{if } i \in N_m, x_j^0 = 1, \\ -\delta, & \text{if } i \in N_m, x_j^0 = 0, \end{cases}$$

where

$$\psi < \delta < \varepsilon/n^{1/p}m^{1/q}s^{1/r}. \quad (7)$$

Hence, according to (4), we obtain

$$\|e_{ik}^0\|_p = \delta n^{1/p}, \quad i \in N_m, \quad k \in N_s,$$

$$\|E_k^0\|_{pq} = \delta n^{1/p}m^{1/q}, \quad k \in N_s,$$

$$\|E^0\|_{pqr} = \delta n^{1/p}m^{1/q}s^{1/r}.$$

It means that $E^0 \in \Omega(\varepsilon)$. In addition, all rows e_{ik}^0 , $i \in N_m$, $k \in N_s$, are identical and they consist of component δ and $-\delta$. Therefore, if we set $c = e_{ik}^0$, $i \in N_m$, $k \in N_s$, we deduce that the correlation

$$c(x - x^0) = -\delta \|x - x^0\|_1 < 0, \quad (8)$$

is valid for any portfolio $x \neq x^0$.

Hence, taking into account (6) and (7), we conclude that for any portfolio $x \in P(x^0, E)$ there exists an index $l \in N_s$ such that

$$\begin{aligned} f_l(x, E_l + E_l^0) - f_l(x^0, E_l + E_l^0) &= \max_{i \in N_m} (e_{il} + e_{il}^0)x - \max_{i \in N_m} (e_{il} + e_{il}^0)x^0 = \\ &= \min_{i \in N_m} \max_{i' \in N_m} (e_{i'l}x - e_{il}x^0 + e_{i'l}^0x - e_{il}^0x^0) = \\ &= f_l(x, E_l) - f_l(x^0, E_l) + c(x - x^0) \leq (\psi - \delta) \|x - x^0\|_1 < 0. \end{aligned}$$

Thus, the following formula holds

$$\forall x \in P(x^0, E) \quad (x \notin X(x^0, E + E^0)). \quad (9)$$

If $X(x^0, E + E^0) = \emptyset$, then it is obvious that portfolio x^0 (it is not Pareto optimal portfolio for problem $Z_m^s(E)$) appears to be optimal in the perturbed problem $Z_m^s(E + E^0)$, i.e. $x^0 \in P^s(E + E^0)$.

If $X(x^0, E + E^0) \neq \emptyset$, then due to the external stability of the Pareto set $P^s(E + E^0)$ (look, for example, [37] p. 34) we can find portfolio $x^* \in P^s(x^0, E + E^0)$. Let us show that $x^* \notin P^s(E)$.

Let assume, on the contrary, that $x^* \in P^s(E)$. Then according to (9), $x^* \notin P^s(x^0, E)$. This leads to two cases.

At first, let have a look at the case $f(x^*, E) = f(x^0, E)$. Then for any index $k \in N_s$ according to (8), we have

$$\begin{aligned} f_k(x^*, E_k + E_k^0) - f_k(x^0, E_k + E_k^0) &= f_k(x^*, E_k) - f_k(x^0, E_k) + c(x^* - x^0) = \\ &= -\delta \|x^* - x^0\|_1 < 0. \end{aligned}$$

Next, let's examine the case where exists an index $h \in N_s$, that

$$f_h(x^*, E_h) < f_h(x^0, E_h).$$

Then, again using (8), we arrive at correlation

$$f_h(x^*, E_h + E_h^0) - f_h(x^0, E_h + E_h^0) = f_h(x^*, E_h) - f_h(x^0, E_h) + c(x^* - x^0) < 0.$$

As a result both cases contradict to the inclusion $x^* \in P^s(x^0, E + E^0)$. This proves that $x^* \notin P^s(E)$. Let us remind that $x^* \in P^s(E + E^0)$.

To summarize, we come to a conclusion that for any number $\varepsilon > n^{1/p}m^{1/q}s^{1/r}\psi$ the existence of the perturbing matrix is guaranteed $E^0 \in \Omega(\varepsilon)$, and there is a portfolio (x^0 or x^*), which is not Pareto optimal portfolio problem $Z_m^s(E)$, which is simultaneously the same in perturbed problem $Z_m^s(E + E^0)$. Thus, the following formula holds

$$\forall \varepsilon > n^{1/p}m^{1/q}s^{1/r}\psi \quad \exists E^0 \in \Omega(\varepsilon) \quad (P^s(E + E^0)) \not\subseteq P^s(E).$$

Therefore, $\rho \leq n^{1/p}m^{1/q}s^{1/r}\psi$.

Finally, we prove inequality

$$\rho_m^s(p, q, r) \leq \sigma.$$

Let x^0 be any non-Pareto optimal portfolio for problem $Z_m^s(E)$, i.e. $x^0 \notin P^s(C)$. Let index $l \in N_m$ be such that

$$\sigma = \|E_l\|_{pq}. \quad (10)$$

Assuming $\varepsilon > \sigma$, let's choose a number δ with condition

$$0 < \delta n^{1/p}m^{1/q} < \varepsilon - \sigma. \quad (11)$$

Next, consider the matrix $V = [v_{ij}] \in \mathbf{R}^{m \times n}$ with elements

$$v_{ij} = \begin{cases} \delta, & \text{if } i \in N_m, x_j^0 = 1, \\ -\delta, & \text{if } i \in N_m, x_j^0 = 0. \end{cases}$$

Then, using (4) we obtain

$$\|V\|_{pq} = \delta n^{1/p} m^{1/q}. \quad (12)$$

Moreover, all the rows of matrix V are the same, and they consist of numbers δ and $-\delta$. Having denoted the line as A , we have

$$A(x - x^0) = -\delta \|x - x^0\|_1 < 0 \quad (13)$$

for any portfolio $x \in X \setminus \{x^0\}$. Cuts E_k^0 , $k \in N_s$ perturbing matrix $E^0 \in \mathbf{R}^{m \times n \times s}$ let us define according to the rule

$$E_k^0 = \begin{cases} V - E_k, & \text{if } k = l, \\ \mathbf{0}, & \text{if } k \neq l, \end{cases}$$

where $\mathbf{0} \in \mathbf{R}^{m \times n}$. Then referring to (10) – (12) we find

$$\|E^0\|_{pqr} = \|E_l^0\|_{pq} = \|V - E_l\|_{pq} \leq \|V\|_{pq} + \|E_l\|_{pq} \leq \delta n^{1/p} m^{1/q} + \sigma < \varepsilon,$$

and taking into consideration the inequality (13), we obtain

$$\begin{aligned} f_l(x, E_l + E_l^0) - f_l(x^0, E_l + E_l^0) &= f_l(x, V) - f_l(x^0, V) = \\ &= A(x - x^0) = -\delta \|x - x^0\|_1 < 0. \end{aligned}$$

It means that $x \notin X(x^0, E + E^0)$, where $E^0 \in \Omega(\varepsilon)$. But as $x^0 \notin X(x^0, E + E^0)$, then $X(x^0, E + E^0) = \emptyset$. Therefore

$$x^0 \in P_m^s(E + E^0).$$

Taking into account $x^0 \notin P^s(E)$, we conclude that $\rho_m^s(p, q, r) \leq \varepsilon$ for any number $\varepsilon > \sigma$. Therefore, $\rho_m^s(p, q, r) \leq \sigma$, together with previously proven inequality

$$\rho_m^s(p, q, r) \leq n^{1/p} m^{1/q} s^{1/r} \psi$$

it provides wanted upper estimate of the stability radius. Theorem 1 is proved.

5 Corollaries

Corollary 1. For $s, m \in \mathbf{N}$ and $p \in [1, \infty]$ the inequalities hold

$$\begin{aligned} \min_{x \notin P^s(E)} \max_{x' \in P(x, E)} \frac{\gamma(x, x')}{\|x' + x\|_1^{1/p'}} &\leq \rho_m^s(p, p, p) \leq \\ &\leq (nms)^{1/p} \min_{x \notin P^s(E)} \max_{x' \in P(x, E)} \frac{\gamma(x, x')}{\|x' - x\|_1}. \end{aligned}$$

Hence, we obtain the following obvious statement indicating the assessment reachability

Corollary 2. If for any two portfolios $x \notin P^s(E)$ and $x' \in P(x, E)$ of the problem $Z_m^s(E)$ the set

$$\{j \in N_n : x_j = x'_j = 1\}$$

is empty, then the following formula holds

$$\rho_m^s(\infty, \infty, \infty) = \varphi = \psi.$$

Corollary 3[32]. For $s, m \in \mathbf{N}$ and $p \in [1, \infty]$, the inequalities hold

$$\begin{aligned} \min_{x \notin P^s(E)} \max_{x' \in P(x, E)} \frac{\gamma(x', x)}{\|x'\|_{p'} + \|x\|_{p'}} &\leq \rho_m^s(p, \infty, p) \leq \\ &\leq (ns)^{1/p} \min_{x \notin P^s(E)} \max_{x' \in P(x, E)} \frac{\gamma(x', x)}{\|x' - x\|_1}. \end{aligned}$$

Corollary 4[33]. For $s, m \in \mathbf{N}$ and $p \in [1, \infty]$, the inequalities hold

$$\begin{aligned} \min_{x \notin P^s(E)} \max_{x' \in P(x, E)} \frac{\gamma(x', x)}{\|x' + x\|_1} &\leq \rho_m^s(\infty, p, p) \leq \\ &\leq (ms)^{1/p} \min_{x \notin P^s(E)} \max_{x' \in P(x, E)} \frac{\gamma(x', x)}{\|x' - x\|_1}. \end{aligned}$$

Corollary 5[34]. For $s, m \in \mathbf{N}$ and $p \in [1, \infty]$, the inequalities hold

$$\begin{aligned} \min_{x \notin P^s(E)} \max_{x' \in P(x, E)} \frac{\gamma(x', x)}{\|x'\|_{p'} + \|x\|_{p'}} &\leq \rho_m^s(p, \infty, \infty) \leq \\ &\leq n^{1/p} \min_{x \notin P^s(E)} \max_{x' \in P(x, E)} \frac{\gamma(x', x)}{\|x' - x\|_1}. \end{aligned}$$

6 The case of linear criteria

With $m = 1$ the investment problem is transformed into vector problem of linear Boolean programming. Let it be written in a convenient form

$$Z_1^s(E) : e_k x \rightarrow \max_{x \in X}, \quad k \in N_s.$$

where $X \subset \mathbf{E}^n$, $e_k \in \mathbf{R}^n$ is the k -th row of the matrix $E = [e_{kj}] \in \mathbf{R}^{s \times n}$. The very case can be interpreted as a situation in which the state of the financial market doesn't raise doubts. As before, let us consider that in the space of projects \mathbf{R}^n and a criterion space \mathbf{R}^s set respectively the Hölder norm l_p и l_r , where $p, r \in [1, \infty]$. For the problem $Z_1^s(E)$, the same notation $P^s(E)$, $P(x, E)$ and etc will be used.

In this linear case, a lower bound for the stability radius ρ_1^s of a problem $Z_1^s(E)$ can be improved. Indeed, the following theorem holds.

Theorem 2. *For the stability radius $\rho_1^s(p, r)$ of non-trivial problem $Z_1^s(E)$, the following bounds are valid*

$$\xi(p) \leq \rho_1^s(p, r) \leq n^{1/p} s^{1/r} \xi(\infty), \quad (14)$$

where

$$\xi(p) = \min_{x \notin P^s(E)} \max_{x' \in P(x, E)} \min_{k \in N_s} \frac{e_k(x' - x)}{\|x' - x\|_p} \quad (15)$$

for any $s \in \mathbf{N}$ and $p, r \in [1, \infty]$.

Proof. The validity of the upper bound, i.e. the inequality

$$\rho_1^s(p, r) \leq n^{1/p} s^{1/r} \xi(\infty),$$

follows from Theorem 1. Next we prove the inequality

$$\rho_1^s(p, r) \geq \xi(p). \quad (16)$$

With $\xi(p) = 0$, it is obvious. Let $\xi(p) > 0$ and $E' \in \mathbf{R}^{s \times n}$ be the perturbing matrix with the rows $e'_k \in \mathbf{R}^n$, $k \in N_s$, and norm

$$\|E'\|_{pr} = \|(\|e'_1\|_p, \|e'_2\|_p, \dots, \|e'_s\|_p)\|_r < \xi(p).$$

According to (15) for every portfolio $x \notin P^s(E)$ there is a Pareto-optimal portfolio $x^0 \in P^s(E)$ such that

$$\|e'_k\|_p \leq \|E'\|_{pr} < \xi(p) \leq \frac{e_k(x' - x)}{\|x' - x\|_p}, \quad k \in N_s.$$

Therefore, by Hölder's inequality (2) for any index $k \in N_s$ we have

$$(e_k + e'_k)(x^0 - x) = e_k(x^0 - x) + e'_k(x^0 - x) \geq e_k(x^0 - x) - \|e'_k\|_p \|x^0 - x\|_{p'} > 0,$$

i.e. $x \notin P^s(E + E')$. Hence, we come to a conclusion that $P^s(E + E') \subseteq P^s(E)$ under any perturbing matrix $E' \in \mathbf{R}^{s \times n}$ with the norm $\|E'\|_{pr} < \xi(p)$. Consequently, inequality (16) holds. Theorem 2 is proved.

Due to Theorem 2, we arrive at the known result, indicating the reachability of lower and upper bounds (14) for the stability radius in the problem $Z_1^s(E)$ with $p = r = \infty$.

Corollary 6[38]. *Given $s \in \mathbf{N}$, for the stability radius of non-trivial problem $Z_1^s(E)$, the next equality holds*

$$\rho_1^s(\infty, \infty) = \min_{x \notin P^s(E)} \max_{x' \in P(x, E)} \min_{k \in N_s} \frac{e_k(x' - x)}{\|x' - x\|_1}.$$

7 Conclusion

We considered a multicriteria discrete version of optimal investment problem. Here venturesome investor optimizes the efficiency of the portfolio under the assumption that the market is in revenue position. We obtained the lower and upper attainable bounds for the stability radius of a vector Boolean investment problem of portfolio optimization with the Hölder norms in the parameter spaces. Conducting similar kind of research for vector investment problems with non-Boolean decision variables could be an interesting direction for further investigations.

References

- [1] Miettinen K. Nonlinear Multiobjective Optimization. Boston, 1999.
- [2] Sergienko I.V., Shylo V.P. Discrete Optimization Problems. Naukova Dumka. Kiev. 2003. (in Russian).
- [3] Greenberg H.J. An annotated bibliography for post-solution analysis in mixed integer programming and combinatorial optimization // Advances in combinatorial and stochastic optimization. Logic programming and heuristic search. Boston. 1998. P. 97–148.
- [4] Ehrgott M., Gandibleux X. A survey and annotated bibliography of multiobjective combinatorial optimization // Operation Research Spectrum. 2000. V. 22. No. 4. P. 425–460.
- [5] Emelichev V.A., Kravtsov M.K., Podkopaev D.P. On the quasistability of trajectory problems of vector optimization // Mathematical notes. Vol 63. Issue 1. P. 19–24.
- [6] Emelichev V.A., Kuzmin K.G., Leonovich A.M. Stability in the Combinatorial Vector Optimization Problems // Automation and Remote Control. 2004. Vol 65. Issue 2. P. 227–240
- [7] Emelichev V.A., Kuzmin K.G. About quasistability of vector Boolean problem of minimizing linear functions modules // Proceedings of the National Academy of Sciences of Belarus (Vesti NAN Belarusi). Ser. fiz.-mat. nauk. 2005. No. 4. P. 55–58. (in Russian).
- [8] Gurevsky E.E. Emelichev V.A., Kuzmin K.G. About two types of stability for lexicographic problem of linear function modules minimization // Cybernetics and Systems Analysis. 2007. No. 4. P. 127–132.
- [9] Emelichev V.A., Kuzmin K.G. Stability criteria in vector combinatorial bottleneck problems in terms of binary relations // Cybernetics and Systems Analysis. 2008. V. 44, No. 3. P. 397–404.
- [10] Emelichev V.A., Kuzmin K.G. Vodennikov A.G. About one type of radius stability of vector combinatorial location problem // Discrete analysis and operations research (Diskretnyi analiz i issledovanie operaciy). Ser. 2. 2007. T. 14. No. 2. P. 32–40. (in Russian).
- [11] Emelichev V.A., Gurevsky E.E., Kuzmin K.G. On stability of some lexicographic integer optimization problem // Control and Cybernetics. 2010. V. 39. No. 3. P. 35–44.

- [12] Emelichev V.A., Karelkina O.V., Kuzmin K.G. Qualitative stability analysis of multicriteria combinatorial minimin problem // Control and Cybernetics. 2012. V. 41. No. 1. P. 57–79.
- [13] Lebedeva T.T., Semenova N.V., Sergienko T.I. Stability of vector problem of integer optimization: Relationship with the stability of sets of optimal and nonoptimal solutions // Cybernetics and System Analysis. 2005. 41 (4). P. 551–558.
- [14] Lebedeva T.T., Sergienko T.I. Different types of stability of vector integer optimization problem: general approach // Cybernetics and Systems Analysis. 2008. 44 (3). P. 429–433.
- [15] Libura M., van der Pourt E.S., Sierksma G., van der Veen J.A.A. Stability aspects of the traveling salesman problem based on k-best solutions // Discrete Applied Mathematics. 1998. V. 87. P. 159–185.
- [16] Sotskov Yu.N., Sotskova N.Yu Scheduling Under Uncertainty. Theory and Algorithms. Minsk. 2004.
- [17] Emelichev V.A., Kravtsov M.K. Stability in vector optimization path problems // Cybernetics and Systems Analysis. 1995. 31 (4). P. 599–604.
- [18] Sotskov Yu.N., Leontev V.K., Gordeev E.N. Some concepts of stability analysis in combinatorial optimization // Discrete Applied Mathematics. 1995. V. 58. No. 2. P. 169–190.
- [19] Emelichev V.A., Girlich E., Nikulin Yu.V., Podkopaev D.P. Stability and regularization of vector problem of integer linear programming // Optimization. 2002. V. 51. No.4. P. 645–676.
- [20] Emelichev V.A., Kuzmin K.G. A general approach to studying the stability of a Pareto optimal solution of a vector integer linear programming problem. Discret. Math. Appl. 2007. 17. No. 4. P. 349–354.
- [21] Emelichev V., Podkopaev D. Quantitative stability analysis for vector problems of 0-1 programming // Discrete Optimization. 2010. V. 7. No. 1–2. P. 48–63.
- [22] Gordeev E.N. Algorithms of polynomial complexity for computing the radius of instability in two classes of trajectory problems // USSR Computational Mathematics and Mathematical Physics. 1987. Vol 27. Issue 4. P. 14–20.
- [23] Chakravarti N., Wagelmans A. Calculation of stability radius for combinatorial optimization problems // Operations Research Letters. 1998. V. 23. No. 1. P. 1–7.

- [24] Gordeev E.N. Stability analysis of the minimum spanning tree problem in the l_1 metric // Computational Mathematics and Mathematical Physics. 1999. 39:5. P. 738–746
- [25] Gordeev E.N. Research of stability optimisation problems on matroids in the metric l_1 // Cybernetics and Systems Analysis. 2001. No. 2. P. 132–144.
- [26] Roland J., Smet Y., Figueira J. On the calculation of stability radius for multi-objective combinatorial optimization problems by inverse optimization // 4or Journal Operations Research. 2012. V. 10. No. 4. P. 379–389.
- [27] Emelichev V.A., Korotkov V.V., Kuzmin K.G. Multicriterial investment problem in conditions of uncertainty and risk // J. of Computer and System Sciences International. 2011. 50. No. 6. P. 1011–1018.
- [28] Emelichev V.A., Korotkov V.V. Stability radius of a vector investment problem with Savage’s minimax risk criteria // Cybernetics and Systems Analysis. 2012. V. 48, No. 3. P. 378–386.
- [29] Emelichev V., Korotkov V. Post-optimal analysis of investment problem with Wald’s ordered maximin criteria // Bulletin of the Academy of Sciences of Moldova. Mathematics. 2012. No. 1 (68). P. 59–69.
- [30] Emelichev V.A., Korotkov V.V. On stability of a vector Boolean investment problem with Wald’s criteria // Discrete Math. Appl. 2012. V. 22, No. 4. P. 367–381.
- [31] Emelichev V.A., Korotkov V.V. Postoptimal Analysis of Bicriteria Boolean Problems of Selecting Investment Projects with Wald’s and Savage’s Criteria // Journal of Computer and Systems Sciences International. 2013. V. 52, No. 4. P. 608–617.
- [32] Bukhtoyarov S.E., Emelichev V.A. On the stability measure of solutions to a vector version of an investment problem // Journal of Applied and Industrial Mathematics. 2015. V. 9, No. 3. P. 328–334.
- [33] Emelichev V.A., Ustilko E.V. Postoptimal analysis of multicriteria investment problem with the extreme optimism criteria // Applied Discrete Mathematics (Prikladnaya Diskretnaya Matematika). 2014. P. 117–123. (in Russian).
- [34] Bukhtoyarov S.E., Emelichev V.A. Stability of Markovitz investment problem with extreme optimism criteria // Proceedings of the National Academy of Sciences of Belarus (Vesti NAN Belarusi). Ser. fiz.-mat. navuk. 2014. No. 3. P. 44–48. (in Russian).

- [35] Emelichev V.A., Mychkov V.I. About stability radius for multicriterial investment problem with Hölder's norms in the parameter space // Discrete Mathematics, Algebra and their Applications: abstracts of the International Scientific Conference. The conference is dedicated to the 100th anniversary of Dmitrii Alexeevich Suprunenko. (Diskretnaya matematika, algebra i ih prilozheniya: tezisyy docladov mezghdunarodnoi nauchnoi konferencii, posviaschennoi 100-letiyu Suprunenko D.A.) (Minsk. 14-18 September 2015) / Institute of mathematics of national academy of sciences of Belarus. Minsk. 2015. P. 101–103. (in Russian).
- [36] Markowitz H.M. Portfolio Selection: Efficient Diversification of Investments. Oxford, 1991.
- [37] Podinovski V.V., Nogin V.D. Pareto-optimal solutions of multicriteria problems. Moskow. 2007 (in Russian).
- [38] Emelichev V.A., Podkopaev D.P. Stability and regularization of vector problems of integer linear programming // Discrete analysis and operations research (Diskretnyi Analiz i Issledovanie Operatsiy). 2001. 8. No. 1. 47–69. (in Russian)

TURKU
CENTRE *for*
COMPUTER
SCIENCE


Joukahaisenkatu 3-5 A, 20520 TURKU, Finland | www.tucs.fi



University of Turku

Faculty of Mathematics and Natural Sciences

- Department of Information Technology
- Department of Mathematics and Statistics
- Turku School of Economics*
- Institute of Information Systems Sciences



Åbo Akademi University

- Computer Science
- Computer Engineering

ISBN 978-952-12-3349-4

ISSN 1239-1891