

# Functional Consequences of Road Traffic Injuries: Preliminary Results from PERSIAN Traffic Cohort (PTC)

Yalda Mousazadeh<sup>1,2</sup>, Homayoun Sadeghi-Bazargani<sup>3\*</sup>, Ali Janati<sup>4</sup>, Hossein Poustchi<sup>5</sup>, Raana Zakeri<sup>3</sup>, Alireza Shafiee-Kandjani<sup>3</sup>, Faramarz Pourasghar<sup>6</sup>, Nasrin Shahedifar<sup>3,7</sup>, Zeynab Iraj<sup>8</sup>, Mahdi Rezaei<sup>3</sup>, Shahrzad Bazargan-Hejazi<sup>9</sup>

1 Department of Public Health, Khalkhal University of Medical Sciences, Khalkhal, Iran.

2 Department of Health Policy and Management, Iranian Center of Excellence in Health Management, School of Management and Medical Informatics, Tabriz University of Medical Sciences, Tabriz, Iran.

3 Road Traffic Injury Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

4 Iranian Center of Excellence in Health Management, School of Management and Medical Informatics, Tabriz University of Medical Sciences, Tabriz, Iran.

5 Liver and Pancreatobiliary Diseases Research Center, Digestive Disease Research Institute, Shariati Hospital, Tehran university of Medical Sciences, Tehran, Iran.

6 Department of Medical Informatics, School of Management and Medical Informatics, Tabriz University of Medical Sciences, Tabriz, Iran.

7 Injury Epidemiology and Prevention Research Group, Turku Brain Injury Center, Division of Clinical Neurosciences, Turku University Hospital and University of Turku, Turku, Finland.

8 Department of Statistics and Epidemiology, Faculty of Health, Tabriz University of Medical Sciences, Tabriz, Iran.

9 Department Psychiatry and Human Behavior, Charles R. Drew University of Medicine and Science & David Geffen of Medicine at University of California at Los Angeles, Los Angeles, CA, USA.

\*Corresponding Author: Prof. Homayoun Sadeghi-Bazargani: Road Traffic Injury Research Center, Tabriz University of Medical Sciences, Tabriz, Iran. Phone: 04133800568, Fax: 04133800538, Email: homayoun.sadeghi@gmail.com.

Received 2021-06-05; Accepted 2021-08-05; Online Published 2021-11-23

## Abstract

**Introduction:** Traffic injuries are one of the main causes of death worldwide. After decreasing mortality rates and improving the recovery of injured patients, long-term functional consequences need to be addressed. The purpose of this study was to assess the functional outcomes of road traffic injuries and their predictors six months after hospital discharge, based on the preliminary results from PTC.

**Methods:** A cross-sectional study based on PERSIAN Traffic Health and Safety Cohort Study was performed. Data were collected using the 12-item World Health Organization Disability Assessment Schedule 2.0 (WHODAS 2.0) with six domains (cognition, mobility, self-care, getting along with others, life activities, and participation), filled-in by 180 injured adults (age >18 years) at six-month follow-up after hospital discharge during October 2019. These patients were hospitalized after road accidents at two referral trauma centers, (from 23 September 2018 to 20 March 2019).

**Results:** The majority of participants were men (82.7%), (Mean age =38.8). The mean score of WHODAS 2.0 was 17.8 (SD=9.1). The highest score was estimated for the self-care dimension 3.3 (SD=1.8), and the lowest score for getting along with others 2.4 (SD=1.2). Age, gender, physiotherapy, injury localization including head and face, spinal cord, and upper extremity were predictors of WHODAS 2.0 score in various dimensions ( $p<0.05$ ).

**Conclusion:** The current study identified some functional disabilities among patients sustaining road traffic injuries. It is evident from the results that a proportion of patients do not recover six months after the injury and suffer a disability, especially in self-care, mobility, and life activities, which potentially prevent them from returning to normalcy. In addition, age, gender, physiotherapy, injury localization was related to WHODAS 2.0 scores

**Keywords:** Accidents, Road Traffic Injury, Traffic Accidents, Functional Status, WHODAS 2.0, Cohort Studies.

## Introduction

Severe injuries are one of the causes of death in the first four decades of life,<sup>1</sup> and more than five million die out of injuries annually. Injuries affect 700 million people all over the world.<sup>2</sup> Multiple trauma is usually caused by a road crash, which is a leading and increasing public health problem in many countries.<sup>3</sup> In Iran, over 21000 Disability-Adjusted Life Years (DALYs) were estimated due to all diseases and injuries, of which 28% were attributed to trauma.<sup>4</sup> Road Traffic Injuries (RTIs) account for about 2600 person-years of life lost<sup>5</sup> that cost Iran about 7.2 billion US Dollars (2.19% of Iran's Gross domestic product (GDP) in 2010).<sup>6</sup>

A considerable number of injured people suffer from long-term physically and mentally impairments, disabilities, and handicaps<sup>3</sup>, differing by their types, the number of injuries, and body location.<sup>7</sup> According to the World Health Organization (WHO) definition, disabilities is an umbrella term, covering impairments, activity restrictions, and participation restrictions.<sup>4</sup> Low and Middle Income Countries (LMICs) bear 90% of trauma-related deaths and related disability-adjusted life years, which increase progressively and reduce national income levels.<sup>8</sup> For assessing functional consequences and disabilities, a trauma system monitors the quality of trauma care. Arising the awareness of the functional consequences would improve the probability of recovery from major trauma, and have recommendations for the improvement of rehabilitation services.<sup>9</sup> In addition, it is a clinical and economic obligation.<sup>3</sup>

A wide variety of tools are available for the assessment of post-injury levels of functioning and disability.<sup>10</sup> The 12-item version of the World Health Organization Disability Assessment Schedule 2.0 (WHODAS 2.0) with a simple sum scoring method is one of the reliable tools developed by the WHO regarding the Diagnostic and Statistical Manual of Mental Disorders (Fifth Edition). According to the International Classification of Functioning, Disability, and Health, WHODAS measures disability and

functional impairments in six domains: cognition, mobility, self-care, getting along with others, life activities, and participation.<sup>11</sup> The WHODAS 2.0, like other generic disability measures was developed to consider problems due to health circumstances include diseases, illnesses, injuries, mental or emotional difficulties, and problems with alcohol or drugs.<sup>12</sup> Regarding the short completion time of the short-form schedule (about five minutes) through either phone or clinical interviewing<sup>13</sup>, it explained that 81% of the variance of the full version, the 12-item WHODAS 2.0 was preferred over its 36-item version<sup>13-14</sup>, in the current study.

The demographic factors (age and gender), and injury-related factors (localization, severity, number of injuries) are major determinants of functional outcome in some studies.<sup>3</sup> Several studies have drawn attention to the influence of psychological factors, such as depression and posttraumatic stress disorder.<sup>15-16</sup> Despite several studies on psychological factors, the association between functional outcomes and demography, physical health, and hospitalization-related factors in trauma patients has received little attention.<sup>17</sup> On the other hand, road traffic injuries are one of the most significant traumatic patients. These patients face many physical and physiological problems. The main objective of the present study was to explore the functional status of patients with traffic injuries, some related hospitalization, and demographic factors.

## Methods

### Design

The current cross-sectional study is a preliminary result of a population-based PERSIAN Traffic Safety and Health Cohort (PTC) study on exposures, predictors, and outcomes related to post-crash, started in 2019.<sup>18</sup>

### Patients and settings

The study was performed in the metropolitan city of Tabriz in the Province of East

Azerbaijan, Iran, where RTIs are among the ten prominent burdens of diseases.<sup>19</sup> The target population included all patients hospitalized after a road crash in two referral trauma centers, Shohada and Imam Reza hospitals, Tabriz, Iran, from 2018 to 2019 (from 23 September 2018 to 20 March 2019). The study data was about post-injury functioning of survivors, which was collected six months after hospital discharge, during October 2019. The sampling method was convenience sampling. Inclusion criteria included enrolment at baseline, a minimum of 18 years old, provision of informed consent, registration in road traffic registry system in two referral trauma centers, and participation in the second follow-up of a post-crash cohort study. Exclusion criteria consisted of being out of reach due to the wrong phone number for contact, immigration, or refusal. Based on the mentioned criteria, a total of 180 patients were included in the analysis.

### Measurement

The post-injury functioning of RTI survivors was examined six months after hospital discharge using the 12-item WHODAS2.0. Based on information gathered at baseline and first follow-up, people eligible to be at the second follow-up (FU2) were chosen and contacted by phone. At FU2, several questionnaires were administered include demographic and economic questions, Post Traumatic Stress Disorder (PTSD), Patient Health Questionnaire-9 (PHQ9), Europe Quality of Life-5 Dimensions-3 Levels (EQ-5D-3L), and WHODAS 12-item tools. This study-utilized data contributed to WHODAS 12-item. Via telephone interview, people were asked about their post-injury functional status as mentioned in the short form. According to WHO simple scoring method, items are responded on a five-point Likert-type scale from 0 to 4, as follows, 0: No Difficulty; 1: Mild Difficulty; 2: Moderate Difficulty; 3: Severe Difficulty; 4: Extreme or cannot do; then scores assigned to each of the items were summed (<https://www.sralab.org/rehabilitation-measures/world-health-organization->

[disabilityassessment-schedule-ii](https://www.sralab.org/rehabilitation-measures/world-health-organization-disabilityassessment-schedule-ii)). Accordingly, the lowest and the highest possible scores of the tool are 0 and 48, respectively. Higher scores reflect higher levels of disability.<sup>20</sup>

The WHODAS 2.0 is a common multi-dimensional questionnaire, which is applicable to measure the level of disability across many conditions and has been validated in several languages.<sup>14</sup> So that, this schedule has been translated into sixteen languages in fourteen countries and has been reported to have an adequate internal consistency, construct and discriminate validity, and one-factor structure for the 12-item WHODAS 2.0.<sup>21</sup> The reliability coefficient was calculated at 0.92 in the present study. The instrument is validated for use in the Persian language through a PERSIAN Traffic Cohort study conducted by RTIRC.

Moreover, some of the information at baseline and hospital information system (HIS) in this study include some crash-related exposures such as crash mechanism, demographic information, and hospitalization-related predictors like having surgery, hospitalized ward, status at discharge time, having physiotherapy, as well as external causes (as mentioned in International Classification of Diseases version 10 (ICD-10), chapter XX), trauma clinical characteristics (ICD-10, chapter XI). This information was merged with those obtained from the questionnaire and then was analyzed.

### Data analysis

Proportions and means (standard deviation (SD)) were calculated for categorical and continuous variables respectively, including patients' characteristics, hospitalization information, and functional status based on 12-item WHODAS 2.0. The normality of the distribution of variables was tested and confirmed by One-Sample Kolmogorov-Smirnov Test. Spearman correlation test was used to express the relationship among dimensions of WHODAS 2.0. In addition, Mann-Whitney, Kruskal–Wallis tests were used to assess the relationship among demographic, hospital

factors, and WHODAS domains. Data were analyzed using the SPSS 25 (IBM software Version 25.0, Chicago, IL, USA). We considered  $p < 0.05$  to be statistically significant.

**Results**

Of the 180 patients in this study, 149 (82.8%) were male. The mean age (SD) was 38.8 (16.7) years old, and 103 injured people (57.2%) were car drivers or passengers. Injured organs of the body are classified into six groups. The most affected organs were head and face (36.7%), and the least was upper extremity (5.7%). Most of the participants were hospitalized in the trauma department (97 patients), and half of the participants had surgery after at Road Traffic Accident (RTA). 85.5% of patients were recovering at discharge. In Figure1-3, some hospital information of injured people is presented.

The mean WHODAS2.0 score was 17.8 (SD=9.1). About 20% of patients had scored more than 24. The highest score was estimated for the self-care dimension 3.3 (SD=1.8), and the lowest score for getting along with others 2.4 (SD=1.3). The scores of other dimensions include cognition, mobility,

life activities, and participation were calculated at 2.8 (SD=1.5), 3.2 (SD=2.4), 2.9 (SD=2.2), and 3 (SD=1.7), respectively. In addition, physiotherapy was not prescribed or performed for 134 (74.4%) patients. Only 46 (25.6%) patients completed their physiotherapy. The frequency and mean score of items based on gender are presented in Table 1.

Statistically, all dimensions of the questionnaire were positively correlated, and these correlations were significant ( $p < 0.001$ ). The highest correlation was between scores of mobility and life activities ( $r = 0.778$ ), and the lowest correlation was between getting along with and mobility ( $r = 0.351$ ) (Table 2).

There was no significant difference between WHODAS 2.0 domains scores and the role of injured people based on Kruskal–Wallis test. In addition, Kruskal–Wallis test between individual domains of WHODAS 2.0 and age group revealed a significant relationship in domains of life activities ( $X^2 = 11.34$ ,  $p = 0.01$ ), mobility ( $X^2 = 11.19$ ,  $p = 0.01$ ), and participation ( $X^2 = 11.34$ ,  $p = 0.02$ ). Therefore, a U Mann-Whitney test was performed to identify the source of the difference (Table 3). Also, U Mann-Whitney analysis among WHODAS 2.0 domains and other qualitative independent variables revealed some significant results that are presented in Table 3.




Figure 1. Role of injured people




Figure 2. Injury localization




Figure 3. Number of patients admitted to different wards of the hospital

**Table 1: Frequency and mean scores of WHODAS 2.0 items**

Dimension	Item	Gender	Frequency of five-point Likert (%)					Mean (SD)
			No problem	Mild problem	Moderate problem	Severe problem	Very severe problem	
<b>Cognition</b>	Learning a new task, for example, learning how to get to a new place?	Male	0	130(87.2)	0	12(8.1)	7(3.9)	Male: 2.8(1.5)
		Female	0	26(83.9)	1(3.2)	4(12.9)	0	
	Concentrating on doing something for ten minutes?	Male	0	108 (72.5)	11(7.4)	19(12.8)	11(7.4)	Female: 2.7(1.2)
		Female	0	23(74.2)	2(6.5)	6(19.4)	0	
<b>Mobility</b>	Standing for long periods such as 30 minutes?	Male	49(53)	13 (8.7)	35 (23.5)	0	22(14.8)	Male: 3.1(2.4)
		Female	12(38.7)	6(19.4)	8(25.8)	0	5(16.1)	
	Walking a long distance such as a kilometer (or equivalent)?	Male	0	78(52.3)	17(11.4)	37(24.8)	17(11.4)	Female: 3.6(2.4)
		Female	0	11(35.5)	5(16.1)	10(32.3)	5(16.1)	
<b>Self-care</b>	Washing your whole body?	Male	0	98(65.8)	13(8.7)	31(20.8)	7(4.7)	Male: 3.2(1.8)
		Female	0	15(48.8)	4(12.9)	11(35.5)	1(3.2)	
	Getting dressed?	Male	0	103(69.1)	12(8.1)	27(18.1)	7(4.7)	Female: 3.6(1.8)
		Female	0	19(61.3)	4(12.9)	7(22.6)	1(3.2)	
<b>Getting along with others</b>	Dealing with people, you do not know?	Male	0	130(87.2)	8(5.4)	5(3.4)	6(4)	Male: 2.5(1.3)
		Female	0	30(96.8)	0	1(3.2)	0	
	Maintaining a friendship?	Male	0	130(87.2)	4(12.7)	10(6.7)	5(3.4)	Female: 2.1(0.74)
		Female	0	28(90.3)	2(6.5)	1(3.2)	0	
<b>life activities</b>	Your day-to-day work/school	Male	78(52.3)	16 (10.7)	40 (26.8)	10 (6.7)	5(3.4)	Male: 2.8(2.1)
		Female	8(25.8)	4(12.9)	17(54.8)	2(6.5)	0	
	Taking care of your household responsibilities?	Male	0	85(57)	13(8.7)	40(26.8)	11(7.4)	Female: 3.7(1.9)
		Female	0	8(25.8)	4(12.9)	16(51.6)	2(6.5)	
<b>Participation</b>	How much of a problem did you have in joining in community activities (for example, festivities, religious or other activities) in the same way as anyone else can?	Male	0	103(69.1)	11(7.4)	28(18.8)	7(4.7)	Male: 3(1.7)
		Female	0	17(54.8)	6(19.4)	7(22.6)	1(3.2)	
	How much have you been emotionally affected by your health problems?	Male	0	121(81.2)	4(2.7)	14(9.4)	10(6.7)	Female: 3.2(1.5)
		Female	0	22(71)	3(9.7)	6(19.4)	0	

**Table 2:** Spearman correlation coefficients matrix and probability values of WHODAS2.0 dimensions

Variable	Cognition	Mobility	Self-care	Getting along with others	Life activities	Participation
<b>Cognition</b>	1					
<b>Mobility</b>	0.572** 0.000	1				
<b>Self-care</b>	0.547** 0.000	0.623** 0.000	1			
<b>Getting along with others</b>	0.573** 0.000	0.351** 0.000	0.406** 0.000	1		
<b>Life activities</b>	0.0.585** 0.000	0.778** 0.000	0.753** 0.000	0.386** 0.000	1	
<b>Participation</b>	0.604** 0.000	0.602** 0.000	0.547** 0.000	0.497** 0.000	0.631** 0.000	1

\*\*  $P < 0.0001$ **Table 3:** Mann-Whitney analysis test results of participants' characteristics and WHODAS 2.0 dimensions

Variable		U Mann-Whitney	Z	Sig (2-tailed)
According to gender				
<b>Dimension</b>	<b>Self-care</b>	3285	-2.43*	0.01
According to having physiotherapy				
<b>Dimension</b>	<b>life activities</b>	2078	-2.45*	0.014
	<b>Cognition</b>	2158	-2.46*	0.014
	<b>Mobility</b>	1818	-3.43**	0.001
	<b>Self-care</b>	1971	-3.07**	0.002
	<b>Participation</b>	2227	-2.01*	0.04
According to head and face injury				
	<b>Self-care</b>	3285	-2.43*	0.015
According to spinal cord injury				
<b>Dimension</b>	<b>life activities</b>	1581	-1.83***	0.066
	<b>Mobility</b>	1578	-1.82***	0.069
	<b>Self-care</b>	1642	-1.66***	0.096
According to upper extremity injury				
<b>Dimension</b>	<b>life activities</b>	695	-3**	0.003
	<b>Mobility</b>	773	-2.53*	0.011
	<b>Self-care</b>	728	-3**	0.003
	<b>Participation</b>	928	-1.87***	0.06
According to 20-40 and 41-60 age group				
<b>Dimension</b>	<b>life activities</b>	893	-2.32*	0.01
	<b>Mobility</b>	1708	-1.78***	0.07
According to 20-40 and upper 60 age group				
<b>Dimension</b>	<b>life activities</b>	893	-2.58*	0.01
	<b>Mobility</b>	853	-2.8**	0.005
	<b>Participation</b>	914	-2.61**	0.009
According to lower 20 and upper 60 age group				
<b>Dimension</b>	<b>life activities</b>	105	-2***	0.058
	<b>Mobility</b>	85	-2.56*	0.01
	<b>Participation</b>	92	-2.49*	0.02
According to lower 20 and 41-60 age group				
<b>Dimension</b>	<b>life activities</b>	184	-1.86***	0.06
	<b>Mobility</b>	183	-1.9***	0.057

\*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.1$

## Discussion

This pilot study in the post-crash study of PERSIAN Traffic Cohort is one of the few studies in Iran that explores the long-term burden of patients due to road traffic crashes admitted to two referral hospitals in Tabriz City. The collected information was extracted from road traffic registry system. It provided a description of functional status among participants, and its probable determinants based on available data. The study was managed to create evidence on the profile of patients affected by the road traffic crashes, which is very important for policymaking to effectively meet their needs.

In the present study, the mean score of WHODAS 2.0 showed a low functioning disorder. However, the mean was over 24 and more for about 20%, which indicated the existence of a disability. There is some research following our findings. A one-year cohort study in three European countries reported some levels of psychological distress one year after a traumatic event as well as physical disability among more than 30%, at the same time, based on WHODAS 2.0 schedule.<sup>22</sup> Another follow-up study of RTIs survivors in the UK hospital revealed that there was a substantial impact on physical activity, large injury costs, potentially high Quality-Adjusted Life Year (QALY) losses, using the Short Form-36 Health Survey version 2 (SF-36v2), European Quality of Life-5 Dimensions (EQ-5D), and Epidemiologic Studies Depression Scale (CES-D) scales.<sup>23</sup> Rissanen et al. reported that even relatively minor RTIs could lead to a considerably lower Health-Related Quality of Life (HRQoL), especially among women, compared to the non-injured reference group.<sup>24</sup>

In this study, the most reported function disabilities were related to domains of self-care,

mobility, participation, and life activities, in a descending order that it may be due to participants' young age. The disrupted daily activities and impaired mobility were barriers to the active population's work and professional activities. Based on studies, about 60% of road crashes victims are young aged 14–45.<sup>25–26</sup> Schluter et al. with the study on the functional capacity outcome of 619 participants showed 54% experienced no limitations, 16% experienced minor limitations, and 29% experienced major limitations, 12 months after hospitalization due to injury.<sup>27</sup> Other studies of patients injured in road crash presented the rates of Post-Traumatic Stress Disorder (PTSD) (up to 45%), the occurrence of depressive symptoms (10%), anxiety symptoms (about 40%), and travel phobia (20%).<sup>28,29</sup> Kuo et al. studied patients with Traumatic Brain Injury (TBI) and Spinal Cord Injury (SCI) using WHODAS and indicated the most functional problems in life activities dimension. However, the score of this dimension was statistically significant between the two groups.<sup>30</sup>

This study revealed that factors like age, gender, injury localization, and physiotherapy were relevant to WHODAS dimensions. A longitudinal cohort study produced previous disability, obesity, and injury severity were related to post-injury disability among hospitalized and non-hospitalized.<sup>31</sup> In another study, ethnicity, length of stay (LOS), education, and comorbidity were significant predictors of outcome based on the Glasgow Outcome Scale Extended (GOS-E).<sup>32</sup> Alharbi et al. in a comprehensive review confirmed some factors associated with functional outcomes among adult trauma patients and clustered into six categories; (1) location and severity of the injury; (2) hospital predictive; (3) demographic features; (4) social and family supports; (5) compensation system process and fault in the road



traffic crash; (6) previous health status.<sup>33</sup> Several of these factors were also confirmed in the present study.

There is no significant difference between WHODAS 2.0 domains scores and the role of injured people in the present study. Being a two-wheel rider or a pedestrian was revealed in the literature as a predictive factor for the risk of poor outcome in some studies.<sup>34-35</sup> Physiotherapy was associated with mobility, cognition, and life activities based on results. Similarly, the results of a single-blind prospective randomized trial on 71 patients were shown to have significant improvement in the severity of neck pain and cervical movement after outpatient physiotherapy one and two months after injury. In addition, 66 patients who received inclusive advice on home mobilization from a physiotherapist presented a similar improvement.<sup>36</sup>

Some main variables affect the recovery process. Kevin et al. reported that patients with severe injuries (LOS  $\geq$  7 days and Injury Severity Score (ISS) 12+) had poorer recovery 12 months after the injury. Also, post-injury mediators include baseline 12-Item Short Form Survey (SF12) scores, baseline pain, baseline psychological status, and compulsory third party insurance (CTP) claimant have an essential role in influencing long-term health outcomes.<sup>37</sup> Tournier et al. resulted in whiplash injuries; the pain was an intermediate factor between whiplash grading, health status, and Quality of Life (QOL).<sup>38</sup> Elbers et al. confirmed some factors that contributed to weak outcomes containing the role of negative perception and the stress-related claims process.<sup>39</sup> It can be concluded from these studies, some hospital factors, comprising the length of stay and the severity of the injury, affect long-term outcomes. On the other hand, mental state, baseline pain, and even the patient's financial status affects long-term health status after trauma.

## Conclusion

The current study identified some functional disabilities among patients sustaining RTIs. It was evident from the results that a proportion of patients don't recover six months after the injury and suffer a disability, especially in self-care, mobility, and life activities. In addition, Age, gender, physiotherapy, injury localization were related to WHODAS 2.0 scores.

These results highlight the importance of a comprehensive understanding of the long-term impact of the injury on an individual. People with poor functional or mental health are at-risk and need special attention. A short screening and assessing disability could speed up identifying disabilities and prevent costly assessment. Also, such screening helps to efficiently allocate recourses and policymaking. Future research with larger samples of patients is necessary to increase our understanding of long-lasting disability. Due to the lack of post-event information, in many parts of Iran, it is recommended to establish an integrated post-injury registry system including items such as the person's mental state, as well as a comorbidity in much more details and so on.

## Acknowledgments

We would like to thank the vice secretary of health, the Iranian Ministry of Health and Medical Education, and the Research Deputy of Tabriz University of Medical Sciences for their financial support. Special thanks are extended to the Road Traffic Injury Prevention Research Center of Tabriz University of Medical Sciences and Mr. Malek Abazari for their general and technical contribution.

## Conflict of Interest Disclosures

None.

## Funding Sources

The project of PERSIAN Traffic Cohort is jointly supported as a national project by the vice secretary of health, the Iranian Ministry of Health and Medical Education, and Tabriz University of Medical Sciences [grant number- 63795]. It is also part of two research projects of Ph.D. theses conducted by Yalda Mousazadeh (under thesis number:58043) and Nasrin Shahedifar (under thesis number: 64813).

## Authors' Contributions

AJ, FP, HP, MR, and HS designed and coordinated the study. YM and RZ conducted the literature review. AJ, RZ, ASH, SHB, and YM contributed to the data collection. YM, AJ, HS, NSH, and ZI analyzed and interpreted the findings. YM and HS wrote the initial draft of the manuscript. All authors have read and approved the final manuscript.

### Ethical Statement

The current study was approved by the Regional Research Ethics Committee of Tabriz University of Medical Sciences (ethical code: IR.TBZMED.REC.1398.543). Although patients' participation in the research was voluntary, and they informed oral consent was obtained before the interview.

### References

- Martino C, Russo E, Santonastaso DP, Gamberini E, Bertoni S, Padovani E, et al. Long-term outcomes in major trauma patients and correlations with the acute phase. *World J Emerg Surg.* 2020;15(6):1-7.
- Mousazadeh Y, Sadeghi-Bazargani H, Janati A, Pouraghaei M, Rahmani F. Prediction of mortality risk in patients with traffic injury: A case study in Tabriz hospitals, Iran. *Arch Trauma Res.* 2019;8(2):104-9.
- Holtslag HR, van Beeck EF, Lindeman E, Leenen LP. Determinants of long-term functional consequences after major trauma. *J Trauma Acute Care Surg.* 2007;62(4):919-27.
- Abedzadeh-kalahroudi M, Razi E, Sehat M, Asadi-Lari M. Psychometric properties of the world health organization disability assessment schedule II-12 Item (WHODAS II) in trauma patients. *Injury.* 2016;47(5):1104-8.
- Saadat S, Yousefifard M, Asady H, Jafari AM, Fayaz M, Hosseini M. The most important causes of death in Iranian population; a retrospective cohort study. *Emerg.* 2015;3(1):16-21.
- Rezaei S, Arab M, Matin BK, Sari AA. Extent, consequences and economic burden of road traffic crashes in Iran. *J. Inj. Violence Res.* 2014;6(2):57-63.
- Soberg HL, Bautz-Holter E, Roise O, Finset A. Long-term multidimensional functional consequences of severe multiple injuries two years after trauma: a prospective longitudinal cohort study. *J. Trauma Acute Care Surg.* 2007;62(2):461-70.
- Maas AI, Menon DK, Adelson PD, Andelic N, Bell MJ, Belli A, et al. Traumatic brain injury: integrated approaches to improve prevention, clinical care, and research. *Lancet Neuro.* 2017;16(12):987-1048.
- Currens JAB, Coats TJ. The timing of disability measurements following injury. *Injury.* 2000;31(2):93-8.
- Van Beeck E. Priorities in injury epidemiology. *Eur J Epidemiol.* 2004; 19(5):401-3.
- Axelsson E, Lindsäter E, Ljytsson B, Andersson E, Hedman-Lagerluf E. The 12-item Self-Report World Health Organization Disability Assessment Schedule (WHODAS) 2.0 Administered via the internet to individuals with anxiety and stress disorders: A psychometric investigation based on data from two clinical trials. *JMIR ment health.* 2017;4(4): e58.
- Puul M, Cieza A, Stucki G. Psychometric properties of the WHODASII in rehabilitation patients. *Qual Life Res.* 2007;16(9):1521-31.
- Ետևոն TB, Chatterji S, Kostanjsek N, Rehm J, Kennedy C, Epping-Jordan J, et al. Developing the World Health Organization disability assessment schedule 2.0. *Bull World Health Organ.* 2010;88(11):815-23.
- Luciano JV, Ayuso-Mateos JL, Fernández A, Serrano-Blanco A, Roca M, Haro JM. Psychometric properties of the twelve item World Health Organization Disability Assessment Schedule II (WHO-DAS II) in Spanish primary care patients with a first major depressive episode. *J Affect Disord.* 2010;121(1-2): 52-8.
- Ristner G, Andersson R, Johansson L, Johansson S-E, Ponzer S. Sense of coherence and lack of control in relation to outcome after orthopaedic injuries. *Injury.* 2000;31(10):751-6.
- Mayou R, Bryant B. Outcome in consecutive emergency department attenders following a road traffic accident. *Br J Psychiatry.* 2001;179(6):528-34.
- Mousazadeh Y. Developing a Hospital Performance Assessment Model for Patients Management with Traffic Injuries [dissertation]. Tabriz, Iran: Tabriz University of Medical Sciences, School of Management and Medical Informatics; 2019.
- Sadeghi Bazargani H, Somi MH, Shahedifar N, Poustchi H, Bazargan-Hejazi SH, Sadeghi V et al., PERSIAN Traffic Safety and Health Cohort: A Study Protocol on Long Term Post-Crash Mental and Physical Health Consequences, PLoS one, (Preprint)
- Road traffic crashes 2020 [Iranian legal medicine organization]. Tehran: legal medicine organization online Resources, Inc; c 2018-19[updated 2019 Mar 16; cited 2019 Jul] Available from: [https://www.lmo.ir/web\\_directory/53999-.html](https://www.lmo.ir/web_directory/53999-.html).
- Andrews G, Kemp A, Sunderland M, Von Korff M, Ustun TB. Normative data for the 12 item WHO Disability Assessment Schedule 2.0. *PLoS one.* 2009;4(12): e8343.
- Horowitz M, Wilner N, Alvarez W. Impact of Event Scale: A measure of subjective stress. *Psychosom Med.* 1979;41(3):209-18.
- Papadakaki M, Ferraro OE, Orsi C, Otte D, Tzamalouka G, VonderGeest M, et al. Psychological distress and physical disability in patients sustaining severe injuries in road traffic crashes: results from a one-year cohort study from three European countries. *Injury.* 2017;48(2):297-306.
- Barnes J, Thomas P. Quality of life outcomes in a hospitalized sample of road users involved in crashes. *Annu Proc Assoc Adv Automot Med.* 2006; 50: 253-268.
- Rissanen R, Iíver J, Hasselberg M, Berg H-Y. Quality of life following road traffic injury: the impact of age and gender. *Qual Life Res.* 2020; 29(6):1587-96.
- Ziyab AH, Akhtar S. Incidence and trend of road traffic injuries and related deaths in Kuwait: 2000-2009. *Injury.* 2012;43(10):2018-22.
- González MPS, Sotos FE, Ponce BT. Impact of provincial characteristics on the number of traffic accident victims on interurban roads in Spain. *Accid Anal Prev.* 2018; 118:178-89.
- Schluter PJ, McClure RJ. Predicting functional capacity outcome 12 months after hospitalized injury. *ANZ J surg.* 2006;76: 886-93.
- Ehring T, Ehlers A, Glucksman E. Do cognitive models help in predicting the severity of posttraumatic stress disorder, phobia, and

---

depression after motor vehicle accidents? A prospective longitudinal study. *J. Consult. Clin. Psychol.* 2008;76(2):219-30.

29. Chahraoui K, Laurent A, Bioy A, Quenot J-P. Psychological experience of patients 3 months after a stay in the intensive care unit: A descriptive and qualitative study. *J. Crit. Care.* 2015;30(3):599-605.

30. Kuo C-Y, Liou T-H, Chang K-H, Chi W-C, Escorpizo R, Yen C-F, et al. Functioning and disability analysis of patients with traumatic brain injury and spinal cord injury by using the world health organization disability assessment schedule 2.0. *Int J Environ Res Public Health.* 2015;12(4):4116-27.

31. Derrett S, Samaranyaka A, Wilson S, Langley J, Ameratunga S, Cameron ID, et al. Prevalence and predictors of sub-acute phase disability after injury among hospitalised and non-hospitalised groups: a longitudinal cohort study. *PLoS one.* 2012;7(9): e44909.

32. Kersten P, Czuba K, Anstiss D, Maheswaran R, Smith G, Kayes N, et al. Predictors of disability outcome after major trauma. *Ann Phys Rehabil Med.* 2018;61: e122.

33. Alharbi R, Mosley I, Miller C, Hillel S, Lewis V. Factors associated with physical, psychological and functional outcomes in adult trauma patients following road traffic crash: a scoping literature review. *Transport res interdis pers* 2019; 3:100061.

34. WHOD V, Prevention I, Violence WH, Prevention I. Organization WH. Global status report on road safety: time for action: World Health Organization. 2009.

35. Mayou R, Bryant B. Consequences of road traffic accidents for different types of road user. *Injury.* 2003;34(3):197-202.

36. McKinney L, Dorman J, Ryan M. The role of physiotherapy in the management of acute neck sprains following road-traffic accidents. *Arch Emerg Med J.* 1989; 6:27-33.

37. Hung KK, Kifley A, Brown K, Jagnoor J, Craig A, et al. Impacts of injury severity on long-term outcomes following motor vehicle crashes. *BMC public health.* 2021; 21:1-13.

38. Tournier C, Hours M, Charnay P, Chossegras L, Tardy H. Five years after the accident, whiplash casualties still have poorer quality of life in the physical domain than other mildly injured casualties: analysis of the ESPARR cohort. *BMC Public Health.* 2015;16(1): 1-3.

39. Elbers, N.A., Akkermans, A.J., Lockwood, K. et al. Factors that challenge health for people involved in the compensation process following a motor vehicle crash: a longitudinal study. *BMC Public Health.* 2015; 15:339-49.