



**UNIVERSITY  
OF TURKU**

This is a self-archived – parallel published version of supplementary data of an original article. This version may differ from the original in pagination and typographic details. When using please cite the original.

AUTHOR Niina M. Santio, Päivi J. Koskinen

TITLE PIM kinases: From survival factors to regulators of cell motility – supplementary data

YEAR 2017


DOI <https://doi.org/10.1016/j.biocel.2017.10.016>

VERSION Author's accepted manuscript

COPYRIGHT License: [CC BY NC ND](#)


CITATION Niina M. Santio, Päivi J. Koskinen, PIM kinases: From survival factors to regulators of cell motility, The International Journal of Biochemistry & Cell Biology, Volume 93, 2017, Pages 74-85, ISSN 1357-2725, <https://doi.org/10.1016/j.biocel.2017.10.016>

## Supplementary Data



### Supplementary Figure 1. *PIM* expression in healthy tissues

Relative *PIM* mRNA levels in healthy human tissue samples were obtained from the IST Online™ ([ist.medisapiens.com](http://ist.medisapiens.com)) database (Kilpinen et al., 2008).



**Supplementary Figure 2. PIM expression in cancer tissues**

Relative *PIM* mRNA levels in human cancer tissue samples were obtained from the IST Online™ (ist.medisapiens.com) database (Kilpinen et al., 2008).

## Reference

Kilpinen S., Autio R., Ojala K., Iljin K., Bucher E., Sara H., Pisto T., Saarela M., Skotheim R.I., Bjorkman M., Mpindi J.P., Haapa-Paananen S., Vainio P., Edgren H., Wolf M., Astola J., Nees M., Hautaniemi S. and Kallioniemi O., 2008. Systematic bioinformatic analysis of expression levels of 17,330 human genes across 9,783 samples from 175 types of healthy and pathological tissues, *Genome Biol.* 9, R139, <https://doi.org/10.1186/gb-2008-9-9-r139>.