

# 1 **MAGIC and MWL monitoring of the blazar TXS** 2 **0506+056 in the 2018/2019 season**

---

**K. Satalecka<sup>\*1</sup>, E. Bernardini<sup>2</sup>, W. Bhattacharyya<sup>1</sup>, M. Cerruti<sup>3</sup>, V. Fallah Ramazani<sup>4</sup>,  
L. Foffano<sup>2</sup>, S. Inoue<sup>5</sup>, E. Prandini<sup>2</sup>, Ch. Righi<sup>6</sup>, N. Sahakyan<sup>7</sup>, F. Tavecchio<sup>6</sup> for the  
MAGIC Collaboration<sup>†</sup>**

<sup>1</sup>DESY, 15738 Zeuthen, Germany; <sup>2</sup>Università di Padova and INFN, I-35131 Padova, Italy;  
<sup>3</sup>Universitat de Barcelona, ICCUB, IEEC-UB, E-08028 Barcelona, Spain; <sup>4</sup>Finnish Centre of  
Astronomy with ESO (FINCA), University of Turku, FI-20014 Turku, Finland; <sup>5</sup>ICRR, The  
University of Tokyo, 277-8582 Chiba, Japan; <sup>6</sup>National Institute for Astrophysics (INAF),  
I-00136 Rome, Italy; <sup>7</sup>ICRANet-Armenia at NAS RA, 0019 Yerevan, Armenia  
E-mail: [konstancja.satalecka@desy.de](mailto:konstancja.satalecka@desy.de), [you@youremail.com](mailto:you@youremail.com),  
[you@youremail.com](mailto:you@youremail.com), ...

The gamma-ray blazar TXS 0506+056, was discovered in VHE gamma-rays by the MAGIC telescopes in 2017 in a follow-up campaign of a high energy neutrino event IceCube-170922A (IC+Fermi+MAGIC++, Science 361, eaat1378 (2018)). Subsequent multivavelength (MWL) observations and theoretical modeling in a frame of hadro-leptonic emission confirmed that this source could be a potential cosmic ray and neutrino emitter (MAGIC Collaboration, Ansoldi et al., (2018)). This is, by far, the most significant association between a high-energy neutrino and an astrophysical source emitting gamma rays and X-rays. TXS 0506+056 is a key object to help the astrophysics community to establish connections between high-energy neutrinos and astrophysical sources. Accurate and contemporaneous MWL spectral measurements are essential ingredients to achieve this goal. In the conference, we present the measurements from the MAGIC and MWL monitoring of this source, spanning the time period from November 2017 till February 2019. These include the lowest VHE gamma-ray emission state measured from this source so far as well as a flaring episode in December 2018.

*36th International Cosmic Ray Conference -ICRC2019-  
July 24th - August 1st, 2019  
Madison, WI, U.S.A.*

---

<sup>\*</sup>Speaker.

<sup>†</sup><https://magic.mpp.mpg.de/> For collaboration list see PoS(ICRC2019)1177

### 3 1. Introduction

4 Active Galactic Nuclei (AGN), especially of the blazar class, dominate the very high energy  
5 (VHE;  $> 100$  GeV) electromagnetic sky. Blazars, as all AGN are highly luminous sources, powered  
6 by a supermassive black hole. Additionally they display relativistic jets, one of which is pointed  
7 near the line of sight of the observer, which results in a relativistic boosting of the emission. Blazars  
8 are also considered as prime candidates for hadronic accelerators [1, 2, 3]. In this scenario, their  
9 VHE photon emission can be explained as secondary radiation from interactions of accelerated  
10 protons and the surrounding photon or matter fields. Another by-product of those interactions are  
11 neutrinos. If observed, they should provide us with a smoking-gun signature of hadronic interac-  
12 tions.

13 IceCube, a  $1\text{km}^3$  neutrino telescope located at the South Pole, has performed several extensive  
14 searches for neutrino emission from AGN [4]. So far, no strong correlation between the cata-  
15 loged AGN and astrophysical neutrinos has been found, and blazar contributions to the all-sky  
16 astrophysical neutrino flux has been constrained at the level of 27% [5]. Nevertheless, the all-sky  
17 astrophysical neutrino flux shows an isotropic distribution, favouring an extragalactic origin and  
18 AGN still remain as promising neutrino sources.

19 TXS 0506+056, a bright gamma-ray emitting blazar, is of special interest for the hadronic  
20 accelerators case. On September 22, 2017 a 290-TeV neutrino event was detected by IceCube  
21 (IceCube-170922A) in spatial and temporal coincidence with an enhanced  $\gamma$ -ray emission state of  
22 this source [6]. The significance of this coincidence was estimated to be at the  $3\sigma$  level. MAGIC  
23 was the first IACT to detect VHE gamma-rays from this object shortly after IceCube issued the  
24 alert. During the subsequent monitoring of the source the MAGIC collaboration and other alerted  
25 instruments collected a large multiwavelength data set, essential for modeling and interpretation of  
26 the TXS 0506+056 emission mechanism. The MAGIC collaboration proposed an interpretation in  
27 the hadro-leptonic spine-layer emission framework [7], where the results of the modeling strongly  
28 support the hypothesis of TXS 0506+056 being a neutrino and cosmic ray emitter.

29 The TXS 0506+056 case is, our most compelling evidence of hadronic emission in blazars to  
30 date. Therefore during the period from November 2017 till February 2019 MAGIC, together with  
31 MWL partners continued to monitor the source behaviour. In the conference, we present the data  
32 collected during this extended multiwavelength monitoring campaign, including longer periods of  
33 low state emission, as well as a VHE  $\gamma$ -ray flaring episode in December 2018.

### 34 2. MAGIC monitoring 2017/2019

35 MAGIC observed the source TXS 0506+056 for a total of about 90 h between November  
36 2017 and February 2019 within a dedicated monitoring program aimed at collecting a long-term  
37 data sample of the source. The analysis was performed on  $\sim 80$  h of good-quality data with zenith  
38 angle range between  $22^\circ$  and  $50^\circ$ , using the MAGIC Analysis and Reconstruction Software [8, 9].

39 During most of the monitored period the source was in *low state* ( $\sim 75$  h), with an average flux  
40 above 80 GeV about 10-15 times lower than the flux observed during the flare in October 2018.  
41 This is the lowest VHE  $\gamma$ -ray emission level observed from this source so far. On December 1<sup>st</sup> and  
42 3<sup>rd</sup> 2018 (MJD 58453 and 58455) an enhanced emission was observed with fluxes comparable with

43 the flare detected by MAGIC in October 2017, shortly after the neutrino alert IceCube-170922A  
 44 [7]. On December 4<sup>th</sup>, the MAGIC collaboration issued an Astronomers Telegram (ATel #12260)  
 45 to encourage further MWL observations of TXS 0506+056. Several ToOs, including the X-ray and  
 46 optical instruments were triggered. Table 1 shows the summary of the TXS 0506+056 flux levels  
 47 as measured by MAGIC.

Table 1: MAGIC measurements of TXS 0506+056

Data set	Duration [h]	Significance	VHE activity
MJD 58453	2.5	$3.8\sigma$	High
MJD 58455	1.8	$5.4\sigma$	Very high
Rest	74.4	$4.0\sigma$	Low

### 48 3. Multiwavelength observations

49 The preliminary multi-band fluxes that were shown at the conference were collected from pub-  
 50 lic archives of each instrument. Large part of those observations were coordinated with MAGIC  
 51 in order to ensure simultaneous exposure. Dedicated ToO observations were performed with  
 52 Swift/XRT [10] and UVOT as well as NuSTAR [11]. The MAGIC observations were usually  
 53 accompanied by the KVA [12] optical telescope, additional measurements were performed with  
 54 REM after the flare in December 2018. TXS 0506+056 is also systematically monitored by the  
 55 ASAS-SN project [13].




Figure 1: Fractional variability parameter for each instrument

56 Clear variability is observed in all wavelengths. In order to quantify it, we calculated the  
 57 *fractional variability parameter*  $F_{var}$  according to [14]. The most pronounced variability  $F_{var} \sim 0.3-$   
 58  $0.5$  is observed in the highest energies (X-rays and  $\gamma$ -rays). The optical and UV bands display a  
 59 moderate variability of  $F_{var} \sim 0.1-0.2$ . As for VHE gamma rays, the very low activity during these  
 60 two years yielded low significance flux measurements with MAGIC for most of the single night  
 61 observations, which precluded to compute a meaningful fractional variability in this band.

62 The two distinct activity levels at VHE gamma rays motivated us to produce two broadband  
 63 SEDs, for the low state and the flaring state, and we modeled this emission using a leptohadronic

64 theoretical scenario as discussed in [15]. The largest contemporaneous MWL data sample was  
65 collected for the VHE  $\gamma$ -ray low emission state. Its spectral energy distribution shows a clear  
66 similarity in shape and flux level at the low energy peak to the one previously observed [7]. The  
67 flux level of the high energy peak was measured to be lower than before. For the flaring episode in  
68 December 2018, the VHE  $\gamma$ -ray flux is compatible with the previously measured flares.

#### 69 4. Summary and Outlook

70 TXS 0506+056 is a key object to help the astrophysics community to establish connections  
71 between high-energy neutrinos and astrophysical sources. Accurate and contemporaneous MWL  
72 spectral measurements are essential ingredients to achieve this goal. In the conference, we pre-  
73 sented the results from the MAGIC and MWL monitoring of this source, spanning the time period  
74 from November 2017 till February 2019.

75 In comparison to the previously published results, TXS 0506+056 displayed a very low VHE  
76  $\gamma$ -ray emission state during most of the observed nights, with the exception of the flaring activity  
77 observed on December 1<sup>st</sup> and 3<sup>rd</sup> 2018. The MWL light curve shows clear signs of variability,  
78 especially in the high energy range (X-ray and HE  $\gamma$ -ray), as quantified by the fractional variability  
79 parameter  $F_{var}$ .

80 We plan to perform a dedicated, low energy optimized analysis of the MAGIC data with a goal  
81 of recovering the signal below the currently obtained energy threshold. These results, along with  
82 an updated theoretical interpretation will be presented in an upcoming publication.

#### 83 5. Acknowledgments

84 The authors gratefully acknowledge financial support from the agencies and organizations  
85 listed here: [https://magic.mpp.mpg.de/acknowledgments\\_ICRC2019/](https://magic.mpp.mpg.de/acknowledgments_ICRC2019/)

86 This project has received funding from the European Union's Horizon2020 research and inno-  
87 vation programme under the Marie Skłodowska-Curie grant agreement no 664931.

#### 88 References

- 89 [1] K. Mannheim, *A&A* **269** (Mar., 1993) 67–76.  
90 [2] K. Mannheim, *Astropart.Phys.* **3** (1995) 295–302.  
91 [3] F. Halzen and E. Zas, *ApJ* **488** (Oct., 1997) 669–674.  
92 [4] M. G. Aartsen et al., *ApJ* **807** (Jul, 2015) 46.  
93 [5] M. G. Aartsen et al., *ApJ* **835** (Jan, 2017) 45.  
94 [6] **IceCube, Fermi-LAT, MAGIC, ++** Collaboration, M. G. Aartsen et al., *Science* **361** (2018)  
95 eaat1378.  
96 [7] S. Ansoldi et al., *ApJ* **863** (Aug, 2018) L10.

- 97 [8] A. Moralejo et al., *arXiv e-prints* (Jul, 2009) arXiv:0907.0943.
- 98 [9] J. Aleksić et al., *Astropart. Phys.* **72** (Jan., 2016) 76–94.
- 99 [10] D. N. Burrows et al., *The Swift X-Ray Telescope*, in *X-Ray and Gamma-Ray Instrumentation*  
100 *for Astronomy XIII* (K. A. Flanagan and O. H. W. Siegmund, eds.), vol. 5165 of *Proceedings*  
101 *of the SPIE*, pp. 201–216, Feb., 2004.
- 102 [11] F. A. Harrison et al., *ApJ* **770** (June, 2013) 103.
- 103 [12] K. Nilsson, , et al., *A&A* **620** (Dec., 2018) A185.
- 104 [13] C. S. Kochanek, B. J. Shappee, K. Z. Stanek, T. W.-S. Holoien, T. A. Thompson, J. L. Prieto,  
105 S. Dong, J. V. Shields, D. Will, C. Britt, D. Perzanowski, and G. Pojmański, *PASP* **129** (Oct.,  
106 2017) 104502.
- 107 [14] S. Vaughan, R. Edelson, R. S. Warwick, and P. Uttley, *MNRAS* **345** (Nov., 2003) 1271–1284.
- 108 [15] M. Cerruti et al., *PoS(ICRC2019)646*, 2019.