

Coronary Volume to Left Ventricular Mass Ratio in Patients With Hypertension



Sophie E. van Rosendaal, MD^a, Alexander R. van Rosendaal, MD^a, Jurrien H. Kuneman, MD^a, Manesh R. Patel, MD^b, Bjarne Linde Nørgaard, MD, PhD^c, Timothy A. Fairbairn, MB, ChB, PhD^d, Koen Nieman, MD, PhD^{e,f}, Takashi Akasaka, MD^g, Daniel S. Berman, MD^h, Lynne M. Hurwitz Kowek, MD^b, Gianluca Pontone, MD, PhDⁱ, Tomohiro Kawasaki, MD^j, Niels Peter Rønnow Sand, MD, PhD^{k,l}, Jesper M. Jensen, MD, PhD^c, Tetsuya Amano, MD^m, Michael Poon, MDⁿ, Kristian A. Øvrehus, MD, PhD^o, Jeroen Sonck, MD^p, Mark G. Rabbat, MD^q, Campbell Rogers, MD^r, Hitoshi Matsuo, MD, PhD^s, Jonathon A. Leipsic, MD^t, Nina Ajmone Marsan, MD, PhD^a, J. Wouter Jukema, MD, PhD^{a,u}, Jeroen J. Bax, MD, PhD^{a,v}, Antti Saraste, MD, PhD^{u,w}, and Juhani Knuuti, MD, PhD^{a,w,*}

The coronary vascular volume to left ventricular mass (V/M) ratio assessed by coronary computed tomography angiography (CCTA) is a promising new parameter to investigate the relation of coronary vasculature to the myocardium supplied. It is hypothesized that hypertension decreases the ratio between coronary volume and myocardial mass by way of myocardial hypertrophy, which could explain the detected abnormal myocardial perfusion reserve reported in patients with hypertension. Individuals enrolled in the multicenter ADVANCE (Assessing Diagnostic Value of Noninvasive FFRCT in Coronary Care) registry who underwent clinically indicated CCTA for analysis of suspected coronary artery disease with known hypertension status were included in current analysis. The V/M ratio was calculated from CCTA by segmenting the coronary artery luminal volume and left ventricular myocardial mass. In total, 2,378 subjects were included in this study, of whom 1,346 (56%) had hypertension. Left ventricular myocardial mass and coronary volume were higher in subjects with hypertension than normotensive patients (122.7 ± 32.8 g vs 120.0 ± 30.5 g, $p = 0.039$, and $3,105.0 \pm 992.0$ mm³ vs $2,965.6 \pm 943.7$ mm³, $p < 0.001$, respectively). Subsequently, the V/M ratio was higher in patients with hypertension than those without (26.0 ± 7.6 mm³/g vs 25.3 ± 7.3 mm³/g, $p = 0.024$). After correcting for potential confounding factors, the coronary volume and ventricular mass remained higher in patients with hypertension (least square mean difference estimate: 196.3 (95% confidence intervals [CI] 119.9 to 272.7) mm³, $p < 0.001$, and 5.60 (95% CI 3.42 to 7.78) g, $p < 0.001$, respectively), but the V/M ratio was not significantly different (least square mean difference estimate: 0.48 (95% CI -0.12 to 1.08) mm³/g, $p = 0.116$). In conclusion, our findings do not support the hypothesis that the abnormal perfusion reserve would be caused by reduced V/M ratio in patients with hypertension. © 2023 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) (Am J Cardiol 2023;199:100–109)

^aDepartment of Cardiology, Leiden University Medical Center, Leiden, The Netherlands; ^bDivision of Cardiology, Department of Medicine, Duke University Medical Center, Duke Clinical Research Institute, Duke University School of Medicine, Durham, North Carolina; ^cDepartment of Cardiology, Aarhus University Hospital, Aarhus, Denmark; ^dDepartment of Cardiology, University of Liverpool, Liverpool Heart and Chest Hospital, Liverpool, United Kingdom; ^eDepartment of Cardiovascular Medicine, Stanford University, Stanford, California; ^fDepartment of Radiology, Stanford University, Stanford, California; ^gDepartment of Cardiovascular Medicine, Wakayama Medical University, Wakayama, Japan; ^hDivision of Nuclear Imaging, Department of Imaging, Cedars-Sinai Heart Institute, Los Angeles, California; ⁱCentro Cardiologico Monzino, Milan, Italy; ^jCardiovascular Center, Shin Koga Hospital, Fukuoka, Japan; ^kDepartment of Cardiology, University Hospital of Southern Denmark, Esbjerg, Denmark; ^lDepartment of Regional Health Research, University of Southern Denmark, Esbjerg, Denmark; ^mDepartment of Cardiology, Aichi Medical University, Aichi, Japan; ⁿDepartment of Noninvasive Cardiac Imaging,

Northwell Health, New York, New York; ^oDepartment of Cardiology, Odense University Hospital, Odense, Denmark; ^pCardiovascular Center Aalst, OLV Clinic, Aalst, Belgium; ^qDivision of Cardiology, Loyola University Chicago, Chicago, Illinois; ^rHeartFlow Inc., Redwood City, California; ^sDepartment of Cardiovascular Medicine, Gifu Heart Center, Gifu, Japan; ^tDepartment of Radiology, St Paul's Hospital, Vancouver, British Columbia, Canada; ^uNetherlands Heart Institute, Utrecht, The Netherlands; ^vHeart Center, Turku University Hospital and University of Turku, Turku, Finland; and ^wTurku PET Center, Turku University Hospital and University of Turku, Turku, Finland. Manuscript received January 4, 2023; revised manuscript received and accepted April 17, 2023.

Funding: none.

See page 108 for Declaration of Conflict of Interest.

*Corresponding author: Tel: +358 500 592 998; fax: +358 2 231 8191.

E-mail address: juhani.knuuti@tyks.fi (J. Knuuti).

Hypertension causes changes in the coronary circulation, characterized by a reduction of the coronary vascular reserve.^{1–10} Left ventricular (LV) hypertrophy, usually a complication of hypertension because of sustained elevated afterload, is associated with a reduction in maximal coronary vasodilator reserve^{11–13} and an increase in myocardial oxygen demand.^{14–16} The ratio of the total epicardial coronary artery luminal volume to LV myocardial mass (V/M ratio) is considered a parameter capable of revealing a potential physiologic imbalance between coronary blood supply and myocardial demand.¹⁷ Low V/M ratios were associated with more advanced coronary artery disease (CAD), reduced myocardial blood flow, and lesion-specific fractional flow reserve <0.80.^{18,19} Based on previous studies observing reduced coronary flow reserve in patients with hypertension, we hypothesized that patients with hypertension may have a lower V/M ratio than normotensive patients.

Methods

ADVANCE (Assessing Diagnostic Value of Noninvasive FFRCT in Coronary Care) is a multinational (38 sites in Europe, North America, and Japan) registry with

prospective follow-up data of patients being investigated for clinically suspected CAD designed to understand the effect of coronary computed tomography angiography (CCTA)-derived fractional flow reserve on clinical practice. The study design has been described earlier in detail.²⁰ In summary, subjects were enrolled from July 15, 2015 to October 20, 2017. Patients aged >18 years with documented stenosis of at least 30% on CCTA were included. Patients with an insufficient CCTA image quality, an inability to comply with follow-up requirements, and a life expectancy <1 year were excluded.

For the present analysis, patients with known hypertension status and available coronary artery luminal volume and LV myocardial mass analysis were included (Figure 1). Patients with diabetes were excluded to reduce the confounding effects of diabetes on V/M.²¹ The study was conducted in accordance with the Declaration of Helsinki. All individuals provided written informed consent after local institutional review board review and approval.

All CCTA scans were performed with ≥64-row multidetector computed tomography scanners. If the prescan heart rate was >60 beats/min, patients received metoprolol before the CCTA scan, unless contraindicated. Sublingual nitrates were administered to all patients before scanning. Coronary




Figure 1. Flowchart of study population.

arteries with a diameter of ≥ 2 mm were evaluated for stenosis severity in accordance with current guidelines according to the clinical site procedures.²² HeartFlow Inc. (Redwood City, California), a central core laboratory, computed the V/M analyses, which has been described previously.^{20,23–26} In short, a patient-specific anatomic epicardial model of the coronary tree was derived from the CCTA images provided. The total coronary arterial luminal volume is calculated by the summation of all the segmented coronary arteries. The volume of the myocardium extracted from CCTA was multiplied by 1.05 g/ml, an average value for myocardial tissue density, resulting in the left ventricle myocardial mass.²⁷ Subsequently, the ratio between the total coronary artery luminal volume and the LV myocardial mass was calculated. Because of software development during the study time period, the analysis of the V/M ratio could not be performed in all patients.

The diagnoses of hypertension were based on the medical history in the electronic case report forms and defined as systolic blood pressure values of ≥ 140 mm Hg and/or diastolic blood pressure values of ≥ 90 mm Hg requiring

treatment. Among patients with anatomically obstructive and without obstructive CAD the coronary artery luminal volume and LV myocardial mass were separately analyzed. Obstructive CAD was defined as $\geq 50\%$ diameter stenosis.

Statistical analyses were performed with SAS version 9.4 (SAS institute, Cary, North Carolina). Continuous variables with a normal distribution are presented as mean $\pm$ SD and were compared using the Student's *t* test or one-way analysis of variance, as appropriate. Non-normally distributed continuous variables are presented as median with (twenty-fifth to seventy-fifth interquartile range) and were compared using the Mann-Whitney *U* test. Categorical variables are presented as absolute numbers and percentages and were compared using the chi-square test. To correct for potential confounding effects on the coronary artery luminal volume, LV myocardial mass, and V/M ratio, analysis of covariance models were used. Age, body mass index (BMI), hyperlipidemia, gender, number of vessels with obstructive CAD, and the degree of maximum stenosis were used as covariates in this analysis. The differences in total coronary artery luminal volume, LV myocardial mass,

Table 1
Baseline characteristics of the overall population and according to hypertension status

	Total (n=2,378)	Hypertension (n=1,346)	No hypertension (n=1,032)	p Value
Age, (y)				
N	2,272	1288	984	<0.001
Mean $\pm$ SD	66.1 $\pm$ 10.4	67.8 $\pm$ 9.6	63.9 $\pm$ 11.0	
Min, max	15.0, 93.0	34.0, 93.0	15.0, 92.0	
Male sex	1,564 (65.8%)	849 (63.1%)	715 (69.3%)	0.002
BMI, (kg/m²)				
N	2,347	1332	1,015	<0.001
Mean $\pm$ SD	26.1 $\pm$ 4.7	26.4 $\pm$ 4.9	25.6 $\pm$ 4.4	
Min, max	14.9, 63.7	15.8, 63.7	14.9, 55.5	
Diamond Forrester CAD likelihood				
N	2,251	1281	970	0.544
Mean $\pm$ SD	50.9 $\pm$ 20.0	51.2 $\pm$ 19.9	50.6 $\pm$ 20.1	
Min, max	5.3, 92.5	8.0, 92.5	5.3, 92.5	
Hyperlipidemia				
Yes	1,368 (57.5%)	888 (66.0%)	480 (46.5%)	<0.001
No	995 (41.8%)	448 (33.3%)	547 (53.0%)	
Unknown	15 (0.6%)	10 (0.7%)	5 (0.5%)	
Tobacco use				
Current smoker	364 (15.3%)	191 (14.2%)	173 (16.8%)	0.072
Ex-smoker	815 (34.3%)	484 (36.0%)	331 (32.1%)	
Never smoked	1,020 (42.9%)	571 (42.4%)	449 (43.5%)	
Unknown	179 (7.5%)	100 (7.4%)	79 (7.7%)	
Angina status				
Typical	465 (19.6%)	264 (19.6%)	201 (19.5%)	0.028
Atypical	868 (36.5%)	467 (34.7%)	401 (38.9%)	
Dyspnea	274 (11.5%)	148 (11.0%)	126 (12.2%)	
Non-cardiac pain	150 (6.3%)	85 (6.3%)	65 (6.3%)	
None	604 (25.4%)	375 (27.9%)	229 (22.2%)	
Unknown	17 (0.7%)	7 (0.5%)	10 (1.0%)	
CCS angina class				
Grade I	109/ 465 (23.4%)	55/ 264 (20.8%)	54/ 201 (26.9%)	0.210
Grade II	264/ 465 (56.8%)	152/ 264 (57.6%)	112/ 201 (55.7%)	
Grade III	42/ 465 (9.0%)	27/ 264 (10.2%)	15/ 201 (7.5%)	
Grade IV	6/ 465 (1.3%)	5/ 264 (1.9%)	1/ 201 (0.5%)	
Unknown	44/ 465 (9.5%)	25/ 264 (9.5%)	19/ 201 (9.5%)	

Data are presented as mean $\pm$ standard deviation or number (percentage), as appropriate.

BMI = body mass index; CAD = coronary artery disease; CCS = Canadian Cardiovascular Society.

and V/M ratio between hypertensive and normotensive patients are presented as least square (LS) mean difference estimate with corresponding 95% confidence intervals (CIs). A 2-sided $p < 0.05$ was considered statistically significant.

Results

A total of 5,083 individuals were enrolled in the ADVANCE registry. Of these, 2,378 patients without diabetes with known hypertension status and measured V/M ratio were included in present analysis. Hypertension was present in 1,346 patients (60%). Baseline patient demographic and clinical characteristics of the enrolled patients

are listed in Table 1. Patients with hypertension were older (67.8 ± 9.6 vs 63.9 ± 11.0 years, $p < 0.001$) and had a higher BMI (26.4 ± 4.9 vs 25.6 ± 4.4 kg/m², $p < 0.001$). In addition, patients with hypertension had more frequently a history of hyperlipidemia ($p < 0.001$) and were more likely to be female ($p = 0.002$).

Patients with hypertension had more frequently obstructive CAD by anatomic CCTA evaluation ($p = 0.017$; Table 2). In the quantitative analysis, the volume of epicardial coronary arteries was higher in patients with hypertension ($3,105.0 \pm 992.0$ mm³ vs $2,965.6 \pm 943.7$ mm³, $p = 0.001$). The LV myocardial mass was higher in patients with hypertension as well (122.7 ± 32.8 g vs 120.0 ± 30.5 g, $p = 0.039$). This resulted in a higher V/M ratio in

Table 2

Coronary computed tomography angiography parameters of patients according to hypertension status

	Total (n=2,378)	Hypertension (n=1,346)	No hypertension (n=1,032)	p Value
CCTA anatomical finding				
Without obstructive stenosis <50%	711 (29.9%)	376 (27.9%)	335 (32.5%)	0.017
Obstructive stenosis ≥50%	1,663 (69.9%)	968 (71.9%)	695 (67.3%)	
Unknown	4 (0.2%)	2 (0.1%)	2 (0.2%)	
Non-severe stenosis ≤70%	1,676 (70.5%)	943 (70.1%)	733 (71.0%)	0.596
Severe stenosis >70%	698 (29.4%)	401 (29.8%)	297 (28.8%)	
Unknown	4 (0.2%)	2 (0.1%)	2 (0.2%)	
Degree stenosis				
Normal (0%)	15 (0.6%)	6 (0.4%)	9 (0.9%)	0.040
Minimal (0%–30%)	136 (5.7%)	62 (4.6%)	74 (7.2%)	
Mild (30%–50%)	560 (23.5%)	308 (22.9%)	252 (24.4%)	
Moderate (50%–70%)	965 (40.6%)	567 (42.1%)	398 (38.6%)	0.004
Severe (70%–90%)	493 (20.7%)	288 (21.4%)	205 (19.9%)	
Sub-total/occluded (≥90%/occluded)	205 (8.6%)	113 (8.4%)	92 (8.9%)	
Unknown	4 (0.2%)	2 (0.1%)	2 (0.2%)	0.004
0	711 (29.9%)	376 (27.9%)	335 (32.5%)	
1	1,062 (44.7%)	592 (44.0%)	470 (45.5%)	
2	420 (17.7%)	259 (19.2%)	161 (15.6%)	0.002
3	181 (7.6%)	117 (8.7%)	64 (6.2%)	
4	0	0	0	
Unknown	4 (0.2%)	2 (0.1%)	2 (0.2%)	
Rate of obstructive CAD per vessel				
LAD stenosis <50%	1,069 (45.0%)	584 (43.4%)	485 (47.0%)	0.080
LAD stenosis ≥50%	1,309 (55.0%)	762 (56.6%)	547 (53.0%)	
LCX stenosis <50%	1,860 (78.2%)	1,030 (76.5%)	830 (80.4%)	
LCX stenosis ≥50%	518 (21.8%)	316 (23.5%)	202 (19.6%)	0.002
RCA stenosis <50%	1,760 (74.0%)	963 (71.5%)	797 (77.2%)	
RCA stenosis ≥50%	618 (26.0%)	383 (28.5%)	235 (22.8%)	
Coronary volume - myocardial mass				
Epicardial coronary artery volume (mm ³)				
N	2,378	1346	1,032	0.001
Mean±SD	3,044.5±973.6	3,105.0±992.0	2,965.6±943.7	
Min, max	704.6, 7,891.2	732.1, 7,891.2	704.6, 7,198.4	
Left ventricle myocardial mass (g)				
N	2,378	1346	1,032	0.039
Mean±SD	121.6±31.8	122.7±32.8	120.0±30.5	
Min, max	54.9, 324.1	54.9, 324.1	56.9, 308.9	
Coronary volume /mass (mm ³ /g)				
N	2,378	1346	1,032	0.024
Mean±SD	25.7±7.5	26.0±7.6	25.3±7.3	
Min, max	6.8, 62.5	6.8, 61.9	7.2, 62.5	

Number of vessels with anatomically obstructive CAD (>50% DS). Data are presented as mean±standard deviation or number (percentage), as appropriate.

CAD = coronary artery disease; CCTA = coronary computed tomography angiography; DS = diameter stenosis; LAD = left anterior descending artery; LCX = left circumflex artery; RCA = right coronary artery.




Figure 2. Bar chart showing the least squares means, after correcting for potential confounding factors, of the coronary volume, left ventricular mass and V/M ratio for patients with and without hypertension. A: total cohort. B: subjects with obstructive coronary artery disease. C: Subjects without obstructive CAD* = The variable "Number of vessels with obstructive coronary artery disease" is removed in current analysis because of collinearity with "maximum stenosis %." Inference did not change, but values changed slightly.

patients with hypertension than patients without hypertension (26.0 ± 7.6 mm³/g vs 25.3 ± 7.3 mm³/g, $p = 0.024$). When correcting for the differences in baseline and CCTA characteristics, the coronary volume and myocardial mass remained significantly higher in patients with hypertension (LS mean difference estimate: 196.3 [95% CI 119.9 to 272.7] mm³, $p < 0.001$; LS mean difference estimate: 5.60 [95% CI 3.42 to 7.78] g, $p < 0.001$, respectively; Figure 2, Table 3). Whereas the V/M ratio showed no significant

difference between hypertensive and normotensive patients (LS mean difference estimate 0.48 [95% CI -0.12 to 1.08] mm³/g, $p = 0.116$).

Because CAD has known effects on coronary volume, the groups with and without obstructive CAD were analyzed separately (Table 4). Obstructive CAD was present in 1,663 subjects (69.9%), of whom 968 (58.2%) had hypertension. In individuals with obstructive CAD, patients with hypertension were more often male ($p = 0.009$), were older

Table 3

Coronary volume, cardiac mass and coronary volume : mass ratio corrected for potential confounding variables

Model effect	LS mean difference (95% CI)	p Value
Total segmented volume		
Hypertension (yes/no)	196.3 (119.9, 272.7)	<0.001
Age		0.735
BMI		<0.001
Hyperlipidemia (yes/no)		0.002
Sex (male/female)		<0.001
Number of vessels with obstructive CAD (0,1,2,3)		<0.001
Maximum stenosis % (0, >0–<30, ≥30–<50, ≥50–≤70, >70–≤90, >90)		<0.001
Myocardial mass		
Hypertension (yes/no)	5.60 (3.42, 7.78)	<0.001
Age		<0.001
BMI		<0.001
Hyperlipidemia (yes/no)		<0.001
Sex (male/female)		<0.001
Number of vessels with obstructive CAD (0,1,2,3)		0.047
Maximum stenosis % (0, >0–<30, ≥30–<50, ≥50–≤70, >70–≤90, >90)		<0.001
Volume : mass ratio		
Hypertension (yes/no)	0.48 (–0.12, 1.08)	0.116
Age		<0.001
BMI		<0.001
Hyperlipidemia (yes/no)		0.629
Sex (male/female)		0.007
Number of vessels with obstructive CAD (0,1,2,3)		<0.001
Maximum stenosis % (0, >0–<30, ≥30–<50, ≥50–≤70, >70–≤90, >90)		<0.001

BMI = body mass index; CAD = coronary artery disease; CI = confidence interval; LS = least squares.

($p < 0.001$), had a higher BMI ($p = 0.004$), and had more frequently a history of hyperlipidemia ($p < 0.001$) (Table 4). Coronary volume did not differ significantly between hypertensive and normotensive patients with obstructive CAD ($3,026.4 \pm 971.5 \text{ mm}^3$ vs $2,937.5 \pm 918.5 \text{ mm}^3$, $p = 0.058$). Moreover, the LV mass was not significantly different between the 2 groups ($123.6 \pm 33.4 \text{ g}$ vs $121.8 \pm 29.4 \text{ g}$; $p = 0.243$). Accordingly, the V/M ratio was comparable between the 2 groups ($25.2 \pm 7.3 \text{ mm}^3/\text{g}$ vs $24.7 \pm 7.2 \text{ mm}^3/\text{g}$, $p = 0.209$). When we corrected for potential confounding variables, the epicardial coronary artery volume and myocardial mass were significantly higher in patients with hypertension than normotensive patients (LS mean difference estimate: 135.21 [95% CI 45.3 to 225.1] mm^3 , $p = 0.003$ and LS mean difference estimate: 4.92 [95% CI 2.30 to 7.55] g , $p < 0.001$ respectively]; Figure 2, Table 5). However, the V/M ratio was not significantly different between the 2 groups (LS mean difference estimate: 0.15 [95% CI –0.54 to 0.84] mm^3/g , $p = 0.671$).

Hypertension was present in 376 of 711 patients (53%) without obstructive CAD. Patients with hypertension were more frequent female ($p = 0.024$), older ($p < 0.001$), had a higher BMI ($p = 0.006$), and had more frequently a history of hyperlipidemia ($p < 0.001$) (Table 4). Coronary volume was higher in patients with hypertension than normotensive patients without obstructive CAD ($3,305.8 \pm 1,019.1 \text{ mm}^3$ vs $3,023.8 \pm 995.4 \text{ mm}^3$, $p < 0.001$), whereas LV mass did not differ significantly between the groups ($120.5 \pm 31.1 \text{ g}$ vs $116.2 \pm 32.4 \text{ g}$, $p = 0.074$). Consequently, the V/M ratio was significantly higher ($28.1 \pm 7.9 \text{ mm}^3/\text{g}$ vs $26.5 \pm 7.2 \text{ mm}^3/\text{g}$, $p = 0.007$) in patients with hypertension than normotensive patients. Coronary artery volume remained

significantly higher in patients with hypertension after correction for potential confounding variables (LS mean difference estimate: 352.20 [95% CI 208.37 to 496.04] mm^3 , $p < 0.001$; Figure 2, Table 6). The myocardial mass after correction for confounding variables was significantly higher in patients with hypertension as well (LS mean difference estimate: 7.24 [95% CI 3.33 to 11.14] g , $p < 0.001$). The V/M ratio remained significant higher in the patients with hypertension (LS mean difference estimate: 1.33 [95% CI 0.15 to 2.51] mm^3/g , $p = 0.028$; Table 5).

Discussion

This study assessed the impact of hypertension on the V/M ratio. The hypothesis was that the known reduced myocardial perfusion reserve in patients with hypertension may be partially explained by an abnormally low V/M ratio, likely because of myocardial hypertrophy not accompanied by increase in vascular volume. The main results demonstrate that the V/M ratio was not decreased in patients with hypertension, suggesting that the increased myocardial mass was compensated by increased vascular volume, leading to preserved V/M ratio.

The V/M ratio has been shown to be reduced in patients with CAD.¹⁸ This is expected because CAD typically affects the coronary lumen and the vasodilatory capacity. We recently found that V/M ratio is reduced also in patients with diabetes, even when CAD was taken into account as a confounding factor.²¹ In this study, we excluded patients with diabetes and also analyzed the patients with and without obstructive CAD separately. An interesting finding was that in patients without obstructive CAD, the V/M ratio was

Table 4

Baseline characteristics and coronary computed tomography and coronary computed tomography angiography parameters of patients with anatomically obstructive and without obstructive CAD according to hypertension status

	Obstructive CAD ($\geq 50\%$ DS)				Without obstructive CAD ($< 50\%$ DS)			
	Total (n=1,663)	Hypertension (n=968)	No hypertension (n=695)	p Value	Total (n=711)	Hypertension (n=376)	No hypertension (n=335)	p Value
Baseline patient characteristics								
Age, (y)								
N	1597	930	667	<0.001	672	357	315	<0.001
Mean $\pm$ SD	66.6 $\pm$ 10.3	68.0 $\pm$ 9.6	64.6 $\pm$ 10.7		65.0 $\pm$ 10.7	67.2 $\pm$ 9.5	62.4 $\pm$ 11.4	
Min, max	26.0, 93.0	40.0, 93.0	26.0, 92.0		15.0, 90.0	34.0, 89.0	15.0, 90.0	
Male sex	1,150 (69.2%)	645 (66.6%)	505 (72.7%)	0.009	412 (57.9%)	203 (54.0%)	209 (62.4%)	0.024
BMI, (kg/m²)								
N	1648	960	688	0.004	695	370	325	0.006
Mean $\pm$ SD	25.9 $\pm$ 4.5	26.2 $\pm$ 4.6	25.5 $\pm$ 4.2		26.4 $\pm$ 5.2	26.9 $\pm$ 5.4	25.9 $\pm$ 4.8	
Min, max	14.9, 53.1	15.8, 53.1	14.9, 42.6		15.9, 63.7	18.0, 63.7	15.9, 55.5	
Diamond forrester CAD likelihood								
N	1585	926	659	0.656	663	354	309	0.206
Mean $\pm$ SD	53.2 $\pm$ 20.0	53.0 $\pm$ 20.0	53.4 $\pm$ 19.9		45.6 $\pm$ 19.0	46.5 $\pm$ 18.9	44.6 $\pm$ 19.2	
Min, max	8.0, 92.5	8.0, 92.5	8.0, 92.5		5.3, 92.5	8.0, 92.5	5.3, 88.9	
Hyperlipidemia								
Yes	959 (57.7%)	636 (65.7%)	323 (46.5%)	<0.001	406 (57.1%)	251 (66.8%)	155 (46.3%)	<0.001
No	697 (41.9%)	327 (33.8%)	370 (53.2%)		297 (41.8%)	120 (31.9%)	177 (52.8%)	
Unknown	7 (0.4%)	5 (0.5%)	2 (0.3%)		8 (1.1%)	5 (1.3%)	3 (0.9%)	
Rate of obstructive CAD per vessel								
LAD stenosis $< 50\%$	354 (21.3%)	206 (21.3%)	148 (21.3%)	0.995	NA	NA	NA	NA
LAD stenosis $\geq 50\%$	1,309 (78.7%)	762 (78.7%)	547 (78.7%)		NA	NA	NA	
LCX stenosis $< 50\%$	1,145 (68.9%)	652 (67.4%)	493 (70.9%)		NA	NA	NA	
LCX stenosis $\geq 50\%$	518 (31.1%)	316 (32.6%)	202 (29.1%)	0.017	NA	NA	NA	NA
RCA stenosis $< 50\%$	1,045 (62.8%)	585 (60.4%)	460 (66.2%)		NA	NA	NA	
RCA stenosis $\geq 50\%$	618 (37.2%)	383 (39.6%)	235 (33.8%)		NA	NA	NA	
Coronary volume - myocardial mass								
Epicardial coronary artery volume (mm³)								
N	1663	968	695	0.058	711	376	335	<0.001
Mean $\pm$ SD	2,989.2 $\pm$ 950.5	3,026.4 $\pm$ 971.5	2,937.5 $\pm$ 918.5		3,172.9 $\pm$ 1,017.1	3,305.8 $\pm$ 1,019.1	3,023.8 $\pm$ 995.4	
Min, max	704.6, 7,415.5	732.1, 7,415.5	704.6, 7,055.6		889.6, 7,891.2	1,181.3, 7,891.2	889.6, 7,198.4	
Left ventricle myocardial mass (g)								
N	1663	968	695	0.243	711	376	335	0.074
Mean $\pm$ SD	122.9 $\pm$ 31.8	123.6 $\pm$ 33.4	121.8 $\pm$ 29.4		118.5 $\pm$ 31.7	120.5 $\pm$ 31.1	116.2 $\pm$ 32.4	
Min, max	54.9, 324.1	54.9, 324.1	56.9, 247.1		58.3, 308.9	63.3, 264.6	58.3, 308.9	
Coronary volume / mass (mm³/g)								
N	1663	968	695	0.209	711	376	335	0.007
Mean $\pm$ SD	25.0 $\pm$ 7.3	25.2 $\pm$ 7.3	24.7 $\pm$ 7.2		27.3 $\pm$ 7.6	28.1 $\pm$ 7.9	26.5 $\pm$ 7.2	
Min, max	6.8, 62.5	6.8, 59.2	7.2, 62.5		9.8, 61.9	10.7, 61.9	9.8, 51.0	

Data are presented as mean $\pm$ standard deviation or number (percentage), as appropriate.

BMI = body mass index; CAD = coronary artery disease; DS = diameter stenosis; LAD = left anterior descending artery; LCX = left circumflex artery; RCA = right coronary artery.

higher in hypertensive patients, despite increased myocardial mass. In patients with obstructive CAD, V/M ratio was not significantly different between patients with and without hypertension, likely because of the confounding effect of CAD on the V/M ratio.

The concept of the V/M ratio was first described by Gould et al²⁸ and the method of assessing the V/M ratio is based on allometric scaling laws. Allometric scaling laws provide a model to predict the functional and structural properties of the cardiovascular system of mammals.²⁹ Choy et al³⁰ investigated the scaling laws of myocardial flow and mass in a porcine heart and reported a very tight linear relation between coronary artery luminal volume and

myocardial mass. Previous studies investigating the V/M ratio have shown that individuals with a low V/M ratio had reduced myocardial blood flow on positron emission tomography compared with patients with a high V/M ratio.¹⁸ Furthermore, Taylor et al¹⁹ concluded that the V/M ratio was independently associated with a fractional flow reserve below the ischemic threshold (≤ 0.80).

We hypothesized that the abnormal myocardial perfusion in patients with hypertension was caused by a reduced V/M ratio. LV hypertrophy is frequently associated with hypertension, increases the myocardial mass, and is considered a mechanism contributing to abnormal myocardial perfusion. However, this study shows a corresponding increase

Table 5

Coronary computed tomography angiography parameters corrected for potential confounding variables in patients with obstructive CAD

Model effect	LS mean difference (95% CI)	p Value
Total segmented volume		
Hypertension (yes/no)	135.21 (45.3, 225.1)	0.003
Age		0.790
BMI		<0.001
Hyperlipidemia (yes/no)		0.002
Sex (male/female)		<0.001
Number of vessels with obstructive CAD (0, 1, 2, 3)		<0.001
Maximum stenosis % (≥ 50 – ≤ 70 , > 70 – ≤ 90 , > 90)		<0.001
Myocardial mass		
Hypertension (yes/no)	4.92 (2.30, 7.55)	<0.001
Age		<0.001
BMI		<0.001
Hyperlipidemia (yes/no)		<0.001
Sex (male/female)		<0.001
Number of vessels with obstructive CAD (0, 1, 2, 3)		0.031
Maximum stenosis % (≥ 50 – ≤ 70 , > 70 – ≤ 90 , > 90)		0.002
Volume : mass ratio		
Hypertension (yes/no)	0.15 (–0.54, 0.84)	0.671
Age		<0.001
BMI		<0.001
Hyperlipidemia (yes/no)		0.371
Sex (male/female)		0.002
Number of vessels with obstructive CAD (0, 1, 2, 3)		<0.001
Maximum stenosis % (≥ 50 – ≤ 70 , > 70 – ≤ 90 , > 90)		<0.001

BMI = body mass index; CAD = coronary artery disease; CI = confidence interval; LS = least squares.

in coronary artery volume, leading to a preserved V/M ratio in patients with hypertension.

The increased coronary luminal volume in patients with hypertension we observed in this study is in line with previous research, showing luminal enlargement of proximal elastic arteries.^{31,32} Carotid and coronary arteries represent

large vessels, often referred to as “elastic arteries” or “conducting arteries” and are both central, predominantly elastic, and transport large volumes of blood away from the left ventricle to perfuse vital organs.³³ In addition, atherosclerotic disease and its potential confounding effect needs to be taken into account when calculating the V/M ratio

Table 6

Coronary computed tomography angiography parameters corrected for potential confounding variables in patients without obstructive CAD

Model effect	LS mean difference (95% CI)	p Value
Total segmented volume		
Hypertension (yes/no)	352.2 (208.4, 496.0)	<0.001
Age		0.950
BMI		0.001
Hyperlipidemia (yes/no)		0.239
Sex (male/female)		<0.001
Maximum stenosis % (0, > 0 – < 30 , ≥ 30 – < 50)		0.352
Myocardial mass		
Hypertension (yes/no)	7.24 (3.33, 11.14)	<0.001
Age		0.014
BMI		<0.001
Hyperlipidemia (yes/no)		0.043
Sex (male/female)		<0.001
Maximum stenosis % (0, > 0 – < 30 , ≥ 30 – < 50)		0.352
Volume : mass ratio		
Hypertension (yes/no)	1.33 (0.15, 2.51)	0.028
Age		0.002
BMI		<0.001
Hyperlipidemia (yes/no)		0.731
Sex (male/female)		0.627
Maximum stenosis % (0, > 0 – < 30 , ≥ 30 – < 50)		0.413

BMI = body mass index; CAD = coronary artery disease; CI = confidence interval; LS = least squares.

because the presence of atherosclerosis and reduced coronary volume has been linked. When the cohort is divided into patients with and without obstructive CAD, patients with obstructive CAD remain to have no significant different V/M ratio between patients with hypertension and normotensive patients. However, we observed in patients with hypertension without obstructive CAD an even higher V/M ratio than normotensive patients. The increase in coronary luminal volume is apparently larger than the increase of the ventricular mass. This effect is diminished in patients with obstructive CAD by the presence of more extensive atherosclerosis. Zhou et al.³⁴ observed that the diameter of the coronary artery is inversely associated with the severity of CAD. In addition, endothelial dysfunction because of atherosclerosis, with a subsequent reduction of vasodilator capacity contributes to a reduced coronary volume in these patients as well.³⁵

The observational design of the study has inherent limitations, including selection bias and unmeasured confounding. The registry may have been subject to referral bias inherent in local practices. In addition, information regarding the severity and duration of hypertension in the patients was lacking, and in our population, the increase of LV mass was small, despite being statistically significant. Antihypertensive treatment has been associated with the reduction of LV hypertrophy and might have a favorable effect on the matching between myocardial mass and perfusion.³⁶ Angiotensin-converting enzyme inhibitors were found to increase cardiac nitric oxide release and reduce oxygen consumption in coronary microvessels.^{37,38} The lack of data regarding antihypertensive treatment could be viewed as a limitation of the present study as well. Equally, this study did not adjust for the presence or absence of other cardiac diseases that affect myocardial blood flow reserve, such as valvular disease and hypertrophic cardiomyopathy. Lastly, the lack of information regarding the total plaque burden can be considered a limitation.

In conclusion, in contrast to our hypothesis, the V/M ratio was not decreased in patients with hypertension compared with patients without hypertension, and the abnormal coronary flow reserve in patients with hypertension is not likely caused by a reduced arterial volume to myocardial mass. Further studies are required using different cohorts to investigate the relation of flow reserve and V/M ratio.

Declaration of Competing Interest

The Department of Cardiology, Leiden University Medical Center, Leiden, the Netherlands has received unrestricted research grants from Bayer, Abbott Vascular, Medtronic, Biotronik, Boston Scientific, GE Healthcare (Little Chalfont, United Kingdom), and Edwards Lifesciences. Dr. Patel has received research grants from HeartFlow, Bayer (Abbott), Janssen, and the National Heart, Lung, and Blood Institute and has served on the advisory board for HeartFlow, Bayer, and Janssen. Dr. Nørgaard has received an unrestricted institutional research grant from HeartFlow Inc. Dr. Fairbairn has served on the Speakers Bureau for Heartflow. Dr. Nieman reports support from the NIH (NIH R01- HL141712; NIH R01 - HL146754) and reports

unrestricted institutional research support from Siemens Healthineers, consulting fees from Siemens Medical Solutions United States And Novartis, and equity in Lumen Therapeutics. Dr. Berman has received unrestricted research support from HeartFlow. Dr. Hurwitz Koweek has received research support and speaking fees from HeartFlow and Siemens. Dr. Pontone has received institutional research grant and/or honorarium a consultant/speaker from GE Healthcare, Boehringer, Bracco, Medtronic, Bayer, and HeartFlow. Dr. Rabbat has served as a consultant for HeartFlow. Dr. Rogers is employee of and owns equity in HeartFlow. Dr. Leipsic has received research grants from GE Healthcare and Edwards Lifesciences and has served as a consultant for and holds stock options in Circle cardiovascular Imaging and HeartFlow Inc. Dr. Jukema/his department has received research grants from and/or was speaker (with or without lecture fees) on a.o.(CME accredited) meetings sponsored by Amarin, Amgen, Athera, Biotronik, Boston Scientific, Dalcov, Daiichi Sankyo, Lilly, Medtronic, Merck-Schering-Plough, Novartis, Novo Nordisk, Pfizer, Roche, Sanofi Aventis (Bridgewater, New Jersey), the Netherlands Heart Foundation, CardioVascular Research the Netherlands, the Netherlands Heart Institute, and the European Community Framework KP7 Program. Dr. Bax received speaker fees from Abbot Vascular. Dr. Ajmone received speaker fees from Abbot Vascular and GE Healthcare. Dr. Saraste received consultancy fees from Amgen, Astra Zeneca, Boehringer Ingelheim, and Pfizer and speaker fees from Abbott, Astra Zeneca, and Bayer. Dr. Knuuti received consultancy fees from GE Healthcare and AstraZeneca and speaker fees from GE Healthcare, Bayer, Lundbeck, Boehringer Ingelheim, Pfizer, and Merck outside of the submitted work. The remaining authors have no conflicts of interest to declare.

1. Treasure CB, Klein JL, Vita JA, Manoukian SV, Renwick GH, Selwyn AP, Ganz P, Alexander RW. Hypertension and left ventricular hypertrophy are associated with impaired endothelium-mediated relaxation in human coronary resistance vessels. *Circulation* 1993;87:86–93.
2. Leschke M, Schoebel FC, Vogt M, Heintzen M, Kelm M, Motz W, Strauer BE. Reduced peripheral and coronary vasomotion in systemic hypertension. *Eur Heart J* 1992;13(suppl D):96–99.
3. Antony I, Nitenberg A, Foulst JM, Aptekar E. Coronary vasodilator reserve in untreated and treated hypertensive patients with and without left ventricular hypertrophy. *J Am Coll Cardiol* 1993;22:514–520.
4. Strauer BE, Vogt M, Motz W. ACE-inhibitors and coronary microcirculation. *Basic Res Cardiol* 1993;88(suppl 1):97–106.
5. Vogt M, Motz W, Strauer BE. Coronary haemodynamics in hypertensive heart disease. *Eur Heart J* 1992;13(suppl D):44–49.
6. Brush JE Jr, Faxon DP, Salmon S, Jacobs AK, Ryan TJ. Abnormal endothelium-dependent coronary vasomotion in hypertensive patients. *J Am Coll Cardiol* 1992;19:809–815.
7. Egashira K, Suzuki S, Hirooka Y, Kai H, Sugimachi M, Imaizumi T, Takeshita A. Impaired endothelium-dependent vasodilation of large epicardial and resistance coronary arteries in patients with essential hypertension. Different responses to acetylcholine and substance P. *Hypertension* 1995;25:201–206.
8. Motz W, Strauer BE. Improvement of coronary flow reserve after long-term therapy with enalapril. *Hypertension* 1996;27:1031–1038.
9. Brush JE Jr, Cannon RO 3rd, Schenke WH, Bonow RO, Leon MB, Maron BJ, Epstein SE. Angina due to coronary microvascular disease in hypertensive patients without left ventricular hypertrophy. *N Engl J Med* 1988;319:1302–1307.
10. Gaudieri V, Acampa W, Rozza F, Nappi C, Zampella E, Assante R, Mannarino T, Mainolfi C, Petretta M, Verberne HJ, Arumugam P, Cuocolo A. Coronary vascular function in patients with resistant

- hypertension and normal myocardial perfusion: a propensity score analysis. *Eur Heart J Cardiovasc Imaging* 2019;20:949–958.
11. Strauer BE. Ventricular function and coronary hemodynamics in hypertensive heart disease. *Am J Cardiol* 1979;44:999–1006.
 12. Opher D, Mall G, Zebe H, Schwarz F, Weihe E, Manthey J, Kübler W. Reduction of coronary reserve: a mechanism for angina pectoris in patients with arterial hypertension and normal coronary arteries. *Circulation* 1984;69:1–7.
 13. Pichard AD, Gorlin R, Smith H, Ambrose J, Meller J. Coronary flow studies in patients with left ventricular hypertrophy of the hypertensive type. Evidence for an impaired coronary vascular reserve. *Am J Cardiol* 1981;47:547–554.
 14. Laine H, Katoh C, Luotolahti M, Yki-Järvinen H, Kantola I, Jula A, Takala TO, Ruotsalainen U, Iida H, Haaparanta M, Nuuti J. Myocardial oxygen consumption is unchanged but efficiency is reduced in patients with essential hypertension and left ventricular hypertrophy. *Circulation* 1999;100:2425–2430.
 15. Galderisi M, de Simone G, Cicala S, Parisi M, D'Errico A, Innelli P, de Divitiis M, Mondillo S, de Divitiis O. Coronary flow reserve in hypertensive patients with hypercholesterolemia and without coronary heart disease. *Am J Hypertens* 2007;20:177–183.
 16. Frohlich ED. Hypertension left ventricular hypertrophy, and coronary flow reserve. *Adv Exp Med Biol* 1997;432:253–262.
 17. Ihdayhid AR, Fairbairn TA, Gulsin GS, Tzimas G, Danehy E, Updegrove A, Jensen JM, Taylor CA, Bax JJ, Sellers SL, Leipsic JA, Nørgaard BL. Cardiac computed tomography-derived coronary artery volume to myocardial mass. *J Cardiovasc Comput Tomogr* 2022;16:198–206.
 18. van Diemen PA, Schumacher SP, Bom MJ, Driessen RS, Everaars H, Stuijzand WJ, Raijmakers PG, van de Ven PM, Min JK, Leipsic JA, Knuuti J, Boellaard PR, Taylor CA, van Rossum AC, Danad I, Knaapen P. The association of coronary lumen volume to left ventricle mass ratio with myocardial blood flow and fractional flow reserve. *J Cardiovasc Comput Tomogr* 2019;13:179–187.
 19. Taylor CA, Gaur S, Leipsic J, Achenbach S, Berman DS, Jensen JM, Dey D, Bøtker HE, Kim HJ, Khem S, Wilk A, Zarins CK, Bezerra H, Lesser J, Ko B, Narula J, Ahmadi A, Øvrehus KA, St Goar F, De Bruyne B, Nørgaard BL. Effect of the ratio of coronary arterial lumen volume to left ventricle myocardial mass derived from coronary CT angiography on fractional flow reserve. *J Cardiovasc Comput Tomogr* 2017;11:429–436.
 20. Chinnaiyan KM, Akasaka T, Amano T, Bax JJ, Blanke P, De Bruyne B, Kawasaki T, Leipsic J, Matsuo H, Morino Y, Nieman K, Nørgaard BL, Patel MR, Pontone G, Rabbat M, Rogers C, Sand NP, Raff G. Rationale, design and goals of the HeartFlow assessing diagnostic value of non-invasive FFR_{CT} in Coronary Care (ADVANCE) registry. *J Cardiovasc Comput Tomogr* 2017;11:62–67.
 21. Kuneman JH, El Mahdiui M, van Rosendaal AR, van den Hoogen IJ, Patel MR, Nørgaard BL, Fairbairn TA, Nieman K, Akasaka T, Berman DS, Hurwitz Koweek LM, Pontone G, Kawasaki T, Rønnow Sand NP, Jensen JM, Amano T, Poon M, Øvrehus KA, Sonck J, Rabbat MG, De Bruyne B, Rogers C, Matsuo H, Bax JJ, Leipsic JA, Knuuti J. Coronary volume to left ventricular mass ratio in patients with diabetes mellitus. *J Cardiovasc Comput Tomogr* 2022;16:319–326.
 22. Leipsic J, Abbara S, Achenbach S, Cury R, Earls JP, Mancini GJ, Nieman K, Pontone G, Raff GL. SCCT guidelines for the interpretation and reporting of coronary CT angiography: a report of the Society of Cardiovascular Computed Tomography Guidelines Committee. *J Cardiovasc Comput Tomogr* 2014;8:342–358.
 23. Grover R, Leipsic JA, Mooney J, Kueh SH, Ohana M, Nørgaard BL, Eftekhari A, Bax JJ, Murphy DT, Hague CJ, Seidman MA, Blanke P, Sedlak T, Sellers SL. Coronary lumen volume to myocardial mass ratio in primary microvascular angina. *J Cardiovasc Comput Tomogr* 2017;11:423–428.
 24. Taylor CA, Fonte TA, Min JK. Computational fluid dynamics applied to cardiac computed tomography for noninvasive quantification of fractional flow reserve: scientific basis. *J Am Coll Cardiol* 2013;61:2233–2241.
 25. Gaur S, Achenbach S, Leipsic J, Mauri L, Bezerra HG, Jensen JM, Bøtker HE, Lassen JF, Nørgaard BL. Rationale and design of the HeartFlowNXT (HeartFlow analysis of coronary blood flow using CT angiography: NeXt sTeps) study. *J Cardiovasc Comput Tomogr* 2013;7:279–288.
 26. Kitabata H, Leipsic J, Patel MR, Nieman K, De Bruyne B, Rogers C, Pontone G, Nørgaard BL, Bax JJ, Raff G, Chinnaiyan KM, Rabbat M, Rønnow Sand NP, Blanke P, Fairbairn TA, Matsuo H, Amano T, Kawasaki T, Morino Y, Akasaka T. Incidence and predictors of lesion-specific ischemia by FFR_{CT}: learnings from the international ADVANCE registry. *J Cardiovasc Comput Tomogr* 2018;12:95–100.
 27. Fairbairn TA, Dobson R, Hurwitz-Koweek L, Matsuo H, Nørgaard BL, Rønnow Sand NP, Nieman K, Bax JJ, Pontone G, Raff G, Chinnaiyan KM, Rabbat M, Amano T, Kawasaki T, Akasaka T, Kitabata H, Binukrishnan S, Rogers C, Berman D, Patel MR, Douglas PS, Leipsic J. Sex differences in coronary computed tomography angiography-derived fractional flow reserve: lessons from ADVANCE. *JACC Cardiovasc Imaging* 2020;13:2576–2587.
 28. Gould KL. Does coronary flow trump coronary anatomy? *JACC Cardiovasc Imaging* 2009;2:1009–1023.
 29. West GB, Brown JH, Enquist BJ. A general model for the origin of allometric scaling laws in biology. *Science* 1997;276:122–126.
 30. Choy JS, Kassab GS. Scaling of myocardial mass to flow and morphometry of coronary arteries. *J Appl Physiol* (1985) 2008;104:1281–1286.
 31. Boutouyrie P, Bussy C, Lacolley P, Girerd X, Laloux B, Laurent S. Association between local pulse pressure, mean blood pressure, and large-artery remodeling. *Circulation* 1999;100:1387–1393.
 32. Hoeks AP, Willekes C, Boutouyrie P, Brands PJ, Willigers JM, Reneman RS. Automated detection of local artery wall thickness based on M-line signal processing. *Ultrasound Med Biol* 1997;23:1017–1023.
 33. Peace A, Van Mil A, Jones H, Thijssen DHJ. Similarities and differences between carotid artery and coronary artery function. *Curr Cardiol Rev* 2018;14:254–263.
 34. Zhou FF, Liu YH, Ge PC, Chen ZH, Ding XQ, Liu JY, Jia QW, An FH, Li LH, Wang LS, Ma WZ, Yang ZJ, Jia EZ. Coronary artery diameter is inversely associated with the severity of coronary lesions in patients undergoing coronary angiography. *Cell Physiol Biochem* 2017;43:1247–1257.
 35. Davignon J, Ganz P. Role of endothelial dysfunction in atherosclerosis. *Circulation* 2004;109(suppl 1):III27–III32.
 36. Parodi O, Neglia D, Palombo C, Sambuceti G, Giorgetti A, Marabotti C, Gallopin M, Simonetti I, L'Abbate A. Comparative effects of enalapril and verapamil on myocardial blood flow in systemic hypertension. *Circulation* 1997;96:864–873.
 37. Zhang X, Xie YW, Nasjletti A, Xu X, Wolin MS, Hintze TH. ACE inhibitors promote nitric oxide accumulation to modulate myocardial oxygen consumption. *Circulation* 1997;95:176–182.
 38. Ancion A, Tridetti J, Nguyen Trung ML, Oury C, Lancellotti P. A review of the role of bradykinin and nitric oxide in the cardioprotective action of angiotensin-converting enzyme inhibitors: focus on perindopril. *Cardiol Ther* 2019;8:179–191.