



## Redescriptions of two poorly known species of *Zodariellum* Andreeva & Tystshenko, 1968 (Araneae, Zodariidae) from Uzbekistan

**Sergei L. Esyunin**

Perm State University, Bukireva Street 15, Perm 614068, Russia.

✉ [sergei.esyunin@psu.ru](mailto:sergei.esyunin@psu.ru)

<https://orcid.org/0000-0003-3813-1316>

**Alireza Zamani**

Zoological Museum, Biodiversity Unit, FI-20014 University of Turku, Turku 20500, Finland.

✉ [zamani.alireza5@gmail.com](mailto:zamani.alireza5@gmail.com)

<https://orcid.org/0000-0002-8084-9666>

**Yuri M. Marusik**

Institute for Biological Problems of the North, FEB RAS, Portovaya Str. 18, Magadan 685000, Russia [1]. Altai State University, Lenina Pr., 61, Barnaul, RF-656049, Russia [2].

✉ [yurmar@mail.ru](mailto:yurmar@mail.ru)

<https://orcid.org/0000-0002-4499-5148>

**Received:**

27 July 2023

**Accepted:**

09 August 2023

**Published:**

16 August 2023

**Subject Editor:**

John T.D. Caleb

**ABSTRACT.** Two poorly known species of the zodariid spider genus *Zodariellum* Andreeva & Tystshenko, 1968 – *Z. spasskyi* (Charitonov, 1946) and *Z. zebra* (Charitonov, 1946) – are redescribed based on type specimens from eastern Uzbekistan. Additionally, a lectotype is designated for the former species.

**Key words:** Aranei, Central Asia, lectotype, Zodariinae, *Z. spasskyi*, *Z. zebra*

**Citation:** Esyunin, S.L., Zamani, A. & Marusik, Y.M. (2023) Redescriptions of two poorly known species of *Zodariellum* Andreeva & Tystshenko, 1968 (Araneae, Zodariidae) from Uzbekistan. *Journal of Insect Biodiversity and Systematics*, 9 (4), 703–709.

### INTRODUCTION

*Zodariellum* Andreeva & Tystshenko, 1968 is a genus of the spider family Zodariidae Thorell, 1881, currently comprising 21 species distributed from the Volga River's delta to eastern China (WSC, 2023). The genus has recently undergone significant taxonomic revisions by Zamani & Marusik (2021, 2022). The highest diversity of *Zodariellum* occurs in Central Asia, from where 15 species are currently known (WSC, 2023). Five species have been documented in Uzbekistan, two of which, *Z. spasskyi* (Charitonov, 1946) and *Z. zebra* (Charitonov, 1946), are poorly illustrated. Both species are known from female specimens collected in the same locality in Aman-Kutap in eastern Uzbekistan (Charitonov, 1946, 1969), with *Z. spasskyi* also known by a doubtful record from Kazakhstan (Logunov & Gromov, 2012). Recently, we had the opportunity to examine the type material of these two species, which are redescribed and illustrated in this paper.

### MATERIAL AND METHODS

Photographs of specimens and their epigynes were obtained using an Olympus OMD EM-10 digital camera, with a Panasonic Lumix H-H025 25 mm f/1.7 lens mounted on a Zeiss microscope. Digital images of different focal planes were stacked with Adobe Photoshop CS6. Leg segments were

**Corresponding author:** Zamani, A., ✉ [zamani.alireza5@gmail.com](mailto:zamani.alireza5@gmail.com)

**Copyright** © 2023, Esyunin et al. This is an open access article distributed under the terms of the Creative Commons NonCommercial Attribution License (CC BY NC 4.0), which permits Share - copy and redistribute the material in any medium or format, and Adapt - remix, transform, and build upon the material, under the Attribution-NonCommercial terms.

measured on the dorsal side. Measurements of legs are listed as: total length (femur, patella, tibia, metatarsus, tarsus). All measurements are given in millimetres. **Abbreviations:** ALE – anterior lateral eye, AME – anterior median eye, Fe – femur, LT – lectotype, PLE – posterior lateral eye, PLT – paralectotype, PME – posterior median eye. **Depository:** PSU – Perm State University, Perm, Russia (S.L. Esyunin).

## RESULTS

### *Taxonomic hierarchy*

**Class Arachnida Lamarck, 1801**

**Order Araneae Clerck, 1757**

**Family Zodariidae Thorell, 1881**

**Genus *Zodariellum* Andreeva & Tystshenko, 1968**

*Zodariellum* Andreeva & Tystshenko, 1968:688.

*Zodariellum*: Marusik & Koponen, 2001:40 (removed from the synonymy of *Zodarion*). – Zamani & Marusik, 2021:184. – Zamani & Marusik, 2022:162.

**Type species.** *Zodariellum surprisum* Andreeva & Tystshenko, 1968 from Tajikistan, by original designation.

**Diagnosis, description and composition.** See Zamani & Marusik (2022).

### *Zodariellum spasskyi* (Charitonov, 1946) (Figs 1–2, 4–5)

*Zodarion spasskyi* Charitonov, 1946:22, fig. 11 (♀).

*Zodarion spasskyi*: Charitonov, 1969:72.

*Zodariellum spasskyi*: Zamani & Marusik, 2022:175.


**Type material. Lectotype** ♀ (PSU, **designated here**), UZBEKISTAN: *Samarqand Region*: Urgut District, near 39°18'N 66°55'E, 1200–1900 m; labelled as “*Zodarion spasskyi* n.sp. D. Charitonov det., Aman-Kutap [mountain area], 40 km SE from Samarqand, 23.V.1942, D.M. Fedotov”. **Paralectotype:** 1♀ (PSU; epigyne mounted on a slide), collected with the lectotype.

**Diagnosis.** *Zodariellum spasskyi* differs from all its congeners by its unique abdominal pattern: light-coloured with a wide dark median band spanning from the anterior to the posterior part (*vs.* a longitudinal dark band absent in other species). The shape of its epigyne is very similar to that of *Z. nenilini* (Eskov, 1995) and is almost indistinguishable. Nevertheless, there are slight differences in the anterior hood (cf. Figs 4, 8).

**Description. Female.** Habitus as in Figs 1–2. Total length 4.85 (LT), 5.01 (PLT). Carapace length 2.25 (LT), 2.38 (PLT), width 1.05 (LT), 1.70 (PLT). Eye sizes (LT): ALE: 0.10, AME: 0.14, PLE: 0.10, PME: 0.08. Colouration (*sensu* Charitonov, 1946, 1969): body light yellow; ocular region black; sternum with narrow reddish brown margin; abdomen dorsally with a longitudinal dark brown band. Measurements of legs (LT): I: 7.08 (1.90, 0.68, 1.50, 1.93, 1.08), II: 7.18 (1.90, 0.75, 1.38, 2.05, 1.10), III: 7.23 (2.00, 0.75, 1.38, 2.13, 0.98), IV: 9.65 (2.63, 0.83, 2.25, 2.75, 1.20). Epigyne as in Figs 4–5; plate twice wider than long, with triangular anterior hood, its anterior margin forming an angle of 120°; receptacles oval, spaced by ca. 4 times their width; copulatory ducts long, twisted 5 times around their axes, converging and forming an angle of ca. 105°.

**Male.** Unknown.

**Distribution.** Confidently known only from the type locality in eastern Uzbekistan. Although Logunov & Gromov (2012) included this species in their list of spiders of Kazakhstan, the original record for *Z. spasskyi* in this country could not be located. According to Zamani & Marusik (2022), it is likely that this record belongs to a different species. Considering that the epigyne of *Z. spasskyi* is almost indistinguishable from that of *Z. nenilini*, it is possible that their record in fact refers to the latter species.



**Figures 1–3.** Habitus of *Zodariellum* species. **1.** *Z. spasskyi*, lectotype; **2.** *Z. spasskyi*, paralectotype, dorsal and lateral views; **3.** *Z. zebra*, dorsal view. Scale bars = 1.0 mm.

***Zodariellum zebra* (Charitonov, 1946) (Figs 3, 6–7)**

*Zodarion zebra* Charitonov, 1946:21, fig. 10 (♀).

*Zodarion zebra*: Charitonov, 1969:71.

*Zodariellum zebra*: Zamani & Marusik, 2022:176.

**Type material.** **Holotype** ♀ (PSU; epigyne mounted on a slide), UZBEKISTAN: *Samarqand Region*: Urgut District, near 39°18'N 66°55'E, 1200–1900 m; labelled as “*Zodarion zebra* n.sp., D. Charitonov det., Aman-Kutap [mountain area], 40 km SE from Samarqand, 21.V.1942, D.M. Fedotov”.

**Diagnosis.** The epigyne of *Z. zebra* is very similar to that of *Z. proszynskii* (Nenilin & Fet, 1895). However, it can be distinguished by its large and well-delimited bell-shaped atrium and wider coils of copulatory ducts. Additionally, the anterior part of the copulatory ducts are contiguous (cf. Figs 6, 9). Regarding abdominal patterning, *Z. zebra* is similar to *Z. testaceofasciatum* (Spassky, 1941). Nevertheless, it can be distinguished from the latter by its smaller size (carapace length of 1.05, *vs.* 1.7).

**Description.** *Female.* Habitus as in Fig. 3. Total length 2.45. Carapace length 1.05, width 0.43. Eye sizes: ALE: 0.07, AME: 0.08, PLE: 0.07, PME: 0.07. Colouration (*sensu* Charitonov, 1946, 1969): carapace yellow, with a brown transverse spot on border of cephalic and thoracic parts; eye field black; abdomen light yellow, with 3 transverse brown bands in anterior half and a herringbone-shaped spot in posterior half. Measurements of legs (unlisted segments missing): I: 2.55 (0.67, 0.27, 0.53, 0.60, 0.48), II: 2.56 (0.70, 0.31, 0.49, 0.63, 0.43), Fe III: 0.77, IV: 3.57 (1.01, 0.35, 0.74, 0.92, 0.55). Epigyne as in Figs 6–7; plate ca. 1.9 times wider than long; with large, bell-shaped fovea having well delimited margins, ca. as long as wide; receptacles round, spaced by ca. 4 diameters; copulatory ducts converging, touching anteriorly, and forming an angle of 100°; each copulatory duct forming 4 complete coils.


*Male.* Unknown.

**Comments.** According to Zamani & Marusik (2022), there is a possibility that *Z. zebra* is a junior synonym of *Z. testaceofasciatum*. The latter species is known solely by a female and a juvenile specimen from western Tajikistan, with no accompanying illustrations of the epigyne (Spassky, 1941). These two species exhibit a similar abdominal pattern, and their respective type localities are approximately 200 km apart. *Zodariellum turanicum* Zamani & Marusik, 2022 is another species with a similar abdominal pattern, known only by the male holotype. Its type locality is situated in Lebap, Turkmenistan, about 155 km away from the type locality of *Z. zebra* and approximately 300 km from the type locality of

*Z. testaceofasciatum*. Until the discovery of the currently unknown sexes of these species and a redescription of the female of *Z. testaceofasciatum* is provided, the three species can tentatively be differentiated from one another by their size and colouration (Table 1).

**Table 1.** Comparison of some of the somatic characters for three species of *Zodariellum*.

	<i>Z. zebra</i> (♀)	<i>Z. testaceofasciatum</i> (♀)	<i>Z. turanicum</i> (♂)
Length of carapace	1.05	1.7	1.6
Colouration - carapace	yellow	black-brown	yellow
Colouration - abdomen; dorsal, ventral	yellow, yellow	chestnut brown, fawn brown	dark marrow, light beige
Transverse stripes	wide	wide	narrow



**Figures 4–9.** Epigynes of *Zodariellum* species. 4–5. *Z. spasskyi* (paralectotype); 6–7. *Z. zebra*; 8. *Z. nenilini*; 9. *Z. proszynskii*, ventral (4, 6, 8, 9) and dorsal (5, 7) views. Fig. 8 reproduced from Zamani & Marusik (2022) with permission from the copyright holder; Fig. 9 reproduced from Zamani & Marusik (2021). Scale bars = 0.1 mm.

## DISCUSSION

Two species of *Zodariellum* were redescribed in this paper based on type specimens from eastern Uzbekistan. Among the 15 species of *Zodariellum* known from Central Asia, four species – *Z. spasskyi*, *Z. tadzhikum* (Andreeva & Tystshenko, 1968), *Z. testaceofasciatum*, and *Z. zebra* – are known only by females, while three species – *Z. surprisum*, *Z. turanicum*, and *Z. turkmenicum* Zamani & Marusik, 2022 – are known only by males. The illustrations provided in this work and previous studies (e.g., Zamani & Marusik, 2022) reveal that the shape of the epigyne is quite homogenous among different species of this genus, despite potentially distinct differences in colouration and male palp. Amongst the Central Asian species, the epigynes of most species are inadequately illustrated. As a result, it is likely that the discovery of previously unknown sexes and the redescriptions of poorly known species will lead to the synonymization of some of these species.

## AUTHOR'S CONTRIBUTION

The authors confirm their contribution in the paper as follows: SE & AZ: Conceptualization; SE: Microscopic photography; SE, AZ & YM: Writing – original draft; AZ: Writing – review and editing. The authors read and approved the final version of the manuscript.

## FUNDING

The research of YM was funded by a state assignment of the Ministry of Science and Higher Education of the Russian Federation (project FZMW-2023-0006 “Endemic, local and invasive arthropods (Arthropoda) of the mountains of South Siberia and Central Asia: a unique gene pool of a biodiversity hotspot”).

## AVAILABILITY OF DATA AND MATERIAL

The specimens listed in this study are deposited in the Arachnological Collection of the Perm State University, Russia and are available from the curator, upon request.

## ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Not applicable.

## CONSENT FOR PUBLICATION

Not applicable.

## CONFLICT OF INTERESTS

The authors declare that there is no conflict of interest regarding the publication of this paper.

## ACKNOWLEDGMENTS

SE is grateful to Gulli Sh. Farzalieva (Perm State University, Perm) for her assistance in preparing the photographs. Also, we thank the reviewers for their valuable comments and suggestions from which the manuscript benefited.

## REFERENCES

- Andreeva, E.M. & Tystshenko, V.P. (1968) Materials on the fauna of spiders (Aranei) of Tadzhikistan. II. Zodariidae. *Zoologicheskii Zhurnal*, 47, 684–689. [in Russian]
- Charitonov, D.E. (1946) New forms of spiders of the USSR. *Izvestiya Yestestvoenno-Nauchnogo Instituta pri Molotovskom Gosudarstvennom Universitete imeni M. Gor'kogo*, 12, 19–32. [in Russian]
- Charitonov, D.E. (1969) Material'y k faune paukov SSR. *Uchenye Zapiski Permskogo ordena Trudovogo Krasnogo Znameni Gosudarstvennogo Universiteta Imeni A. M. Gor'kogo*, 179, 59–133. [in Russian]
- Logunov, D.V. & Gromov, A.V. (2012) *Spiders of Kazakhstan*. Siri Scientific Press, Manchester. 232 p.
- Marusik, Y.M. & Koponen, S. (2001) Spiders of the family Zodariidae from Mongolia (Arachnida: Araneae). *Reichenbachia*, 34, 39–48.
- Spassky, S. (1941) Araneae palaearticae novae VI. *Folia Zoologica et Hydrobiologica, Rigā*, 11, 12–26.

- WSC (2023) *World Spider Catalog*. Version 24.0. Natural History Museum Bern, Bern. Available from: <https://wsc.nmbe.ch> [Accessed 20 July 2023]
- Zamani, A. & Marusik, Y.M. (2021) Revision of the spider family Zodariidae (Arachnida, Araneae) in Iran and Turkmenistan, with seventeen new species. *ZooKeys*, 1035, 145–193. <https://doi.org/10.3897/zookeys.1035.65767>
- Zamani, A. & Marusik, Y.M. (2022) New taxonomic considerations in *Zodariellum* Andreeva & Tyshchenko, 1968 (Araneae: Zodariidae), with notes on the presence of cymbial diverticulum in different zodariid genera. *Zootaxa*, 5178 (2), 161–177. <https://doi.org/10.11646/zootaxa.5178.2.3>

## توصیف مجدد دو گونه کم‌شناخته شده از جنس *Zodariellum* Andreeva & Tystshenko, 1968 (Araneae, Zodariidae) از ازبکستان

سرگی ال. اسیونین<sup>۱</sup>، علیرضا زمانی<sup>۲\*</sup> و یوری ام. ماروسیک<sup>۳،۴</sup>

۱ دانشگاه ایالتی پرم، پرم، روسیه.

۲ موزه جانورشناسی، واحد تنوع زیستی، دانشگاه تورکو، فنلاند.

۳ انستیتو مسائل زیستی شمالگان، شاخه خاور دور آکادمی علوم روسیه، روسیه.

۴ دانشگاه ایالتی آلتای، بارنائول، روسیه.

\* پست الکترونیک نویسنده مسئول مکاتبه: [zamani.alireza5@gmail.com](mailto:zamani.alireza5@gmail.com)

تاریخ دریافت: ۰۵ مرداد ۱۴۰۲ | تاریخ پذیرش: ۱۸ مرداد ۱۴۰۲ | تاریخ انتشار: ۲۵ مرداد ۱۴۰۲ |

**چکیده:** دو گونه کم شناخته شده از عنکبوت‌های خانواده *Zodariidae* متعلق به جنس *Zodariellum* *Z. zebra* (Charitonov, 1946) و *Z. spasskyi* (Charitonov, 1946) شامل Andreeva & Tystshenko, 1968 بر اساس نمونه‌های تایپ از شرق ازبکستان مجدداً توصیف شدند. همچنین، برای گونه نخست نمونه لکتوتایپ تعیین شد.

**واژگان کلیدی:** Aranei، آسیای مرکزی، لکتوتایپ، *Zodariinae*، *Z. zebra*، *Z. spasskyi*