



This is a self-archived – parallel-published version of an original article. This version may differ from the original in pagination and typographic details. When using please cite the original.

Gunasekar Saravanan, Sudharsanan Baskaran, Balasubramaniam Vanithakumari and Sahsene Altinkaya

Bi-Univalent Characteristics For a Particular Class of Bazilevic Functions Generated By Convolution Using Balancing Polynomials

2025

<https://doi.org/10.47013/18.3.5>

Publisher's PDF

Saravanan, G., et al. (2025) "Bi-Univalent Characteristics For a Particular Class of Bazilevic Functions Generated By Convolution Using Balancing Polynomials". Jordan Journal of Mathematics and Statistics, vol. 18, no. 3, pp. 381-392.

<https://doi.org/10.47013/18.3.5>

Bi-Univalent Characteristics For a Particular Class of Bazilevič Functions Generated By Convolution Using Balancing Polynomials

Gunasekar Saravanan¹, Sudharsanan Baskaran², Balasubramaniam Vanithakumari^{3,4}, Şahsene Altunkaya^{5,*}

- (1) PG and Research Department of Mathematics, Pachaiyappas College, Chennai-600030, Tamil Nadu, India,
- (2) Department of Mathematics, Dwaraka Doss Goverdhan Doss Vaishnav College (Autonomous), Chennai600106, Tamil Nadu, India.,
- (3) Department of Mathematics, Agurchand Manmull Jain College, Meenambakkam, Chennai600061, Tamil Nadu, India.,
- (4) Department of Mathematics, Amrita School of Engineering, Amrita Vishwa Vidyapeetham, Chennai-601103, Tamil Nadu, India ,
- (5) Department of Mathematics and Statistics, Faculty of Sciences, University of Turku, Turku, Finland.

Received: March 24, 2024

Accepted : Oct. 24, 2024

Abstract: Bi-univalent functions, which are analytic and univalent in the open unit disk, have been a significant area of research in complex analysis due to their rich geometric properties. This study introduces a novel subclass of bi-univalent functions, called the Bazilevič functions class, which is created via the convolution of Balancing polynomials. We calculated the bounds for the Fekete-Szegő inequality as well as the initial coefficients $|a_2|$ and $|a_3|$. We have also included several relevant corollaries.

Keywords: Analytic function; Bi-Univalent function; Bazilevič functions; Convolution; Balancing polynomials.

2010 Mathematics Subject Classification. 30C45.

1 Introduction and preliminaries

A function that is complex-valued and depends on one or more complex variables is considered analytic if it is differentiable at every point within its domain. Any such analytic function that is normalized can be represented as a series in the form

$$f(z) = z + \sum_{t=2}^{\infty} a_t z^t. \quad (1)$$

This series is in the complex variable z and converges within the set $\mathcal{U} = \{z : z \in \mathbb{C}, |z| < 1\}$. The set \mathcal{A} is composed of all such functions. Indicate by \mathcal{S} the subclass of \mathcal{A} consisting of univalent functions.

Let c be a Schwarz function defined as

$$c(z) = \sum_{n=1}^{\infty} c_n z^n \quad (c(0) = 0, |c(z)| < 1).$$

Consider two functions, f_1 and f_2 , which are part of the class \mathcal{A} . We say that f_1 is *subordinate* to f_2 if there exists a function $c(z)$ that is analytic in \mathcal{U} , and makes $f_1(z) = f_2(c(z))$ hold true. This relationship is denoted as $f_1(z) \prec f_2(z)$. Here, it is known that for the Schwarz function c , the coefficient inequality $|c_n| < 1$ holds [8], and specifically, we have

$$|c_1| \leq 1, \quad |c_n| \leq 1 - |c_1|^2 \quad (n \in \mathbb{N} \setminus \{1\}).$$

Comprehensive information on the subordination concept is available in monographs authored by Miller and Mocanu [18].

* Corresponding author e-mail: sahsenealtinkaya@gmail.com

The Hadamard product of $f(z)$ and $k(z)$, where $k(z)$ is an element of the set \mathcal{A} and

$$k(z) = z + \sum_{t=2}^{\infty} k_t z^t$$

is expressed by

$$(f * k)(z) = z + \sum_{t=2}^{\infty} a_t k_t z^t.$$

Consider f as a function in \mathcal{S} . The function $f(z)$ is classified as *bi-univalent* if its inverse, denoted by $f^{-1}(w)$, extends analytically to $|w| < 1$ within the w -plane. That is, according to the *Koebe-One Quarter Theorem* [8], it provides that the image of \mathcal{U} under every univalent function $f \in \mathcal{A}$ contains a disc of radius $1/4$. Thus, clearly, every such univalent function $f \in \mathcal{A}$ has an inverse f^{-1} satisfying $f^{-1}(f(z)) = z$ and $f(f^{-1}(w)) = w$ ($|w| < r_0(f)$, $r_0(f) \geq \frac{1}{4}$), where

$$f^{-1}(w) = w - a_2 w^2 + (2a_2^2 - a_3) w^3 - (5a_2^3 - 5a_2 a_3 + a_4) w^4 + \dots$$

We use σ to represent the collection of all such bi-univalent functions present in \mathcal{U} . Here, we present several examples of functions belonging to the class σ which have greatly renewed interest in the study of bi-univalent functions:

$$f(z) = \frac{z}{1-z}, \quad g(z) = -\log(1-z), \quad h(z) = \frac{1}{2} \log \frac{1+z}{1-z}$$

with their respective inverses

$$f^{-1}(w) = \frac{w}{1+w}, \quad g^{-1}(w) = \frac{e^w - 1}{e^w}, \quad h^{-1}(w) = \frac{e^{2w} - 1}{e^{2w} + 1}.$$

The concept was first introduced by Lewin [17] in 1967, who also provided an estimate for the second coefficient of these functions, stating that $|a_2| < 1.51$. This estimation was later refined by Brannan and Clunie [5], who proposed that $|a_2| \leq \sqrt{2}$. Over the years, the initial coefficients of bi-univalent functions have been extensively studied and estimated, contributing to a rich body of literature on the subject. Not much is known about the bounds on the general coefficient $|a_n|$ for $n \geq 4$. In the literature, there are only limited works determining the general coefficient bounds. Hence, the coefficient estimate problem for each of the coefficients $|a_n|$ ($n \in \mathbb{N} \setminus \{1, 2\}$; $\mathbb{N} = \{1, 2, 3, \dots\}$) is still an open problem.

Table 1: Summary of findings on coefficients a_n for functions in Σ .

Researchers	Findings	Estimates
Lewin 1967	For all functions in Σ	$ a_2 \leq 1.51$ [17]
Brannan and Clunie 1980	For all functions in Σ	$ a_2 \leq \sqrt{2}$ [5]
Netanyahu 1969	Proven maximum for $ a_2 $	$\max a_2 = \frac{4}{3}$ [19]
Tan 1984	The best estimate for $ a_2 $	$ a_2 \leq 1.485$ [29]
Various researchers	Estimates for Maclaurin coefficients $ a_2 $ and $ a_3 $	See [16, 24, 26, 27, 28, 31, 32, 33, 34, 35]
Open Problem	Estimating the coefficient $ a_n $	Unresolved [2, 6, 12, 25]

These studies not only enrich the theoretical foundation of complex analysis but also have practical implications in solving problems related to differential equations, approximation theory, and other applied mathematical disciplines.

The process of determining precise bounds for $|a_3 - \kappa a_2^2|$ within any compact function family is referred to as the *Fekete-Szegő* problem. Specifically, when $\kappa = 1$, the function represents the Schwarzian derivative. The Schwarzian derivative plays a significant role in the theory of geometric functions [9].

The literature is rich with various integer number sequences, including well-known ones like Fibonacci, Lucas, Pell, and more. Recently, a new sequence known as Balancing numbers was introduced by Behera and Panda [4]. Over the past quarter-century, this sequence has been extensively studied, revealing many of its unique properties. Research has been conducted, and generalizations have certainly been made. For those interested in delving deeper into Balancing numbers, comprehensive information can be found in the references [7, 10, 11, 13, 14, 15, 20, 21, 22].

An interesting extension of Balancing numbers is the concept of Balancing polynomials. These polynomials, defined and explored in [23], represent a natural progression from Balancing numbers and exhibit some fascinating characteristics.

Definition 1. The Balancing polynomials, for any x in the complex plane \mathbb{C} , are expressed by the recurrence relation:

$$\mathcal{B}_n(x) = 6x\mathcal{B}_{n-1}(x) - \mathcal{B}_{n-2}(x).$$

Here, $\mathcal{B}_0(x) = 0$ and $\mathcal{B}_1(x) = 1$ serve as the initial conditions.

Similar to other number polynomials, Balancing polynomials can also be derived from certain generating functions. Here is one such example:

Lemma 1([3]). The ordinary generating function for Balancing polynomials is expressed as follows:

$$\mathcal{B}(x, z) = \sum_{n=0}^{\infty} \mathcal{B}_n(x)z^n = \frac{z}{1 - 6xz + z^2}.$$

This paper contributes to the understanding of Balancing polynomials by offering insights into their structural properties and their implications across various values of x in the complex plane. We establish a novel class $\mathfrak{BM}_{\mu, \nu}^x(\mathfrak{f} * \mathfrak{k})$ and compute bounds for the initial coefficients $|a_2|$ and $|a_3|$ for functions belonging to this newly defined class. Additionally, we explore the Fekete-Szegő inequality, enhancing our comprehension of the analytic properties and bounds of Balancing polynomials and their applications in mathematical analysis.

2 Coefficient Bounds For The Function Class $\mathfrak{BM}_{\mu, \nu}^x(\mathfrak{f} * \mathfrak{k})$

In this section, we establish a novel class $\mathfrak{BM}_{\mu, \nu}^x(\mathfrak{f} * \mathfrak{k})$, and calculate the bounds for the initial coefficients $|a_2|$ and $|a_3|$ for functions that belong to $\mathfrak{BM}_{\mu, \nu}^x(\mathfrak{f} * \mathfrak{k})$.

Definition 2. Assume $\mathfrak{f}, \mathfrak{k} \in \sigma$, $\mu \in \mathbb{C} \setminus \{0\}$, and $\nu \geq 0$. A function $(\mathfrak{f} * \mathfrak{k})(z) \in \sigma$ is said to belong to the class $\mathfrak{BM}_{\mu, \nu}^x(\mathfrak{f} * \mathfrak{k})$ if it satisfies the following conditions:

$$(1 - \mu) \left(\frac{(\mathfrak{f} * \mathfrak{k})(z)}{z} \right)^\nu + \mu \left(\frac{z(\mathfrak{f} * \mathfrak{k})'(z)}{(\mathfrak{f} * \mathfrak{k})(z)} \right) \left(\frac{(\mathfrak{f} * \mathfrak{k})(z)}{z} \right)^\nu \prec \frac{\mathcal{B}(x, z)}{z} = I(x, z) \tag{2}$$

and

$$(1 - \mu) \left(\frac{(\mathfrak{f} * \mathfrak{k})^{-1}(w)}{w} \right)^\nu + \mu \left(\frac{w((\mathfrak{f} * \mathfrak{k})^{-1}(w))'}{(\mathfrak{f} * \mathfrak{k})^{-1}(w)} \right) \left(\frac{(\mathfrak{f} * \mathfrak{k})^{-1}(w)}{w} \right)^\nu \prec \frac{\mathcal{B}(x, w)}{w} = I(x, w). \tag{3}$$

Example 1. We show that the class $\mathfrak{BM}_{\mu, \nu}^x(\mathfrak{f} * \mathfrak{k})$ is non-empty. We consider the function

$$\mathfrak{f}(z) = z + \frac{a}{4}z^2, \quad z \in \mathcal{U},$$

where

$$a = \frac{\sqrt{2}|6x|\sqrt{|6x|}}{|k_2|\sqrt{|36x^2(\nu + 2\mu)(\nu + 1) - 2(36x^2 - 1)(\mu + \nu)^2|}}.$$

By some simple calculation, we can obtain its inverse as follows

$$\mathfrak{f}^{-1}(w) = \frac{2(-1 + \sqrt{1 + aw})}{a}, \quad w \in \mathcal{U}.$$

For $x = \frac{1}{2}$, $\mu = 0$, $\nu = 1$ and $\mathfrak{k}(z) = \frac{z}{1-z}$, we see that $\frac{(\mathfrak{f} * \mathfrak{k})(z)}{z} \prec I(\frac{1}{2}, z)$ and $\frac{(\mathfrak{f} * \mathfrak{k})^{-1}(w)}{w} \prec I(\frac{1}{2}, w)$. Figures 1 and 2 show that the transformation of \mathcal{U} under $\mathfrak{f}(z)$ and $\mathfrak{f}^{-1}(w)$, respectively.




Fig. 1: The transformation of \mathcal{U} under $f(z)$.




Fig. 2: The transformation of \mathcal{U} under $f^{-1}(w)$.

Remark. (i) For $\nu = 1$, we define the new class $\mathfrak{BM}_{\mu,1}^{\alpha}(f * k)$ consisting of functions $(f * k)(z) \in \sigma$ that satisfy

$$(1 - \mu) \left(\frac{(f * k)(z)}{z} \right) + \mu (f * k)'(z) \prec I(x, z) \quad (z \in \mathcal{U})$$

and

$$(1 - \mu) \left(\frac{(f * k)^{-1}(w)}{w} \right) + \mu ((f * k)^{-1}(w))' \prec I(x, w) \quad (w \in \mathcal{U}).$$

(ii) For $\mu = 1$, we get the new class $\mathfrak{M}_{1,\nu}^{\alpha}(f * k)$ as follows:

$$\left(\frac{z(f * k)'(z)}{(f * k)(z)} \right) \left(\frac{(f * k)(z)}{z} \right)^{\nu} \prec I(x, z) \quad (z \in \mathcal{U})$$

and

$$\left(\frac{w((f * k)^{-1}(w))'}{(f * k)^{-1}(w)} \right) \left(\frac{(f * k)^{-1}(w)}{w} \right)^v \prec I(x, w) \quad (w \in \mathcal{U}).$$

(iii) For $\mu = 1$ and $\nu = 1$, the new class $\mathfrak{BM}_{1,1}^x(f * k)$ is defined as follows:

$$(f * k)'(z) \prec I(x, z) \quad (z \in \mathcal{U})$$

and

$$((f * k)^{-1}(w))' \prec I(x, w) \quad (w \in \mathcal{U}).$$

(iv) For $\mu = 1$ and $\nu = 0$, the new class $\mathfrak{BM}_{1,0}^x(f * k)$ is defined as follows:

$$\frac{z(f * k)'(z)}{(f * k)(z)} \prec I(x, z) \quad (z \in \mathcal{U})$$

and

$$\frac{w((f * k)^{-1}(w))'}{(f * k)^{-1}(w)} \prec I(x, w) \quad (w \in \mathcal{U}).$$

Remark. (v) We define the new class $\mathfrak{BM}_{\mu,\nu}^x(f * \frac{z}{1-z})$ as follows:

$$(1 - \mu) \left(\frac{f(z)}{z} \right)^{\nu} + \mu \left(\frac{zf'(z)}{f(z)} \right) \left(\frac{f(z)}{z} \right)^{\nu} \prec I(x, z) \quad (z \in \mathcal{U})$$

and

$$(1 - \mu) \left(\frac{f^{-1}(w)}{w} \right)^{\nu} + \mu \left(\frac{w(f^{-1}(w))'}{f^{-1}(w)} \right) \left(\frac{f^{-1}(w)}{w} \right)^{\nu} \prec I(x, w) \quad (w \in \mathcal{U}).$$

(vi) For $\nu = 1$, the new class $\mathfrak{BM}_{\mu,1}^x(f * \frac{z}{1-z})$ is defined as follows:

$$(1 - \mu) \left(\frac{f(z)}{z} \right) + \mu f'(z) \prec I(x, z) \quad (z \in \mathcal{U})$$

and

$$(1 - \mu) \left(\frac{f^{-1}(w)}{w} \right) + \mu (f^{-1}(w))' \prec I(x, w) \quad (w \in \mathcal{U}).$$

(vii) For $\mu = 1$ and $\nu = 1$, the new class $\mathfrak{BM}_{1,1}^x(f * \frac{z}{1-z})$ is defined as follows:

$$f'(z) \prec I(x, z) \quad (z \in \mathcal{U})$$

and

$$(f^{-1}(w))' \prec I(x, w) \quad (w \in \mathcal{U}).$$

(viii) For $\mu = 1$ and $\nu = 0$, the new class $\mathfrak{BM}_{1,0}^x(f * \frac{z}{1-z})$ is defined as follows:

$$\frac{zf'(z)}{f(z)} \prec I(x, z) \quad (z \in \mathcal{U})$$

and

$$\frac{w(f^{-1}(w))'}{f^{-1}(w)} \prec I(x, w) \quad (w \in \mathcal{U}).$$

This class was introduced and studied by Aktaş and Karaman [1].

Theorem 1. Assume that f , as defined in (1), belongs to the class $\mathfrak{BM}_{\mu, \nu}^{\times}(f * k)$. Then

$$|a_2| \leq \frac{\sqrt{2}|6x|\sqrt{|6x|}}{|k_2|\sqrt{|36x^2(\nu + 2\mu)(\nu + 1) - 2(36x^2 - 1)(\mu + \nu)^2|}}$$

and

$$|a_3| \leq \frac{1}{|k_3|} \left(\left| \frac{6x}{\nu + 2\mu} \right| + \left| \frac{36x^2}{(\mu + \nu)^2} \right| \right).$$

Proof. If $(f * k) \in \mathfrak{BM}_{\mu, \nu}^{\times}(f * k)$, then from (2) and (3), there exist Schwarz functions $l(z)$ and $m(w)$, mapping \mathcal{U} to \mathcal{U} , such that

$$(1 - \mu) \left(\frac{(f * k)(z)}{z} \right)^{\nu} + \mu \left(\frac{z(f * k)'(z)}{(f * k)(z)} \right) \left(\frac{(f * k)(z)}{z} \right)^{\nu} = I(x, l(z))$$

and

$$(1 - \mu) \left(\frac{(f * k)^{-1}(w)}{w} \right)^{\nu} + \mu \left(\frac{w((f * k)^{-1}(w))'}{(f * k)^{-1}(w)} \right) \left(\frac{(f * k)^{-1}(w)}{w} \right)^{\nu} = I(x, m(w)).$$

The functions $l(z)$ and $m(w)$ are defined as follows:

$$l(z) = l_1z + l_2z^2 + l_3z^3 + \dots$$

and

$$m(w) = m_1w + m_2w^2 + m_3w^3 + \dots$$

These functions are analytic within \mathcal{U} , with $l(0)=0$ and $m(0) = 0$. Additionally, the absolute values of $l(z)$ and $m(w)$ are less than 1 for all $z, w \in \mathcal{U}$. It is important to note that if the following conditions are met:

$$|l(z)| = |l_1z + l_2z^2 + l_3z^3 + \dots| < 1 \quad (z \in \mathcal{U})$$

and

$$|m(w)| = |m_1w + m_2w^2 + m_3w^3 + \dots| < 1 \quad (w \in \mathcal{U}),$$

then, the absolute values of the coefficients l_t and m_t (for $t = 1, 2, 3, \dots$) are less than or equal to 1 [8]:

$$|l_t| \leq 1, \quad |m_t| \leq 1 \quad (t = 1, 2, 3, \dots).$$

Since

$$\begin{aligned} & (1 - \mu) \left(\frac{(f * k)(z)}{z} \right)^{\nu} + \mu \left(\frac{z((f * k)(z))'}{(f * k)(z)} \right) \left(\frac{(f * k)(z)}{z} \right)^{\nu} \\ &= 1 + (\nu + \mu)a_2k_2z + \left((\nu + 2\mu)a_3k_3 + \left(\frac{(\nu + 2\mu)(\nu - 1)}{2} \right) a_2^2k_2^2 \right) z^2 + \dots \end{aligned}$$

and

$$\begin{aligned} & (1 - \mu) \left(\frac{(f * k)^{-1}(w)}{w} \right)^{\nu} + \mu \left(\frac{w((f * k)^{-1}(w))'}{(f * k)^{-1}(w)} \right) \left(\frac{(f * k)^{-1}(w)}{w} \right)^{\nu} \\ &= 1 - (\nu + \mu)a_2k_2w + \left((\nu + 2\mu)(2a_2^2k_2^2 - a_3k_3) + \left(\frac{(\nu + 2\mu)(\nu - 1)}{2} \right) a_2^2k_2^2 \right) w^2 + \dots \end{aligned}$$

or, equivalently

$$(1 - \mu) \left(\frac{(f * k)(z)}{z} \right)^{\nu} + \mu \left(\frac{z((f * k)(z))'}{(f * k)(z)} \right) \left(\frac{(f * k)(z)}{z} \right)^{\nu} = \mathcal{B}_1(x)l_1z + (\mathcal{B}_1(x)l_2 + \mathcal{B}_2(x)l_1^2) z^2 + \dots$$

and

$$(1 - \mu) \left(\frac{(f * k)^{-1}(w)}{w} \right)^v + \mu \left(\frac{w((f * k)^{-1}(w))'}{(f * k)^{-1}(w)} \right) \left(\frac{(f * k)^{-1}(w)}{w} \right)^v = \mathcal{B}_1(x)m_1w + (\mathcal{B}_1(x)m_2 + \mathcal{B}_2(x)m_1^2)w^2 + \dots,$$

we get following equations

$$(v + \mu)a_2k_2 = \mathcal{B}_2(x)l_1, \tag{4}$$

$$(v + 2\mu)a_3k_3 + \frac{(v + 2\mu)(v - 1)}{2}a_2^2k_2^2 = \mathcal{B}_2(x)l_2 + \mathcal{B}_3(x)l_1^2, \tag{5}$$

$$-(v + \mu)a_2k_2 = \mathcal{B}_2(x)m_1 \tag{6}$$

and

$$(v + 2\mu)(2a_2^2k_2^2 - a_3k_3) + \frac{(v + 2\mu)(v - 1)}{2}a_2^2k_2^2 = \mathcal{B}_2(x)m_2 + \mathcal{B}_3(x)m_1^2. \tag{7}$$

By adding equations (4) and (6), we obtain the following result

$$l_1 = -m_1. \tag{8}$$

Additionally, squaring and adding (4) and (6), we derive

$$2(\mu + v)^2k_2^2a_2^2 = \mathcal{B}_2^2(x)(l_1^2 + m_1^2). \tag{9}$$

The addition of (5) and (7) yields

$$(v + 2\mu)(v + 1)a_2^2k_2^2 = \mathcal{B}_2(x)(l_2 + m_2) + \mathcal{B}_3(x)(l_1^2 + m_1^2).$$

From these equations, we can deduce

$$[\mathcal{B}_2^2(x)(v + 2\mu)(v + 1) - 2\mathcal{B}_3(x)(\mu + v)^2] a_2^2k_2^2 = \mathcal{B}_2^3(x)(l_2 + m_2). \tag{10}$$

A minor computation leads to the following upper bound for $|a_2|$:

$$|a_2| \leq \frac{\sqrt{2}|\mathcal{B}_2(x)|^{\frac{3}{2}}}{|k_2| \sqrt{|\mathcal{B}_2^2(x)(v + 2\mu)(v + 1) - 2\mathcal{B}_3(x)(\mu + v)^2|}}.$$

To find the bound for $|a_3|$, we subtract equation (7) from (5), resulting in

$$2a_3k_3(v + 2\mu) - 2(v + 2\mu)a_2^2k_2^2 = \mathcal{B}_2(x)(l_2 - m_2) + \mathcal{B}_3(x)(l_1^2 - m_1^2). \tag{11}$$

By substituting equations (8) and (9) into (11), we obtain

$$a_3k_3 = \frac{\mathcal{B}_2(x)(l_2 - m_2)}{2(v + 2\mu)} + \frac{\mathcal{B}_2^2(x)(l_1^2 + m_1^2)}{2(\mu + v)^2}.$$

After applying the value of $\mathcal{B}_2(x)$ and taking the modulus, we get the desired bound for $|a_3|$:

$$|a_3| \leq \frac{1}{|k_3|} \left(\left| \frac{6x}{v + 2\mu} \right| + \left| \frac{36x^2}{(\mu + v)^2} \right| \right).$$

Corollary 1. If f , given by (1), is in $\mathfrak{BM}_{\mu,1}^x(f * k)$, then

$$|a_2| \leq \frac{|6x|\sqrt{|6x|}}{|k_2| \sqrt{|(36x^2 - 1)\mu^2 - (2\mu + 1)|}}$$

and

$$|a_3| \leq \frac{1}{|k_3|} \left(\left| \frac{6x}{1 + 2\mu} \right| + \left| \frac{36x^2}{(1 + \mu)^2} \right| \right).$$

Corollary 2. If f , given by (1), is in $\mathfrak{BM}_{1,v}^{\alpha}(f * k)$, then

$$|a_2| \leq \frac{|6x|\sqrt{|6x|}}{|k_2|\sqrt{|(18x^2-1)v^2+(9x^2-1)2v-1|}}$$

and

$$|a_3| \leq \frac{1}{|k_3|} \left(\left| \frac{6x}{v+2} \right| + \left| \frac{36x^2}{(1+v)^2} \right| \right).$$

Corollary 3. If f , given by (1), is in $\mathfrak{BM}_{1,1}^{\alpha}(f * k)$, then

$$|a_2| \leq \frac{|3x|\sqrt{|6x|}}{|k_2|\sqrt{|9x^2-1|}}$$

and

$$|a_3| \leq \frac{1}{|k_3|} (2|x| + 9|x^2|).$$

Corollary 4. If f , given by (1), is in $\mathfrak{BM}_{1,0}^{\alpha}(f * k)$, then

$$|a_2| \leq \frac{1}{|k_2|} (|6x|\sqrt{|6x|})$$

and

$$|a_3| \leq \frac{1}{|k_3|} (3|x| + 36|x^2|).$$

Corollary 5. If f , given by (1), is in $\mathfrak{BM}_{\mu,v}^{\alpha}(f * \frac{z}{1-z})$, then

$$|a_2| \leq \frac{\sqrt{2}|6x|\sqrt{|6x|}}{\sqrt{|36x^2(v+2\mu)(v+1)-2(36x^2-1)(\mu+v)^2|}}$$

and

$$|a_3| \leq \left| \frac{6x}{v+2\mu} \right| + \left| \frac{36x^2}{(\mu+v)^2} \right|.$$

Corollary 6. If f , given by (1), is in $\mathfrak{BM}_{\mu,1}^{\alpha}(f * \frac{z}{1-z})$, then

$$|a_2| \leq \frac{|6x|\sqrt{|6x|}}{\sqrt{|(36x^2-1)\mu^2-(2\mu+1)|}}$$

and

$$|a_3| \leq \left| \frac{6x}{1+2\mu} \right| + \left| \frac{36x^2}{(1+\mu)^2} \right|.$$

Corollary 7. If f , given by (1), is in $\mathfrak{BM}_{1,v}^{\alpha}(f * \frac{z}{1-z})$, then

$$|a_2| \leq \frac{|6x|\sqrt{|6x|}}{\sqrt{|(18x^2-1)v^2+(9x^2-1)2v-1|}}$$

and

$$|a_3| \leq \left| \frac{6x}{v+2} \right| + \left| \frac{36x^2}{(1+v)^2} \right|.$$

Corollary 8. If f , given by (1), is in $\mathfrak{BM}_{1,0}^{\alpha}(f * \frac{z}{1-z})$, then

$$|a_2| \leq |6x|\sqrt{|6x|}$$

and

$$|a_3| \leq 3|x| + 36|x^2|.$$

3 Fekete-Szegő Inequalities For The Function Class $\mathfrak{BM}_{\mu, \nu}^{\kappa}(\mathfrak{f} * \mathfrak{k})$

In this section, we will be determining the Fekete-Szegő inequality, denoted as $|a_3 - \kappa a_2^2|$, where $\kappa \in \mathbb{R}$, for functions that belong to the class $\mathfrak{BM}_{\mu, \nu}^{\kappa}(\mathfrak{f} * \mathfrak{k})$.

Theorem 2. *If the function \mathfrak{f} , as defined in (1), belongs to the class $\mathfrak{BM}_{\mu, \nu}^{\kappa}(\mathfrak{f} * \mathfrak{k})$, then the following inequality holds*

$$|a_3 - \kappa a_2^2| \leq \begin{cases} \left| \frac{6x}{\nu+2\mu} \right|, & |\Psi(\mu, \nu, \mathfrak{k}, x)| \leq \frac{1}{2|\mathfrak{k}_3(\nu+2\mu)|} \\ |12x\Psi(\mu, \nu, \mathfrak{k}, x)|, & |\Psi(\mu, \nu, \mathfrak{k}, x)| \geq \frac{1}{2|\mathfrak{k}_3(\nu+2\mu)|} \end{cases},$$

where

$$\Psi(\mu, \nu, \mathfrak{k}; x) = \frac{36x^2 \left(\frac{\mathfrak{k}_2^2}{\mathfrak{k}_3} - \kappa \right)}{\mathfrak{k}_2^2(36x^2(\nu+2\mu)(\nu+1) - 2(36x^2-1)(\mu+\nu)^2)}.$$

Proof. For $\kappa \in \mathbb{R}$, it follows from (8) and (11) that

$$a_3 - \kappa a_2^2 = \frac{1}{\mathfrak{k}_3} \frac{\mathcal{B}_2(x)(l_2 - m_2)}{2(\nu+2\mu)} + \left(\frac{\mathfrak{k}_2^2}{\mathfrak{k}_3} - \kappa \right) a_2^2.$$

By using (10) in the above equality, we obtain

$$\begin{aligned} a_3 - \kappa a_2^2 &= \mathcal{B}_2(x) \left[\frac{l_2 - m_2}{2\mathfrak{k}_3(\nu+2\mu)} + \left(\frac{\mathfrak{k}_2^2}{\mathfrak{k}_3} - \kappa \right) \frac{\mathcal{B}_2^2(x)(l_2 + m_2)}{\mathfrak{k}_2^2(\mathcal{B}_2^2(x)(\nu+2\mu)(\mu+1) - 2\mathcal{B}_3(x)(\mu+\nu)^2)} \right] \\ &= \mathcal{B}_2(x) \left[\left(\frac{1}{2\mathfrak{k}_3(\nu+2\mu)} + \Psi(\mu, \nu, \mathfrak{k}; x) \right) l_2 + \left(\frac{-1}{2\mathfrak{k}_3(\nu+2\mu)} + \Psi(\mu, \nu, \mathfrak{k}; x) \right) m_2 \right], \end{aligned}$$

where

$$\Psi(\mu, \nu, \mathfrak{k}; x) = \frac{\left(\frac{\mathfrak{k}_2^2}{\mathfrak{k}_3} - \kappa \right) \mathcal{B}_2^2(x)}{\mathfrak{k}_2^2(\mathcal{B}_2^2(x)(\nu+2\mu)(\nu+1) - 2\mathcal{B}_3(x)(\mu+\nu)^2)}.$$

Therefore, we have

$$|a_3 - \kappa a_2^2| \leq \begin{cases} \left| \frac{6x}{\nu+2\mu} \right|, & |\Psi(\mu, \nu, \mathfrak{k}, x)| \leq \frac{1}{2|\mathfrak{k}_3(\nu+2\mu)|} \\ |12x\Psi(\mu, \nu, \mathfrak{k}, x)|, & |\Psi(\mu, \nu, \mathfrak{k}, x)| \geq \frac{1}{2|\mathfrak{k}_3(\nu+2\mu)|} \end{cases}.$$

Corollary 9. *If \mathfrak{f} , given by (1), is in $\mathfrak{BM}_{\mu, 1}^{\kappa}(\mathfrak{f} * \mathfrak{k})$ and $\kappa \in \mathbb{R}$. Then*

$$|a_3 - \kappa a_2^2| \leq \begin{cases} \left| \frac{6x}{1+2\mu} \right|, & |\Psi(\mu, 1, \mathfrak{k}; x)| \leq \frac{1}{2|\mathfrak{k}_3(1+2\mu)|} \\ |12x\Psi(\mu, 1, \mathfrak{k}, x)|, & |\Psi(\mu, 1, \mathfrak{k}; x)| \geq \frac{1}{2|\mathfrak{k}_3(1+2\mu)|} \end{cases},$$

where

$$\Psi(\mu, 1, \mathfrak{k}; x) = \frac{36x^2 \left(\frac{\mathfrak{k}_2^2}{\mathfrak{k}_3} - \kappa \right)}{\mathfrak{k}_2^2(36x^2(1+2\mu)(\nu+1) - 2(36x^2-1)(\mu+1)^2)}.$$

Corollary 10. *If \mathfrak{f} , given by (1), is in $\mathfrak{BM}_{1, \nu}^{\kappa}(\mathfrak{f} * \mathfrak{k})$ and $\kappa \in \mathbb{R}$, then*

$$|a_3 - \kappa a_2^2| \leq \begin{cases} \left| \frac{6x}{\nu+2} \right|, & |\Psi(1, \nu, \mathfrak{k}, x)| \leq \frac{1}{2|\mathfrak{k}_3(\nu+2)|} \\ |12x\Psi(1, \nu, \mathfrak{k}, x)|, & |\Psi(1, \nu, \mathfrak{k}, x)| \geq \frac{1}{2|\mathfrak{k}_3(\nu+2)|} \end{cases},$$

where

$$\Psi(1, \nu, \mathfrak{k}; x) = \frac{36x^2 \left(\frac{\mathfrak{k}_2^2}{\mathfrak{k}_3} - \kappa \right)}{\mathfrak{k}_2^2(36x^2(\nu+2)(\nu+1) - 2(36x^2-1)(1+\nu)^2)}.$$

Corollary 11. If f , given by (1), is in $\mathfrak{BM}_{1,1}^{\kappa}(f * k)$ and $\kappa \in \mathbb{R}$, then

$$|a_3 - \kappa a_2^2| \leq \begin{cases} 2|x|, & |\Psi(1, 1, k, x)| \leq \frac{1}{6|k_3|} \\ |12x\Psi(1, 1, k, x)|, & |\Psi(1, 1, k, x)| \geq \frac{1}{6|k_3|} \end{cases},$$

where

$$\Psi(1, 1, k; x) = \frac{9x^2\left(\frac{k_2^2}{k_3} - \kappa\right)}{2k_2^2(1 - 9x^2)}.$$

Corollary 12. If f , given by (1), is in $\mathfrak{BM}_{1,0}^{\kappa}(f * k)$ and $\kappa \in \mathbb{R}$, then

$$|a_3 - \kappa a_2^2| \leq \begin{cases} 3|x|, & |\Psi(1, 0, k, x)| \leq \frac{1}{4|k_3|} \\ |12x\Psi(1, 0, k, x)|, & |\Psi(1, 0, k, x)| \geq \frac{1}{4|k_3|} \end{cases},$$

where

$$\Psi(1, 0, k; x) = \frac{18x^2\left(\frac{k_2^2}{k_3} - \kappa\right)}{k_2^2}.$$

Corollary 13. If f , given by (1), is in $\mathfrak{BM}_{\mu, \nu}^{\kappa}(f * \frac{z}{1-z})$ and $\kappa \in \mathbb{R}$, then

$$|a_3 - \kappa a_2^2| \leq \begin{cases} \left| \frac{6x}{v+2\mu} \right|, & |\Psi(\mu, \nu, x)| \leq \frac{1}{2|v+2\mu|} \\ |12x\Psi(\mu, \nu, x)|, & |\Psi(\mu, \nu, x)| \geq \frac{1}{2|v+2\mu|} \end{cases},$$

where

$$\Psi(\mu, \nu; x) = \frac{36x^2(1 - \kappa)}{36x^2(v + 2\mu)(v + 1) - 2(36x^2 - 1)(\mu + \nu)^2}.$$

Corollary 14. If f , given by (1), is in $\mathfrak{BM}_{\mu, 1}^{\kappa}(f * \frac{z}{1-z})$ and $\kappa \in \mathbb{R}$, then

$$|a_3 - \kappa a_2^2| \leq \begin{cases} \left| \frac{6x}{1+2\mu} \right|, & |\Psi(\mu, 1, x)| \leq \frac{1}{2|1+2\mu|} \\ |12x\Psi(\mu, 1, x)|, & |\Psi(\mu, 1, x)| \geq \frac{1}{2|1+2\mu|} \end{cases},$$

where

$$\Psi(\mu, 1, x) = \frac{36x^2(1 - \kappa)}{36x^2(1 + 2\mu)(v + 1) - 2(36x^2 - 1)(\mu + 1)^2}.$$

Corollary 15. If f , given by (1), is in $\mathfrak{BM}_{1, \nu}^{\kappa}(f * \frac{z}{1-z})$ and $\kappa \in \mathbb{R}$, then

$$|a_3 - \kappa a_2^2| \leq \begin{cases} \left| \frac{6x}{v+2} \right|, & |\Psi(1, \nu, x)| \leq \frac{1}{2|v+2|} \\ |12x\Psi(1, \nu, x)|, & |\Psi(1, \nu, x)| \geq \frac{1}{2|v+2|} \end{cases},$$

where

$$\Psi(1, \nu, k; x) = \frac{36x^2(1 - \kappa)}{36x^2(v + 2)(v + 1) - 2(36x^2 - 1)(1 + \nu)^2}.$$

Corollary 16. If f , given by (1), is in $\mathcal{BM}_{1,0}^{\kappa}(f * k)$ and $\kappa \in \mathbb{R}$, then

$$|a_3 - \kappa a_2^2| \leq \begin{cases} 3|x|, & |\Psi(1, 0, x)| \leq \frac{1}{4} \\ |12x\Psi(1, 0, x)|, & |\Psi(1, 0, x)| \geq \frac{1}{4} \end{cases},$$

where

$$\Psi(1, 0, x) = 18x^2(1 - \kappa).$$

4 Conclusion

In this study, we have introduced and thoroughly investigated a novel subclass of bi-univalent functions known as Bazilevič functions, which are constructed through the convolution of Balancing polynomials. Our primary objective was to establish rigorous bounds for the Fekete-Szegő inequality and to compute the initial coefficients $|a_2|$ and $|a_3|$ for functions belonging to this subclass.

Furthermore, the corollaries derived from our analysis offer additional insights and extensions, underscoring the versatility and applicability of Bazilevič functions in mathematical research. These results pave the way for further exploration into the deeper implications and potential applications of Bazilevič functions across different parameters.

In conclusion, our study contributes to the advancement of mathematical analysis by establishing a solid theoretical foundation for Bazilevič functions and by providing concrete computational results. Future research can build upon these findings to explore more intricate aspects of Bazilevič functions and to investigate their role in solving complex mathematical problems.

Declarations

Competing interests: The authors declare that they have no competing interests.

Authors' contributions: Conceptualization, S.G., B.S., B.V and S.A.; methodology, S.G., B.S., B.V and S.A.; software, S.G., B.S and S.A.; validation, S.G., B.S., B.V and S.A.; formal analysis, S.G., B.S., B.V and S.A.; investigation, S.G., B.S. and S.A.; resources, S.G., B.S., B.V and S.A.; writing original draft preparation, S.G., B.S., B.V and S.A.; writing review and editing, S.G., B.S., B.V and S.A.; visualization, S.G., B.S., B.V and S.A.; supervision, S.A.; All authors have read and agreed to the published version of the manuscript.

Funding: No funding was received for this study.

Availability of data and materials: No data or materials were used in this study. All relevant information is included in the manuscript.

References

- [1] I. Aktas, I. Karaman, On some new subclasses of bi-univalent functions defined by Balancing polynomials, *KMU Journal of Engineering and Natural Sciences*, **5(1)**(2023), 25-32.
- [2] Ş. Altınkaya, S. Yalçın, Faber polynomial coefficient bounds for a subclass of bi-univalent functions, *Comptes Rendus. Mathématique*, **353(12)**(2015), 1075-1080.
- [3] S. Baskaran, G. Saravanan, B. Vanithakumari, Sakaguchi type function defined by (p, q) -fractional operator using Laguerre polynomials, *Palestine Journal of Mathematics*, **11** (2022), 41-47.
- [4] A. Behera, G. K. Panda, On the square roots of triangular numbers, *Fibonacci Quart.*, **37**(1999), 98-105.
- [5] D. A. Brannan, J. G. Clunie (Eds.), *Aspects of contemporary complex analysis*, (Proceedings of the NATO Advanced Study Institute held at the University of Durham, July 1-20, 1979), Academic Press, New York and London, (1980).
- [6] S. Bulut, Faber polynomial coefficient estimates for a comprehensive subclass of analytic bi-univalent functions, *Comptes Rendus. Mathématique*, **352(12)**(2014), 479-484.
- [7] R. K. Davala, G. K. Panda, On sum and ratio formulas for balancing numbers, *J. Indian Math. Soc.*, **82(1-2)**(2015), 23-32.
- [8] P. L. Duren, *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften, Springer, New York, 1983.
- [9] M. Fekete, G. Szegő, Eine Bemerkung über ungerade schlichte Funktionen, *J. London Math. Soc.*, **s1-8(2)**(1933), 85-89.
- [10] R. Frontczak, L. Baden Württemberg, A note on hybrid convolutions involving balancing and Lucas-balancing numbers, *Appl. Math. Sci.*, **12(25)**(2018), 1201-1208.

- [11] R. Frontczak, L. Baden Württemberg, Sums of balancing and Lucas-balancing numbers with binomial coefficients, *Int. J. Math. Anal.*, **12(12)**(2018), 585-594.
- [12] S. G. Hamidi, J. M. Jahangiri, Faber polynomial coefficient estimates for analytic bi-close-to-convex functions, *Comptes Rendus. Mathematique*, **352**(2014), 17-20.
- [13] A. Hussien, M. S. A. Madi, A. M. M. Abominjil, Bounding coefficients for certain subclasses of bi-univalent functions related to Lucas-Balancing polynomials, *AIMS Mathematics* **9(7)**(2024), 1803418047.
- [14] R. Keskin, O. Karaatli, Some new properties of balancing numbers and square triangular numbers, *J. Integer Seq.*, **15(1)**(2012), 1-13.
- [15] T. Komatsu, G. K. Panda, On several kinds of sums of balancing numbers, arXiv:1608.05918, (2016), 1-21.
- [16] V. Madaana, A. Kumar, V. Ravichandran, Estimates for initial coefficients of certain bi-univalent functions, *Filomat*, **35(6)**(2021), 19932009.
- [17] M. Lewin, On a coefficient problem for bi-univalent functions, *Proc. Amer. Math. Soc.*, **18(1)**(1967), 63-68.
- [18] S. S. Miller, P. T. Mocanu, *Differential Subordinations, Monographs and Textbooks in Pure and Applied Mathematics*, 225, Marcel Dekker, Inc., New York, (2000).
- [19] E. Netanyahu, The minimal distance of the image boundary from the origin and the second coefficient of a univalent function in $|z| < 1$, *Arc. for rational mechanics and analysis*, **32(2)**(1969), 100-112.
- [20] B. K. Patel, N. Irmak, P. K. Ray, Incomplete balancing and Lucas-balancing numbers, *Math. Rep.*, **20(70)**(2018), 59-72.
- [21] P. K. Ray, Some congruences for Balancing and Lucas-Balancing numbers and their applications, *Integers*, **14**(2014), 1-8.
- [22] P. K. Ray, Balancing and Lucas-balancing sums by matrix methods, *Math. Rep.*, **17(2)**(2015), 225-233.
- [23] P. K. Ray, On the properties of k -balancing numbers, *Ain Shams Engineering Journal*, **9(3)**(2018), 395-402.
- [24] G. Saravanan, S. Baskaran, B. Vanithakumari, S. Bulut, Sakaguchi type functions defined by balancing polynomials. *Mathematica Bohemica*, first online, pp. 1-13, DOI: 10.21136/MB.2024.0173-23.
- [25] G. Saravanan, K. Muthunagai, Coefficient estimates for the class of Bi-Quasi-Convex functions using Faber polynomials, *Far East J. Math. Sci.*, **102(10)**(2017), 2267-2276.
- [26] H. M. Srivastava, A. K. Mishra, P. Gochhayat, Certain subclasses of analytic and bi-univalent functions, *Appl. Math. Lett.* **23(10)**(2010), 11881192.
- [27] H. M. Srivastava, A. K. Wanas, H. Ö. Güney, New families of bi-univalent functions associated with the Bazilevic functions and the k -Pseudo-starlike functions, *Iran. J. Sci. Technol. Trans. Sci.*, **45**(2021), 17991804.
- [28] H. M. Srivastava, S. S. Eker, R. M. Ali, Coefficient bounds for a certain class of analytic and bi-univalent functions, *Filomat*, **29(8)**(2015), 18391845.
- [29] D. L. Tan, Coefficient estimates for bi-univalent functions, *Chinese Annals of Mathematics, Series A*, **5**(1984), 559-568.
- [30] D. K. Thomas, N. Tuneski, A. Vasudevarao, *Univalent Functions: A Primer*, Berlin, Boston: De Gruyter, (2018).
- [31] B. Vanithakumari, G. Saravanan, B. Baskaran and Yalin, S., 2024. Horadam Polynomials For A New Subclass Of Sakaguchi-Type Bi-Univalent Functions Defined by (p,q) -derivative operator, *Commun. Korean Math. Soc.*, **39(2)**, **2024**, 461-470.
- [32] R. Vijaya, T. V. Sudharsan, S. Sivasubramanian, Coefficient estimates for certain subclasses of bi-univalent functions defined by convolution, *International Journal of Analysis*, **2016**(2016), 1-5.
- [33] Q. H. Xu, Y. C. Gui, H. M. Srivastava, Coefficient estimates for a certain subclass of analytic and bi-univalent functions, *Appl. Math. Lett.*, **25(6)**(2012), 990-994.
- [34] Q. H. Xu, H. G. Xiao, H. M. Srivastava, A certain general subclass of analytic and bi-univalent functions and associated coefficient estimate problems, *Applied Mathematics and Computation*, **218(23)**(2012), 11461-11465.
- [35] S. Yalçın, K. Muthunagai, G. Saravanan, A subclass with bi-univalence involving (p,q) -Lucas polynomials and its coefficient bounds, *Bol. Soc. Mat. Mex.*, **26**(2020), 1015-1022.